

**SOUTH DAKOTA STATEWIDE FISHERIES SURVEY**  
**Burke, Gregory County**  
**FTR-Lake-3197-000**  
**2018**

**Lake Information**

<b>Name:</b>	Burke	<b>Maximum Depth:</b>	16 Feet
<b>County:</b>	Gregory	<b>Mean Depth:</b>	9 Feet
<b>Legal Description:</b>	T97-R71-S32		
<b>Surface Area:</b>	29 Acres		

**Surveys and Investigations**

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
boat shocker (night)	Oct 16, 2018	3600 seconds
boat shocker (night)	Oct 31, 2018	3600 seconds

## **Common Fish Species Present**

Largemouth Bass

Bluegill

Black Crappie

---

## Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- **AFS std gill net** – 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- **std experimental gill net for non-Missouri River waters** - 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- **std experimental gill net for Missouri River reservoirs** – 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$CPUE = \frac{\text{number of fish}}{\text{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left( \frac{\text{number of fish} \geq \text{quality length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

$$PSD - P = \left( \frac{\text{number of fish} \geq \text{preferred length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (**W**) compared to a length-specific standard weight (**Ws**) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left( \frac{W}{W_s} \right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (S), quality (Q), preferred (P), memorable (M) and trophy (T). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

## **Catch Summary of Stock Length Fish**

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

\* **Methods/Species that ignore stock length**

Gear	Species	Sample Size (n)	Abundance		Stock Density Indices			Condition		
			CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
boat shocker (night)	Largemouth Bass	60	13.0	3.5	77	13	46	15	112	2

## **10-Year Catch Per Unit Effort by Gear and Species**

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

Gear	Species	CPUE										Avg
		2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	
AFS std frame net	Black Bullhead									0.1		0.1
	Black Crappie									10.9		10.9
	Bluegill									11.4		11.4
	Largemouth Bass									0.1		0.1
	Northern Pike									0.1		0.1
	Yellow Perch									0.6		0.6
boat shocker (night)	Largemouth Bass			52.5			18.0		35.5	33.0	13.0	30.4
frame net (std 3/4 in)	Black Bullhead			12.4			0.8					6.6
	Black Crappie			10.4			3.9					7.2
	Bluegill			12.6			6.9					9.8
	Green Sunfish			1.4			0.1					0.8
	Northern Pike			0.4			0.2					0.3
	Yellow Perch			0.3			2.0					1.2

## 10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

Gear	Species	Index	Year									
			2009	2010	2011	2012	2013	2014	2015	2016	2017	2018
AFS std frame net	Black Crappie	PSD										51
		PSD-P										0
		Wr										105
	Bluegill	PSD										83
		PSD-P										5
		Wr										107
	Largemouth Bass	PSD										0
		PSD-P										0
		Wr										93
boat shocker (night)	Largemouth Bass	PSD			66			87		34	27	77
		PSD-P			23			60		30	12	46
		Wr			107			118		106	103	112
frame net (std 3/4 in)	Black Crappie	PSD			64			72				
		PSD-P			0			8				
		Wr			107			104				
	Bluegill	PSD			75			62				
		PSD-P			2			12				
		Wr			115			112				

## **Back-Calculated Lengths**

Mean species back-calculated total length (mm) at age, standard error (SE), and sample size (N).

Species: Largemouth Bass

Year Class	Age	N	Mean back-calculated length (SE) at age																	
			1	2	3	4	5	6	7	8	9	10								
2017	1	6	98 (8.7)																	
2016	2	2	86 (9.6)	173 (8.2)																
2015	3	8	81 (5.4)	165 (9.7)	247 (10.2)															
2014	4	6	84 (3.7)	179 (13.7)	262 (13.6)	326 (7.4)														
2013	5	2	108 (17.7)	198 (18.1)	312 (.1)	373 (11.4)	406 (7.5)													
2012	6	3	100 (13.1)	195 (13.5)	344 (16.6)	413 (17.3)	451 (14.9)	471 (11.9)												
2011	7	2	96 (9.7)	204 (21.3)	324 (74.7)	418 (69.2)	468 (58.5)	501 (42)	515 (35.8)											
2010	8	1	93	200	280	366	422	453	492	518										
Weighted Mean		30	90	180	279	367	440	478	507	518										
Year Class	Age	N	11	12	13	14	15	16	17	18	19	20								
2017	1	6																		
2016	2	2																		
2015	3	8																		
2014	4	6																		
2013	5	2																		
2012	6	3																		
2011	7	2																		
2010	8	1																		
Weighted Mean		30																		



## Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Black Crappie

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2017	109				199 (109)						
2014	75	110 (39)	165 (1)	202 (23)	239 (6)	249 (5)	243 (1)				
2011	82		153 (7)	194 (37)	217 (24)	221 (11)	233 (2)				

Species: Bluegill

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2017	114	128 (8)	134 (6)	168 (41)	181 (53)	199 (6)					
2014	70	74 (1)	113 (20)	146 (14)	185 (5)	192 (13)	193 (13)	202 (3)			
2011	101			141 (39)	166 (42)		188 (4)	187 (9)	195 (5)		193 (2)

Species: Largemouth Bass

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2018	39	135 (14)	241 (2)	283 (8)	366 (6)	430 (2)	488 (4)	525 (2)	527 (1)		
2014	54	164 (40)	285 (1)	351 (4)	407 (1)	409 (3)	433 (2)	423 (1)		454 (1)	496 (1)
2011	35	200 (1)	268 (8)	310 (13)	351 (5)	402 (3)	409 (3)	460 (1)	424 (1)		

## **Fish Condition**


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

Species	Year	Length Groups							
		S-Q		Q-P		P-M		M	
		N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Black Crappie Frame Net	2014	11	110 (2.8)	25	102 (1.1)	3	99 (1.2)	0	
	2017	53	107 (0.9)	56	104 (0.8)	0		0	
Bluegill Frame Net	2014	26	117 (9.1)	35	108 (1.9)	8	110 (2.0)	0	
	2017	19	115 (2.6)	89	106 (1.4)	6	100 (3.4)	0	
Largemouth Bass Electro Fishing	2014	2	115 (6.1)	4	119 (3.4)	9	117 (3.7)	0	
	2016	47	106 (1.1)	3	107 (5.6)	18	109 (2.8)	3	104 (0.4)
	2017	24	102 (1.8)	5	106 (5.9)	3	110 (1.2)	1	98
	2018	6	110 (2.8)	8	111 (2.7)	9	115 (3.0)	3	109 (2.7)


## Length Frequency Distribution

Length frequency histogram of species sampled by year.


Species: Black Crappie  
Gear: AFS std frame net




Species: Black Crappie  
Gear: frame net (std 3/4 in)







Species: Bluegill  
Gear: AFS std frame net



Species: Bluegill  
Gear: frame net (std 3/4 in)




Species: Largemouth Bass  
Gear: boat shocker (night)




## Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.


Species: Black Crappie  
Gear: AFS std frame net




Species: Black Crappie  
Gear: frame net (std 3/4 in)



Species: Bluegill  
Gear: AFS std frame net



Species: Bluegill  
Gear: frame net (std 3/4 in)



Species: Largemouth Bass  
Gear: boat shocker (night)

