SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Fort Meade BLM Pond, Meade County LBF-Lake-43-000 2018

Lake Information

Name: Fort Meade BLM Pond

County: Meade

Surface Area: 12 Acres

Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort	
AFS std gill net	May 15, 2018	2 net-nights	
frame net (1/4 inch)	May 15, 2018	1 net-nights	
frame net (std 3/4 in)	May 15, 2018	3 net-nights	
rod and reel	May 15, 2018	3 minutes	

Common Fish Species Present

Largemouth Bass

Yellow Perch

Rainbow Trout

Black Bullhead

Northern Pike

Black Crappie

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- AFS std gill net 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- std experimental gill net for non-Missouri River waters 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- std experimental gill net for Missouri River reservoirs 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$\mathit{CPUE} = \frac{\mathit{number of fish}}{\mathit{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{number\ of\ fish \ge quality\ length}{number\ of\ fish \ge stock\ length}\right) \times 100$$

$$\textit{PSD} - \textit{P} = \left(\frac{number\ of\ fish\ \geq preferred\ length}{number\ of\ fish\ \geq stock\ length}\right) \ge 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{Ws}\right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (S), quality (Q), preferred (P), memorable (M) and trophy (T). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

	St	ock	Qu	ality	Preferred		Memorable		Trophy	
Species Name	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

* Methods/Species that ignore stock length

			Abundance		St	ock Der	Condition			
Gear	Species	Sample Size (n)	CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
AFS std gill net	Black Bullhead	2	1.0	3.1	100		100		117	13
	Northern Pike	1	0.5	1.5	100		0		85	
	Yellow Perch	1	0.5	1.5	0		0		100	
frame net (1/4	Black Bullhead	2	2.0		100		0		85	5
inch)	Largemouth Bass	1	1.0		100		0		103	
frame net (std 3/4	Black Bullhead	96	32.0	30.3	88	5	0		82	1
in)	Black Crappie	1	0.3	0.6	0		0		71	

10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

							CPUE					
Gear	Species	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	Avg
AFS std gill net	Black Bullhead										1.0	1.0
	Northern Pike										0.5	0.5
	Yellow Perch										0.5	0.5
frame net (1/4	Black Bullhead										2.0	2.0
inch)	Largemouth Bass										1.0	1.0
frame net (std	Black Bullhead						0.3				32.0	16.2
3/4 in)	Black Crappie										0.3	0.3
	Green Sunfish						3.0					3.0
	White Sucker						0.3					0.3
	Yellow Perch						67.0					67.0

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

		Year										
Gear	Species	Index	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018
AFS std gill net	Black Bullhead	PSD										100
		PSD-P										100
		Wr										117
	Northern Pike	PSD										100
		PSD-P										0
		Wr										85
	Yellow Perch	PSD										0
		PSD-P										0
		Wr										100
frame net (1/4 inch)	Black Bullhead	PSD										100
		PSD-P										0
		Wr										85
	Largemouth Bass	PSD										100
		PSD-P										0
		Wr										103
frame net (std	Black Bullhead	PSD						100				88
3/4 in)		PSD-P						0				0
		Wr						83				82
	Black Crappie	PSD										0
		PSD-P										0
		Wr										71
	Yellow Perch	PSD						36				
		PSD-P						0				
		Wr						87				

Fish Condition


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

		Length Groups							
			S-Q		Q-P		P-M		M
Species	Year	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Black Bullhead Gill Net	2018	0		0		2	117 (10.2)	0	
Black Crappie Frame Net	2018	1	71	0		0		0	
Northern Pike Gill Net	2018	0		1	85	0		0	
Yellow Perch Gill Net	2018	1	100	0		0		0	


Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.


Species: Black Bullhead Gear: AFS std gill net




Species: Black Crappie Gear: frame net (std 3/4 in)



Species: Northern Pike Gear: AFS std gill net



Species: Yellow Perch Gear: AFS std gill net



Fish Stocking

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2008	Largemouth Bass	Adult	80
2008	Largemouth Bass	Fingerling	1,000
2009	Largemouth Bass	Fingerling	2,000
2010	Largemouth Bass	Fingerling	1,000
2011	Largemouth Bass	Fingerling	7,000
2014	Largemouth Bass	Fingerling	400
2015	Largemouth Bass	Fingerling	2,000
2017	Largemouth Bass	Fingerling	1,000