

SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Owen Dam, Perkins County

LMO-Lake-397-000

2018

Lake Information

Name: Owen Dam

County: Perkins

Surface Area: 136 Acres

Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
AFS std gill net	Jul 10, 2018	1 net-nights
AFS std gill net	Jul 11, 2018	1 net-nights
frame net (std 3/4 in)	Jul 10, 2018	3 net-nights
frame net (std 3/4 in)	Jul 11, 2018	3 net-nights

Common Fish Species Present

Yellow Perch

Northern Pike

Largemouth Bass

Black Bullhead

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- **AFS std gill net** – 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- **std experimental gill net for non-Missouri River waters** - 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- **std experimental gill net for Missouri River reservoirs** – 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$CPUE = \frac{\text{number of fish}}{\text{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{\text{number of fish} \geq \text{quality length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

$$PSD - P = \left(\frac{\text{number of fish} \geq \text{preferred length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (*W*) compared to a length-specific standard weight (*Ws*) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{W_s} \right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (S), quality (Q), preferred (P), memorable (M) and trophy (T). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

*** Methods/Species that ignore stock length**

Gear	Species	Sample Size (n)	Abundance		Stock Density Indices			Condition		
			CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
frame net (std 3/4 in)	Black Bullhead	24	4.0	1.3	46	16	42	16	108	2

10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

Gear	Species	CPUE										Avg
		2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	
AFS std frame net	Black Bullhead									1.7		1.7
	Black Crappie									0.2		0.2
	Bluegill									0.1		0.1
	Largemouth Bass									0.1		0.1
	Northern Pike									0.4		0.4
AFS std gill net	Black Bullhead									6.3		6.3
	Northern Pike									2.8		2.8
frame net (std 3/4 in)	Black Bullhead		1.5		65.3	3.5	7.8	67.3	4.4		4.0	22.0
	Black Crappie						0.3	0.1	0.2			0.2
	Bluegill						1.6	20.5	0.2			7.4
	Green Sunfish							0.1				0.1
	Largemouth Bass								0.2			0.2
	Northern Pike		10.8		4.8	1.5	4.5	1.4	1.8			4.1
	Yellow Perch		0.3						0.1			0.2
std exp gill net	Black Bullhead				91.0		1.5	2.5	1.0			24.0
	Green Sunfish								0.5			0.5
	Northern Pike		8.0		8.0	9.0	7.5	7.5	4.0			7.3
	Yellow Perch		2.0				1.5					1.8

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

Gear	Species	Index	Year										
			2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	
AFS std frame net	Black Bullhead	PSD											100
		PSD-P											100
		Wr											125
	Largemouth Bass	PSD											100
		PSD-P											100
		Wr											123
	Northern Pike	PSD											75
		PSD-P											25
		Wr											89
AFS std gill net	Black Bullhead	PSD										100	
		PSD-P										100	
		Wr										137	
	Northern Pike	PSD										82	
		PSD-P										18	
		Wr										93	
frame net (std 3/4 in)	Black Bullhead	PSD		100		37	100	27	17	50		46	
		PSD-P		17		2	29	6	1	0		42	
		Wr		107		107	114	93	106	101		108	
	Largemouth Bass	PSD									100		
		PSD-P									100		
		Wr									127		
	Northern Pike	PSD		0		89	83	64	55	67			
		PSD-P		0		5	33	3	0	33			
		Wr		88		94	93	87	94	92			
	Yellow Perch	PSD		0						100			
		PSD-P		0						100			
		Wr		103									
std exp gill net	Black Bullhead	PSD				21		33	20	0			
		PSD-P				0		0	0	0			
		Wr				110		130	114	106			
	Northern Pike	PSD		0		75	100	33	73	88			
		PSD-P		0		0	11	0	20	0			
		Wr											

Gear	Species	Index	Year									
			2009	2010	2011	2012	2013	2014	2015	2016	2017	2018
std exp gill net	Northern Pike	Wr		92		98	96	86	96	95		
	Yellow Perch	PSD		0				33				
		PSD-P		0				0				
		Wr		93				82				

Fish Condition


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

Species	Year	Length Groups							
		S-Q		Q-P		P-M		M	
		N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Black Bullhead Gill Net	2014	4	135 (6.6)	2	122 (0.0)	0		0	
	2015	8	112 (1.7)	2	120 (0.0)	0		0	
	2016	4	106 (2.3)	0		0		0	
	2017	0		0		25	137 (2.4)	0	
Northern Pike Gill Net	2014	20	85 (1.0)	10	89 (1.3)	0		0	
	2015	8	96 (3.6)	16	96 (1.6)	2	89 (0.0)	4	101 (1.9)
	2016	2	81 (0.0)	14	97 (2.3)	0		0	
	2017	2	100 (1.7)	7	90 (3.3)	2	93 (0.2)	0	
Yellow Perch Gill Net	2014	4	72 (2.8)	2	103 (0.0)	0		0	


Length Frequency Distribution

Length frequency histogram of species sampled by year.


Species: Black Bullhead
Gear: AFS std gill net




Species: Black Bullhead
Gear: std exp gill net






Species: Northern Pike
Gear: AFS std gill net



Species: Northern Pike
Gear: std exp gill net







Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.


Species: Black Bullhead
Gear: AFS std gill net




Species: Black Bullhead
Gear: std exp gill net




Species: Northern Pike
Gear: AFS std gill net



Species: Northern Pike
Gear: std exp gill net



Species: Yellow Perch
Gear: std exp gill net



Fish Stocking

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2009	Yellow Perch	Adult	250
2010	Largemouth Bass	Fingerling	5,560
2010	Northern Pike	Fry	97,600
2012	Bluegill	Adult	835
2012	Largemouth Bass	Adult	320
2012	Yellow Perch	Adult	572
2014	Largemouth Bass	Juvenile	250
2014	Yellow Perch	Adult	800
2017	Black Crappie	Adult	200
2017	Bluegill	Adult	25
2017	Yellow Perch	Adult	25
2018	Bluegill	Adult	200
2018	Northern Pike	Adult	15
