SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Brush, Brookings County MBS-Lake-253-800 2018

Lake Information

Name: Brush

County: Brookings

Surface Area: 553 Acres

Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
AFS std gill net	Jul 02, 2018	6 net-nights

Common Fish Species Present

Yellow Perch

Walleye

Black Bullhead

Northern Pike

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- AFS std gill net 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- std experimental gill net for non-Missouri River waters 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- std experimental gill net for Missouri River reservoirs 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$\mathit{CPUE} = \frac{\mathit{number of fish}}{\mathit{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{number\ of\ fish \ge quality\ length}{number\ of\ fish \ge stock\ length}\right) \times 100$$

$$\textit{PSD} - \textit{P} = \left(\frac{number\ of\ fish\ \geq preferred\ length}{number\ of\ fish\ \geq stock\ length}\right) \ge 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{Ws}\right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (S), quality (Q), preferred (P), memorable (M) and trophy (T). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

	St	ock	Qu	ality	Preferred		Mem	orable	Tro	pphy
Species Name	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

* Methods/Species that ignore stock length

				dance	St	ock Der	sity Indic	ity Indices		ndition
Gear	Species	Sample Size (n)	CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
AFS std gill net	Black Bullhead	70	11.7	2.4	59	9	13	6		
	Northern Pike	1	0.2	0.2	0		0		106	
	Walleye	82	13.7	1.6	54	8	12	5	97	1
	Yellow Perch	95	15.8	5.3	51	7	34	7	105	1

10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

							CPUE					
Gear	Species	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	Avg
AFS std gill ne	et Black Bullhead									8.5	11.7	10.1
	Northern Pike									1.2	0.2	0.7
	Walleye									23.7	13.7	18.7
	White Sucker									0.2		0.2
	Yellow Perch									2.2	15.8	9.0

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

							Ye	ar				
Gear	Species	Index	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018
AFS std gill net	Black Bullhead	PSD					,		,	,	49	59
		PSD-P									4	13
	Northern Pike	PSD									100	0
		PSD-P									29	0
		Wr									91	106
	Walleye	PSD									78	54
		PSD-P									3	12
		Wr									94	97
	Yellow Perch	PSD									62	51
		PSD-P									8	34
		Wr									96	105

Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Walleye

				Mean Ler	igth (expar	nded sam	ple numb	er) at capt	ure by ag	е	
Year	N	1	2	3	4	5	6	7	8	9	10+
2018	82	287 (37)	410 (12)	447 (12)	502 (13)			533 (1)	562 (2)	540 (1)	652 (3)

Fish Condition


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

					Length	Group	S		
			S-Q		Q-P	P-M			M
Species	Year	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Northern Pike Gill Net	2017	0		5	91 (4.2)	2	90 (6.4)	0	
	2018	1	106	0		0		0	
Walleye Gill Net	2017	31	92 (1.3)	107	95 (0.8)	4	84 (2.4)	0	
	2018	38	95 (0.9)	34	100 (1.0)	7	96 (2.3)	3	85 (4.1)
Yellow Perch Gill Net	2017	5	100 (2.1)	7	92 (2.8)	1	101	0	
	2018	47	108 (1.2)	16	102 (1.6)	28	102 (1.1)	4	98 (2.6)


Length Frequency Distribution

Length frequency histogram of species sampled by year.


Species: Black Bullhead Gear: AFS std gill net



Species: Walleye Gear: AFS std gill net




Species: Yellow Perch Gear: AFS std gill net




Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.


Species: Black Bullhead Gear: AFS std gill net




Species: Northern Pike Gear: AFS std gill net



Species: Walleye Gear: AFS std gill net



Species: Yellow Perch Gear: AFS std gill net



Fish Stocking

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2009	Walleye	Fingerling	8,181
2009	Yellow Perch	Fingerling	244,339
2010	Walleye	Small Fingerling	39,550
2011	Yellow Perch	Small Fingerling	206,640
2012	Black Crappie	Juvenile	770
2012	Yellow Perch	Fingerling	165,360
2014	Walleye	Fingerling	48,603
2015	Walleye	Small Fingerling	28,160
2017	Walleye	Small Fingerling	27,550
2018	Yellow Perch	Small Fingerling	199,710