

Waubay Lake Survey Summary

Waubay Lake, located on the southeastern edge of Grenville, is managed as a walleye and yellow perch fishery but other fish species (e.g., smallmouth bass, white bass) also contribute to the fishery.

- **Smallmouth bass.** More smallmouth bass were sampled by day electrofishing in 2019 (16.0/hour) than combined day and night electrofishing in 2015 (8.0/hour). In 2019, sampled fish ranged in length from 7.5 to 16.5 inches, 4 of the 16 fish caught were ≥ 14.0 inches.
- **Walleye.** Gill net CPUE's have remained relatively stable in recent years (2016 – 2019). In 2019, relative abundance was considered moderate to high at 7.9/gill net. Sampled walleyes ranged in length from 7.9 to 28.0 inches, of those that were at least 10.0 inches 72% were 15.0 inches or longer. Year classes produced in 2011 and 2016, both of which coincided with fry stockings, were the most abundant and accounted for more than 70% of walleyes sampled. Growth of the 2011 cohort was slow with a mean length at capture of 12.0 inches at age 3. Meanwhile, growth of the 2016 year class has been faster to age 3 with a mean length at capture of 15.6 inches.
- **White bass.** White bass have been among the most abundant fish species in the gill net catch since 2016. In 2019, the mean gill net CPUE was 7.3 and relative abundance was considered low to moderate. Those sampled ranged in length from 7.5 to 16.5 inches, most (88%) were ≥ 12.0 inches and 53% were 15.0 inches or longer.
- **Yellow perch.** Yellow perch numbers were higher in 2019 than 2018. Relative abundance was low to moderate at 10.3/gill net. Sampled yellow perch ranged in length from 5.1 to 12.6 inches, 12% were ≥ 8.0 inches and 9% were 10.0 inches or longer. Individuals from seven year classes produced between 2009 and 2018 contributed to the catch, those from the 2018 (age-1) cohort were the most abundant accounting for 88% of fish in the sample. Yellow perch growth appears to be good with mean length at capture values at age 3 exceeding 9.0 inches from 2010 to 2019. In 2019, the mean length at capture of age-3 fish was 9.7 inches.

For more detailed results see the computer generated South Dakota Statewide Fisheries Survey for Waubay (Day; below).

Common Fish Species Present

Northern Pike

Smallmouth Bass

Walleye

Yellow Perch

Emerald Shiner

White Bass

Black Crappie

Bluegill

Common Carp

Rock Bass

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- **AFS std gill net** – 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- **std experimental gill net for non-Missouri River waters** - 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- **std experimental gill net for Missouri River reservoirs** – 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$CPUE = \frac{\text{number of fish}}{\text{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{\text{number of fish} \geq \text{quality length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

$$PSD - P = \left(\frac{\text{number of fish} \geq \text{preferred length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (*W*) compared to a length-specific standard weight (*Ws*) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{W_s} \right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (S), quality (Q), preferred (P), memorable (M) and trophy (T). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

* Methods/Species that ignore stock length

Gear	Species	Sample Size (n)*	Abundance		Stock Density Indices			Condition		
			CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
AFS std gill net	Black Bullhead	6	0.4	0.2	83		50		95	7
	Bluegill	1	0.1	0.1	0		0		133	
	Common Carp	16	1.0	0.3	81		75		98	3
	Rock Bass	15	0.9	0.4	71		7		111	2
	Smallmouth Bass	12	0.6	0.3	50	28	30		99	4
	Walleye	130	7.9	1.2	72	6	3		88	1
	White Bass	116	7.3	2.1	89	4	88	4	95	1
	Yellow Perch	165	10.3	2.0	12	4	9	3	111	1
boat shocker (day)	Smallmouth Bass	16	16.0	6.3	63	20	25		96	2
fall night EF-WAE*	Walleye	7	10.5	8.4						

10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

*Include night sampling completed prior to 2018; ** Methods/Species that ignore stock length

Gear	Species	CPUE										Avg
		2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	
AFS std gill net	Black Bullhead							0.4	0.3	0.3	0.4	0.4
	Bluegill							0.0	0.1	0.0	0.1	0.1
	Common Carp							0.1	0.5	0.3	1.0	0.5
	Lake Herring							0.3	0.1	0.1	0.0	0.1
	Northern Pike							0.1	0.1	0.0	0.0	0.1
	Rock Bass							0.4	0.8	0.3	0.9	0.6
	Smallmouth Bass							1.3	1.3	0.9	0.6	1.0
	Walleye							6.3	4.6	5.9	7.9	6.2
	White Bass							13.2	12.9	6.9	7.3	10.1
	White Sucker							0.0	0.1	0.0	0.0	0.0
Yellow Perch							5.4	8.3	6.4	10.3	7.6	
boat shocker (day)*	Smallmouth Bass				62.8		8.0				16.0	28.9
fall night EF-WAE	Walleye		6.0	5.0	1.0	15.0	1.2	1.5	7.0	0.0	10.5	5.2
frame net (std 3/4 in)	Black Bullhead	0.8	0.4	1.5	3.5	2.0						1.6
	Black Crappie	0.2	0.3	1.3	1.5	2.6						1.2
	Bluegill	0.6	0.7	0.9	0.4	0.3						0.6
	Common Carp	0.5	0.5	0.5	0.3	0.2						0.4
	Northern Pike	0.4	0.1	0.2	0.3	0.6						0.3
	Rock Bass	0.9	0.6	0.9	2.6	1.2						1.2
	Smallmouth Bass	6.3	6.1	5.1	6.2	3.5						5.4
	Walleye	5.5	3.1	2.9	2.5	2.8						3.4
	White Bass	3.1	6.5	5.1	3.8	2.5						4.2
	White Sucker	0.1	0.1	0.1	0.1	0.0						0.1
Yellow Perch	0.2	0.0	0.1	0.0	0.0						0.1	
std exp gill net	Black Bullhead	0.0	0.2	4.3	4.1	1.4	0.1					1.7
	Bluegill	0.0	0.0	0.3	0.0	0.0	0.0					0.1
	Common Carp	0.3	0.0	0.5	0.0	0.5	0.1					0.2
	Lake Herring	1.4	0.3	0.1	0.4	0.3	0.3					0.5
	Northern Pike	0.0	0.0	0.1	0.5	1.0	0.4					0.3
	Rock Bass	0.1	0.2	1.4	1.3	2.0	0.4					0.9
	Smallmouth Bass	0.0	0.0	0.0	0.3	0.3	0.0					0.1
	Spottail Shiner**	0.1	0.0	0.0	0.0	0.0	0.0					0.0
	Walleye	5.3	5.3	11.1	11.8	19.3	14.1					11.2
	White Bass	0.3	1.0	1.5	17.6	8.1	23.9					8.7
White Sucker	0.1	0.0	0.0	0.3	0.1	0.0					0.1	
Yellow Perch	6.6	9.2	28.1	21.9	18.5	19.5					17.3	

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

*Includes night sampling completed prior to 2018

Gear	Species	Index	Year										
			2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	
AFS std gill net	Walleye	PSD							28	81	52	72	
		PSD-P							5	3	1	3	
		Wr							86	88	89	88	
	White Bass	PSD								100	99	100	89
		PSD-P								100	99	98	88
		Wr								98	92	98	95
	Yellow Perch	PSD								71	62	88	12
		PSD-P								38	37	34	9
		Wr								109	109	109	111
boat shocker (day)*	Smallmouth Bass	PSD				52			88			63	
		PSD-P				21			38			25	
		Wr				89			85			96	
std exp gill net	Walleye	PSD	25	42	48	28	17	8					
		PSD-P	0	0	7	2	1	0					
		Wr	88	83	83	82	84	85					
	White Bass	PSD	100	83	100	100	100	100					
		PSD-P	100	22	75	99	100	99					
		Wr	101	98	97	93	98	97					
	Yellow Perch	PSD	59	72	85	79	87	83					
		PSD-P	22	22	32	36	41	38					
		Wr	115	114	117	115	115	117					

Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Smallmouth Bass

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2019	12		206 (2)	270 (4)	288 (4)	318 (2)					
2015	8			260 (1)	294 (1)	331 (5)		414 (1)			
2013	63		208 (5)	253 (29)	312 (15)	369 (8)	418 (1)	441 (4)	443 (1)		

Species: Walleye

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2019	130	253 (10)	341 (13)	395 (72)		460 (6)	466 (1)	553 (1)	467 (24)		548 (3)
2018	96	264 (7)	327 (37)	430 (7)	402 (8)	444 (2)	472 (3)	434 (33)			
2017	92	223 (21)	354 (1)	394 (4)	479 (2)		410 (58)		410 (2)	445 (1)	668 (2)
2016	100		320 (6)	396 (1)	396 (1)	365 (83)	334 (1)	485 (4)			691 (3)
2015	117	215 (4)	280 (1)		332 (104)		387 (4)				417 (4)
2014	157	228 (3)		304 (120)		386 (21)	399 (3)			435 (9)	
2013	113	235 (4)	259 (61)	350 (2)	374 (28)	427 (2)		374 (3)	447 (11)	416 (1)	724 (1)
2012	217	213 (131)	326 (14)	367 (30)	402 (7)	489 (1)	365 (2)	447 (30)			502 (1)
2011	129	248 (5)	322 (50)	368 (17)			407 (54)	391 (3)	454 (1)		
2010	129	270 (47)	347 (24)			380 (55)	395 (3)				

Species: Yellow Perch

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2019	165	146 (145)		247 (9)	282 (6)	274 (1)	306 (2)	311 (1)			327 (1)
2018	102		211 (46)	242 (29)	257 (7)	281 (3)		296 (10)	274 (4)	295 (1)	
2017	133	152 (39)	207 (31)	247 (21)	269 (7)	273 (4)	279 (18)	300 (4)	286 (7)		
2016	87	153 (22)	205 (7)	238 (18)	258 (16)	267 (21)	242 (4)				
2015	159	136 (17)	190 (14)	237 (38)	248 (56)	261 (14)	267 (19)	312 (1)			
2014	152	138 (9)	186 (12)	233 (62)	261 (33)	251 (31)	299 (3)	343 (1)	303 (1)		
2013	175	150 (5)	198 (51)	237 (39)	250 (62)	244 (4)	280 (10)		323 (1)	269 (4)	
2012	228	144 (26)	217 (63)	244 (111)	264 (13)	261 (7)			305 (1)	335 (1)	270 (7)
2011	221	153 (23)	206 (131)	242 (26)	270 (32)	280 (9)					
2010	159	151 (60)	217 (31)	245 (61)	250 (7)		278 (3)				

Fish Condition


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

Species	Year	Length Groups							
		S-Q		Q-P		P-M		M	
		N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Smallmouth Bass Electro Fishing	2015	1	76	4	86 (2.4)	3	87 (5.8)	0	
	2019	6	98 (3.2)	6	95 (3.9)	4	95 (0.9)	0	
Walleye Gill Net	2015	104	86 (0.4)	9	81 (1.1)	0		0	
	2016	72	87 (0.5)	23	86 (0.9)	2	80 (2.5)	3	83 (2.5)
	2017	14	88 (1.5)	57	88 (0.7)	0		2	89 (3.6)
	2018	46	88 (0.8)	48	89 (0.9)	1	87	0	
	2019	36	88 (0.7)	87	87 (0.5)	3	86 (1.5)	1	90
White Bass Gill Net	2015	0		1	265	182	96 (0.4)	8	84 (2.1)
	2016	0		0		169	99 (0.4)	42	94 (0.9)
	2017	2	97 (2.5)	0		120	94 (0.4)	84	90 (0.5)
	2018	0		2	97 (1.1)	39	98 (1.0)	69	99 (0.7)
	2019	13	100 (1.3)	1	105	41	96 (0.5)	61	94 (0.6)
Yellow Perch Gill Net	2015	26	114 (2.0)	71	120 (1.1)	57	116 (1.4)	2	112 (2.6)
	2016	25	107 (1.7)	29	111 (1.7)	33	109 (1.9)	0	
	2017	51	109 (1.4)	33	113 (1.6)	39	111 (1.3)	10	96 (3.0)
	2018	12	112 (2.9)	55	111 (1.0)	32	106 (1.6)	3	91 (3.0)
	2019	145	112 (0.7)	5	115 (3.8)	10	116 (3.1)	5	97 (3.9)

Length Frequency Distribution


Length frequency histogram of species sampled by year.

Species: Smallmouth Bass
Gear: boat shocker (day)




2019


Species: Walleye
Gear: AFS std gill net




2016





2017





2018




Species: Walleye
 Gear: std exp gill net




Species: White Bass
 Gear: AFS std gill net






2018




2019

Species: White Bass
Gear: std exp gill net




2014




2015


Species: Yellow Perch
Gear: AFS std gill net




2016



2017




2018




2019

Species: Yellow Perch
Gear: std exp gill net



2014




2015


Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.


Species: Smallmouth Bass
Gear: boat shocker (day)




Species: Smallmouth Bass
Gear: boat shocker (night, DC)




Species: Walleye
Gear: AFS std gill net




Species: Walleye
Gear: std exp gill net




Species: White Bass
Gear: AFS std gill net




Species: White Bass
Gear: std exp gill net



Species: Yellow Perch
Gear: AFS std gill net



Species: Yellow Perch
Gear: std exp gill net



Fish Stocking

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2009	Walleye	Fry	4,000,000
2011	Walleye	Fry	8,000,000
2012	Walleye	Fry	8,000,000
2014	Walleye	Fry	8,500,000
2016	Walleye	Fry	8,500,000
2017	Walleye	Fry	8,000,000
2019	Walleye	Fry	4,000,000