

**Note:** Curlyleaf pondweed is present in Roy Lake. Care should be taken by all user groups to prevent the spread to other waters. For more information regarding this and other aquatic invasive species please visit <https://sdleastwanted.sd.gov/>

### **Roy Lake Survey Summary**

Roy Lake, located 2.0 miles south and 1.0 miles west of Lake City, is managed as a multiple species fishery including panfish (i.e., bluegill and yellow perch), black bass (largemouth and smallmouth), northern pike, and walleye; other fish species are present and contribute to the fishery.

In 2019, experimental gill nets were the only fish sampling gear deployed at Roy Lake. Thus, the following summary will focus on those fish species assessed using gill nets (i.e., northern pike, walleye, and yellow perch).

- **Northern pike.** Northern pike numbers have declined since 2016. In 2019, relative abundance was considered low to moderate at 1.0/gill net; sampled northern pike ranged in length from 16.1 to 22.8 inches.
- **Walleye.** Walleyes were not abundant (3.6/gill net). Gill net captured walleyes ranged in length from 7.1 to 24.8 inches, of those that were at least 10.0 inches 65% were  $\geq 15.0$  inches and 35% were 20.0 inches or longer. Walleyes from 9 year classes (1998, 2008, 2010, 2011, 2013, 2014, and 2016 – 2018) contributed to the catch, most (6 of 9) were represented by 7 or fewer individuals. Those from the 2016 (age-3) cohort, which coincided with a fry stocking, and the naturally-produced 2017 (age-2) year class were the most abundant accounting for nearly 60% of the sample. The oldest walleye collected was from the 1998 (age-21) year class. Walleyes appear to grow well with mean length at capture values at age 3 from 13.8 to 16.2 inches since 2010. In 2019, the mean length of age-3 walleyes was 16.2 inches.
- **Yellow perch.** Yellow perch CPUE were higher in 2019 than 2018. At 8.3/gill net relative abundance was considered low to moderate in 2019. Of the 101 individuals sampled, only 1 exceeded 8.0 inches. Four cohorts (2014 – 2017) contributed to the catch; fish from the 2017 (age-2) year class were the most abundant accounting for 76% of sampled yellow perch. Growth is slow with mean length at capture values at age 3 from 5.1 to 7.2 inches since 2010. In 2019, the mean length at capture of age-3 fish was 6.6 inches.

For more detailed results see the computer generated South Dakota Statewide Fisheries Survey for Roy (Marshall; below).

# SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Roy, Marshall County

UJA-Lake-866-001

2019

## Lake Information

<b>Name:</b>	Roy	<b>Maximum Depth:</b>	21 Feet
<b>County:</b>	Marshall	<b>Mean Depth:</b>	10 Feet
		<b>OHWM Elevation:</b>	1,796
<b>Surface Area:</b>	2,113 Acres	<b>Outlet Elevation:</b>	1,795

## Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
AFS std gill net	Jul 08, 2019	4 net-nights
AFS std gill net	Jul 09, 2019	4 net-nights
AFS std gill net	Jul 10, 2019	4 net-nights

## **Common Fish Species Present**

Walleye

Smallmouth Bass

Northern Pike

Largemouth Bass

Bluegill

Black Crappie

Yellow Perch

White Sucker

Black Bullhead

Common Carp

---

## Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- **AFS std gill net** – 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- **std experimental gill net for non-Missouri River waters** - 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- **std experimental gill net for Missouri River reservoirs** – 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$CPUE = \frac{\text{number of fish}}{\text{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left( \frac{\text{number of fish} \geq \text{quality length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

$$PSD - P = \left( \frac{\text{number of fish} \geq \text{preferred length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (*W*) compared to a length-specific standard weight (*Ws*) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left( \frac{W}{Ws} \right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (S), quality (Q), preferred (P), memorable (M) and trophy (T). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

## Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

\* **Methods/Species that ignore stock length**

Gear	Species	Sample Size (n)*	Abundance		Stock Density Indices			Condition		
			CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
AFS std gill net	Black Bullhead	5	0.4	0.4	100		100		86	3
	Black Crappie	18	1.4	0.6	24		12		117	3
	Bluegill	5	0.4	0.2	60		20		112	5
	Common Carp	5	0.4	0.3	100		20		101	4
	Largemouth Bass	5	0.4	0.4	60		40		125	2
	Northern Pike	12	1.0	0.4	58	24	0		94	3
	Smallmouth Bass	14	1.2	0.8	100		93		95	2
	Walleye	45	3.6	1.3	65	11	35	11	90	1
	White Sucker	8	0.7	0.4	100		100		97	4
	Yellow Perch	101	8.3	3.2	1		0		96	1

## 10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

\* Methods/Species that ignore stock length; \*\*AFS standard frame nets used in 2017

Gear	Species	CPUE										Avg
		2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	
AFS std gill net	Black Bullhead							1.3	0.9	2.1	0.4	1.2
	Black Crappie							0.3	0.2	0.8	1.4	0.7
	Bluegill							0.3	0.3	1.1	0.4	0.5
	Common Carp							0.0	0.0	0.1	0.4	0.1
	Largemouth Bass							0.1	0.0	0.1	0.4	0.2
	Northern Pike							3.2	2.3	1.7	1.0	2.1
	Smallmouth Bass							2.6	4.2	1.1	1.2	2.3
	Walleye							2.4	3.4	2.2	3.6	2.9
	White Sucker							2.8	2.8	1.9	0.7	2.1
	Yellow Perch							7.4	2.8	4.2	8.3	5.7
boat shocker	Largemouth Bass	26.3		36.7		58.8		44.0				41.5
boat shocker	Smallmouth Bass	42.4		110.8		19.7		3.0				44.0
fall night EF-WAE*	Walleye	154.0	466.5	4.0	286.0	90.0	27.0	87.0	24.5	38.0		130.8
frame net (std 3/4 in)**	Black Bullhead	0.5	0.6	8.2	6.5	3.3	3.0		1.4			3.4
	Black Crappie	0.2	0.5	0.6	0.6	0.2	0.3		0.2			0.4
	Bluegill	8.2	7.2	12.9	8.0	8.0	56.6		22.5			17.6
	Common Carp	0.1	0.0	0.1	0.1	0.0	0.0		0.2			0.1
	Green Sunfish	0.1	0.0	0.7	0.2	0.0	0.2		2.0			0.5
	Largemouth Bass	0.0	0.0	0.0	0.0	0.0	0.0		0.1			0.0
	Northern Pike	0.5	0.5	1.5	1.2	1.0	0.5		0.7			0.8
	Smallmouth Bass	0.5	0.8	0.3	0.2	0.9	0.3		0.0			0.4
	Walleye	0.3	0.1	0.5	0.2	0.2	0.0		0.1			0.2
	White Sucker	0.0	0.0	0.0	0.1	0.0	0.2		0.0			0.0
Yellow Perch	20.9	19.6	21.3	9.8	4.1	2.0		7.8			12.2	
std exp gill net	Black Bullhead	0.0	0.0	4.3	1.2	1.5	0.3					1.2
	Black Crappie	0.1	0.0	0.2	0.0	0.3	0.2					0.1
	Bluegill	0.0	0.0	0.0	0.3	0.0	0.7					0.2
	Common Carp	0.2	0.1	0.0	0.0	0.0	0.2					0.1
	Northern Pike	0.9	2.6	10.3	7.5	6.3	6.0					5.6
	Smallmouth Bass	0.3	0.1	0.5	2.3	4.0	2.3					1.6
	Walleye	1.1	0.6	2.8	8.3	6.3	6.5					4.3
	White Sucker	1.6	2.4	6.7	4.7	8.5	5.0					4.8
	Yellow Perch	17.0	26.8	99.3	82.2	10.0	23.3					43.1

## 10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

Gear	Species	Index	Year										
			2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	
AFS std gill net	Northern Pike	PSD							89	82	65	58	
		PSD-P							5	4	5	0	
		Wr							89	89	92	94	
	Walleye	PSD								76	90	69	65
		PSD-P								34	34	38	35
		Wr								91	87	94	90
	Yellow Perch	PSD								2	0	4	1
		PSD-P								1	0	0	0
		Wr								97	102	95	96
std exp gill net	Northern Pike	PSD	81	66	65	51	61	81					
		PSD-P	19	15	5	0	5	6					
		Wr	93	90	90	88	85	88					
	Walleye	PSD	45	64	41	36	68	77					
		PSD-P	10	27	41	14	18	8					
		Wr	87	94	86	91	88	90					
	Yellow Perch	PSD	0	0	7	13	7	3					
		PSD-P	0	0	0	0	0	0					
		Wr	100	103	101	91	99	92					



## Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Walleye

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2019	45	182 (1)	293 (15)	412 (11)		487 (5)	559 (2)		549 (7)	575 (2)	538 (2)
2018	28	213 (2)	326 (8)	397 (3)	466 (3)		521 (2)	524 (4)	557 (3)	526 (1)	612 (2)
2017	42	215 (1)		395 (13)	450 (1)		511 (16)	510 (4)	547 (4)		661 (3)
2016	30		285 (6)	371 (2)	455 (1)	476 (9)	486 (4)	597 (2)	626 (1)		637 (5)
2015	42	194 (3)	282 (2)	356 (7)	427 (23)	443 (4)	575 (1)				678 (2)
2014	40		232 (2)	377 (19)	408 (10)	482 (2)	592 (2)	476 (1)			652 (4)
2013	50		300 (21)	367 (15)	424 (6)	535 (3)					631 (5)
2012	23	197 (6)	294 (2)	351 (8)	530 (1)						622 (6)
2011	16	183 (4)	276 (5)	398 (1)	485 (2)	464 (1)		513 (1)			661 (2)
2010	24	177 (3)	294 (12)	384 (1)	464 (2)	523 (3)	473 (2)			483 (1)	

Species: Yellow Perch

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2019	101		145 (77)	167 (19)	183 (1)	193 (4)					
2018	50		141 (9)	160 (32)	175 (8)	205 (1)					
2017	34		138 (10)	157 (23)	186 (1)						
2016	89		140 (27)	153 (37)	165 (22)	195 (1)	249 (2)				
2015	728	99 (187)	112 (423)	153 (113)	194 (2)	219 (3)					
2014	275	97 (55)	116 (172)	129 (23)	182 (22)	214 (4)					
2013	1069	99 (563)	138 (44)	167 (258)	187 (165)	202 (33)	205 (9)				
2012	644	102 (45)	149 (412)	184 (106)	195 (80)						
2011	1506	99 (764)	128 (548)	156 (194)							
2010	1178	97 (208)	121 (926)	158 (44)							

## Fish Condition

Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).





Species	Year	Length Groups							
		S-Q		Q-P		P-M		M	
		N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Northern Pike Gill Net	2015	7	93 (2.8)	27	87 (2.3)	2	82 (1.7)	0	
	2016	4	90 (2.4)	32	88 (1.2)	2	95 (4.1)	0	
	2017	5	98 (3.4)	22	87 (1.4)	1	83	0	
	2018	7	99 (4.6)	12	89 (0.9)	1	70	0	
	2019	5	103 (0.9)	7	87 (2.1)	0		0	
Walleye Gill Net	2015	9	90 (0.8)	27	90 (1.0)	1	89	2	79 (2.6)
	2016	7	92 (1.3)	12	94 (2.0)	7	90 (1.2)	3	81 (3.8)
	2017	4	84 (2.6)	23	89 (1.0)	12	86 (1.7)	2	77 (3.4)
	2018	8	93 (1.6)	8	99 (1.5)	9	92 (2.5)	1	77
	2019	15	91 (1.6)	13	90 (1.7)	14	88 (0.9)	1	91
Yellow Perch Gill Net	2015	136	93 (0.9)	4	87 (1.5)	0		0	
	2016	87	97 (0.8)	1	93	1	81	0	
	2017	33	102 (1.2)	0		0		0	
	2018	48	95 (1.1)	2	96 (1.2)	0		0	
	2019	99	96 (1.0)	1	93	0		0	

## Length Frequency Distribution



Length frequency histogram of species sampled by year.

Species: Northern Pike




Gear: AFS std gill net




Species: Northern Pike  
 Gear: std exp gill net




Species: Walleye  
 Gear: AFS std gill net






2019

Species: Walleye  
Gear: std exp gill net




2014




2015


Species: Yellow Perch  
Gear: AFS std gill net




2016



2017




2018




2019

Species: Yellow Perch  
Gear: std exp gill net



2014




2015


## Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.


Species: Northern Pike  
Gear: AFS std gill net




Species: Northern Pike  
Gear: std exp gill net



Species: Walleye  
Gear: AFS std gill net




Species: Walleye  
Gear: std exp gill net






Species: Yellow Perch  
Gear: AFS std gill net



Species: Yellow Perch  
Gear: std exp gill net



## **Fish Stocking**

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2013	Walleye	Fry	850,000
2016	Walleye	Fry	1,000,000
2018	Walleye	Fry	1,030,000
2019	Walleye	Fry	1,030,000