SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Isabel, Dewey County GRA-Lake-613-000 2019

Lake Information

Name: Isabel Maximum Depth: 25 Feet

County: Dewey Mean Depth: 9 Feet

Legal Description: T17-R22-S16

Surface Area: 107 Acres

Surveys and Investigations

Survey methods used by gear type, date, and effort.

| Gear | Date | Effort |
|----------------------|--------------|--------------|
| boat shocker (night) | Oct 01, 2019 | 3600 seconds |
| boat shocker (night) | Sep 18, 2019 | 3600 seconds |

Common Fish Species Present

Walleye

Largemouth Bass

Bluegill

Black Crappie

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- AFS std gill net 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- std experimental gill net for non-Missouri River waters 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- std experimental gill net for Missouri River reservoirs 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$\mathit{CPUE} = \frac{\mathit{number of fish}}{\mathit{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{number\ of\ fish \ge quality\ length}{number\ of\ fish \ge stock\ length}\right) \times 100$$

$$PSD - P = \left(\frac{number\ of\ fish \ge preferred\ length}{number\ of\ fish \ge stock\ length}\right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{Ws}\right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (S), quality (Q), preferred (P), memorable (M) and trophy (T). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

| | St | Stock Quality Preferred | | Memorable | | Trophy | | | | |
|-----------------|------|-------------------------|------|-----------|------|--------|------|------|------|------|
| Species Name | (in) | (cm) | (in) | (cm) | (in) | (cm) | (in) | (cm) | (in) | (cm) |
| Black Bullhead | 6 | 15 | 9 | 23 | 12 | 30 | 15 | 38 | 18 | 46 |
| Black Crappie | 5 | 13 | 8 | 20 | 10 | 25 | 12 | 30 | 15 | 38 |
| Bluegill | 3 | 8 | 6 | 15 | 8 | 20 | 10 | 25 | 12 | 30 |
| Brown Trout | 8 | 20 | 12 | 30 | 16 | 40 | 20 | 50 | 18 | 46 |
| Channel Catfish | 11 | 28 | 16 | 41 | 24 | 61 | 28 | 71 | 36 | 91 |
| Freshwater Drum | 8 | 20 | 12 | 30 | 15 | 38 | 20 | 51 | 25 | 63 |
| Lake Trout | 12 | 30 | 20 | 50 | 26 | 65 | 31 | 80 | 39 | 100 |
| Largemouth Bass | 8 | 20 | 12 | 30 | 15 | 38 | 20 | 51 | 25 | 63 |
| Muskellunge | 20 | 51 | 30 | 76 | 38 | 97 | 42 | 107 | 50 | 127 |
| Northern Pike | 14 | 35 | 21 | 53 | 28 | 71 | 34 | 86 | 44 | 112 |
| Pumpkinseed | 3 | 8 | 6 | 15 | 8 | 20 | 10 | 25 | 12 | 30 |
| Rainbow Trout | 10 | 25 | 16 | 40 | 20 | 50 | 26 | 65 | 31 | 80 |
| Rudd | 6 | 15 | 10 | 25 | 12 | 30 | 15 | 38 | 19 | 48 |
| Sauger | 8 | 20 | 12 | 30 | 15 | 38 | 20 | 51 | 25 | 63 |
| Smallmouth Bass | 7 | 18 | 11 | 28 | 14 | 35 | 17 | 43 | 20 | 51 |
| Walleye | 10 | 25 | 15 | 38 | 20 | 51 | 25 | 63 | 30 | 76 |
| White Bass | 6 | 15 | 9 | 23 | 12 | 30 | 15 | 38 | 18 | 46 |
| White Crappie | 5 | 13 | 8 | 20 | 10 | 25 | 12 | 30 | 15 | 38 |
| Yellow Bullhead | 4 | 10 | 7 | 18 | 9 | 23 | 11 | 28 | 14 | 36 |
| Yellow Perch | 5 | 13 | 8 | 20 | 10 | 25 | 12 | 30 | 15 | 38 |

Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

* Methods/Species that ignore stock length

| | | | Abun | dance | Stock Density Indices | | | | | Condition | |
|-------------------------|-----------------|--------------------|------|-------|-----------------------|-------|-------|-------|-----|-----------|--|
| Gear | Species | Sample Size (n) | CPUE | CI-80 | PSD | CI-80 | PSD-P | CI-80 | Wr | CI-80 | |
| boat shocker (night) | Largemouth Bass | 71 | 35.5 | 8.4 | 75 | 8 | 46 | 9 | 105 | 2 | |

10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

| | | | | | | | CPUE | | | | | |
|------------------|-----------------|------|------|------|------|------|------|------|------|------|------|-------|
| Gear | Species | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 | Avg |
| AFS std gill net | Black Crappie | | | | | | | | | 1.3 | | 1.30 |
| | Bluegill | | | | | | | | | 0.3 | | 0.30 |
| | Northern Pike | | | | | | | | | 2.0 | | 2.00 |
| | Walleye | | | | | | | | | 0.8 | | 0.80 |
| | Yellow Perch | | | | | | | | | 31.3 | | 31.30 |
| boat shocker | Largemouth Bass | | 27.0 | | | 13.0 | | | | 39.5 | 35.5 | 28.75 |
| (night) | Smallmouth Bass | | 0.0 | | | 3.0 | | | | 0.0 | 0.0 | 0.75 |
| | Walleye | | 23.0 | | | 9.0 | | | | 0.0 | 0.0 | 8.00 |
| frame net (std | Black Bullhead | | 3.8 | | | 4.8 | | | | 0.0 | | 2.87 |
| 3/4 in) | Black Crappie | | 9.3 | | | 8.0 | | | | 17.4 | | 11.57 |
| | Bluegill | | 1.6 | | | 1.0 | | | | 9.2 | | 3.93 |
| | Northern Pike | | 0.3 | | | 0.5 | | | | 8.0 | | 0.53 |
| | Walleye | | 0.0 | | | 0.1 | | | | 0.0 | | 0.03 |
| | Yellow Perch | | 0.3 | | | 0.7 | | | | 9.8 | | 3.60 |
| std exp gill net | Black Bullhead | | 23.5 | | | 2.0 | | | | | | 12.75 |
| | Black Crappie | | 0.0 | | | 2.5 | | | | | | 1.25 |
| | Bluegill | | 0.0 | | | 0.0 | | | | | | 0.00 |
| | Northern Pike | | 0.5 | | | 3.0 | | | | | | 1.75 |
| | Walleye | | 0.5 | | | 5.0 | | | | | | 2.75 |
| | Yellow Perch | | 0.5 | | | 5.5 | | | | | | 3.00 |

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

| | | | | | | | Ye | ai | | | | |
|------------------|-----------------|-------|------|------|------|------|------|------|------|------|------|------|
| Gear | Species | Index | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 |
| AFS std gill net | Black Crappie | PSD | | | | | | | | | 80 | |
| | | PSD-P | | | | | | | | | 0 | |
| | | Wr | | | | | | | | | 98 | |
| | Bluegill | PSD | | | | | | | | | 100 | |
| | | PSD-P | | | | | | | | | 0 | |
| | | Wr | | | | | | | | | 93 | |
| | Walleye | PSD | | | | | | | | | 100 | |
| | | PSD-P | | | | | | | | | 67 | |
| | | Wr | | | | | | | | | 89 | |
| boat shocker | Largemouth Bass | PSD | | 52 | | | 100 | | | | 81 | 75 |
| (night) | | PSD-P | | 11 | | | 69 | | | | 43 | 46 |
| | | Wr | | 108 | | | 104 | | | | 107 | 105 |
| | Walleye | PSD | | 64 | | | 56 | | | | | |
| | | PSD-P | | 0 | | | 11 | | | | | |
| | | Wr | | 90 | | | 85 | | | | | |
| frame net (std | Black Crappie | PSD | | 70 | | | 85 | | | | 63 | |
| 3/4 in) | | PSD-P | | 2 | | | 20 | | | | 3 | |
| | | Wr | | 99 | | | 97 | | | | 102 | |
| | Bluegill | PSD | | 31 | | | 80 | | | | 48 | |
| | | PSD-P | | 0 | | | 40 | | | | 0 | |
| | | Wr | | 120 | | | 107 | | | | 101 | |
| | Walleye | PSD | | | | | 100 | | | | 0 | |
| | | PSD-P | | | | | 100 | | | | 0 | |
| | | Wr | | | | | 80 | | | | | |
| std exp gill net | Black Crappie | PSD | | | | | 100 | | | | | |
| | | PSD-P | | | | | 40 | | | | | |
| | | Wr | | | | | 94 | | | | | |
| | Walleye | PSD | | 0 | | | 90 | | | | | |
| | | PSD-P | | 0 | | | 50 | | | | | |
| | | Wr | | | | | 91 | | | | | |

Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Black Crappie

| Mean Length (expanded sample number) at capture by age Year N | 9 246 (6) 9 | 10 24 (2 |
|--|----------------------|------------|
| Company | 9 | (2 |
| Control Cont | | 10 |
| Mean Length (expanded sample number) at capture by age Year N | | 10 |
| Mean Length (expanded sample number) at capture by age Year N | | 10 |
| Year N 1 2 3 4 5 6 7 8 2018 92 116 134 154 186 174 182 2014 10 126 126 172 172 186 204 2011 16 96 170 173 173 173 173 174 186 204 170 173 173 174 186 204 170 173 173 174 186 204 170 173 173 174 186 204 170 173 173 174 186 204 170 173 173 173 173 173 186 204 180 170 173 173 180 180 180 180 170 173 180 180 180 180 180 180 180 180 180 180 180 180 180 180 180 180 | | 10 |
| 2018 92 116 134 154 186 174 182 (22) (14) (27) (6) (14) (11) (20) (21) (14) (11) (11) (11) (11) (11) (12) (2) (2) (2) (2) (2) (2) (2) (2) (2) (| | 10 |
| Company | 211 | |
| Control of the cont | 211 | |
| Mean Length (expanded sample number) at capture by age Year N 1 2 3 4 5 6 7 8 | (2) | |
| Mean Length (expanded sample number) at capture by age Year N 1 2 3 4 5 6 7 8 2018 71 166 194 243 292 325 342 403 385 (5) (3) (10) (5) (10) (15) (8) (5) 2014 14 133 376 387 403 (1) 376 387 403 (5) (2) (5) | | |
| Year N 1 2 3 4 5 6 7 8 2018 71 166 194 243 292 325 342 403 385 (5) (3) (10) (5) (10) (15) (8) (5) 2014 14 133 376 387 403 (1) (5) (2) (5) | | |
| 2018 71 166 194 243 292 325 342 403 385 (5) (3) (10) (5) (10) (15) (8) (5) 2014 14 133 376 387 403 (1) (5) (2) (5) | | |
| (5) (3) (10) (5) (10) (15) (8) (5) 2014 14 133 376 387 403 (1) (5) (2) (5) | 9 | 10 |
| (1) (5) (2) (5) | 436 (3) | 439 (10 |
| 2014 24 402 244 245 242 244 402 442 | 533 (1) | |
| 2011 31 193 241 215 312 314 400 448 (5) (4) (1) (16) (2) (2) (1) | | |
| Species: Walleye | | |
| Mean Length (expanded sample number) at capture by age | | |
| Year N 1 2 3 4 5 6 7 8 | 9 | 10 |
| 2014 12 232 400 543 (3) (4) (5) | | |

Fish Condition


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

| | | | Length Groups | | | | | | | | |
|------------------------------------|------|----|---------------|-----|--------------|----|--------------|---|--------------|--|--|
| | | | S-Q | | Q-P | | P-M | | M | | |
| Species | Year | N | Wr (SE) | N | Wr (SE) | N | Wr (SE) | N | Wr (SE) | | |
| Black Crappie Frame Net | 2018 | 64 | 110 (0.9) | 105 | 99 (0.7) | 5 | 86 (1.0) | 0 | | | |
| Bluegill Frame Net | 2018 | 48 | 104 (0.9) | 44 | 99 (1.1) | 0 | | 0 | | | |
| Largemouth Bass Electro Fishing | 2018 | 15 | 105 (1.7) | 30 | 105 (1.4) | 34 | 109 (1.6) | 0 | | | |
| | 2019 | 18 | 103 (3.2) | 20 | 107 (3.1) | 30 | 105 (1.5) | 3 | 108 (6.1) | | |
| Walleye Gill Net | 2018 | 0 | | 1 | 82 | 1 | 89 | 1 | 96 | | |

Length Frequency Distribution

Length frequency histogram of species sampled by year.

Species: Black Crappie Gear: frame net (std 3/4 in)



19

21

23

Species: Bluegill

0

5

Gear: frame net (std 3/4 in)


9

11

13

15

17 Total Length (cm)





27

29


25

Species: Largemouth Bass Gear: boat shocker (night)






Species: Walleye Gear: std exp gill net



Historic Fish Sizes and Relative Abundance


Size distribution per net by color for species sampled by year.

Species: Black Crappie Gear: frame net (std 3/4 in)




Species: Bluegill


Gear: frame net (std 3/4 in)




Species: Largemouth Bass Gear: boat shocker (night)



Species: Walleye Gear: AFS std gill net



Species: Walleye Gear: std exp gill net



Fish Stocking

Number of fish stocked by year, species, and size.

| Year | Species | Size | Number |
|------|---------|------------------|--------|
| 2008 | Walleye | Large Fingerling | 910 |
| 2010 | Walleye | Small Fingerling | 8,100 |
| 2013 | Walleye | Large Fingerling | 688 |
| 2015 | Walleye | Large Fingerling | 540 |
| 2017 | Walleye | Large Fingerling | 2,800 |
| 2019 | Walleye | Large Fingerling | 2,868 |
| | | | |