SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Henry, Bon Homme County LJA-Lake-588-000 2019

Lake Information

Name: Henry Maximum Depth: 37 Feet

County: Bon Homme Mean Depth: 14 Feet

Legal Description: T96-R58-Sec.9-10

Surface Area: 104 Acres

Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
boat shocker (day)	Jun 10, 2019	5400 seconds
frame net (std 3/4 in)	Jun 03, 2019	5 net-nights

Common Fish Species Present

Largemouth Bass

Black Crappie

Bluegill

White Sucker

Black Bullhead

Common Carp

Yellow Perch

Sunfish Hybrid

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- AFS std gill net 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- std experimental gill net for non-Missouri River waters 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- std experimental gill net for Missouri River reservoirs 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$\mathit{CPUE} = \frac{\mathit{number of fish}}{\mathit{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{number\ of\ fish \ge quality\ length}{number\ of\ fish \ge stock\ length}\right) \times 100$$

$$PSD - P = \left(\frac{number\ of\ fish \ge preferred\ length}{number\ of\ fish \ge stock\ length}\right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{Ws}\right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (S), quality (Q), preferred (P), memorable (M) and trophy (T). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

	Stock		Qu	ality	Pref	erred	Mem	orable	Trophy	
Species Name	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

* Methods/Species that ignore stock length

			Abundance		St	tock Der	es	Condition		
Gear	Species	Sample Size (n)	CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
boat shocker (day)	Largemouth Bass	141	91.3	21.5	33	6	24	5	95	2
frame net (std 3/4	Black Bullhead	7	1.2	1.2	0		0			
in)	Black Crappie	121	24.2	29.7	19	5	1		85	1
	Bluegill	32	6.4	6.2	53	13	0		87	2
	Common Carp	2	0.4	0.6	100		50			
	Largemouth Bass	2	0.4	0.6	100		100		107	10
	Sunfish Hybrid	1	0.0	0.0						
	White Sucker	7	1.4	1.1	100		100			
	Yellow Perch	1	0.2	0.3	100		0		92	

10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

							CPUE					
Gear	Species	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	Avg
AFS std frame	Black Bullhead				'				0.4			0.40
net	Black Crappie								4.0			4.00
	Bluegill								1.8			1.80
	Green Sunfish								0.2			0.20
	White Sucker								1.0			1.00
AFS std gill net	Black Bullhead								0.3			0.30
	Black Crappie								3.0			3.00
	Common Carp								0.0			0.00
	Green Sunfish								0.2			0.20
	Largemouth Bass								0.2			0.20
	Saugeye								0.2			0.20
	White Sucker								4.0			4.00
	Yellow Perch								5.0			5.00
boat shocker (day)	Largemouth Bass										91.3	91.30
boat shocker (night)	Largemouth Bass				33.0	37.5	25.5					32.00
frame net (std	Black Bullhead		1.5		0.7	0.3	0.0	0.2		0.0	1.2	0.56
3/4 in)	Black Crappie		20.0		16.1	16.0	16.9	2.2		10.4	24.2	15.11
	Bluegill		54.4		21.1	24.6	20.5	16.1		5.0	6.4	21.16
	Channel Catfish		0.0		0.0	0.9	0.1	0.1		0.0	0.0	0.16
	Common Carp		0.2		0.4	0.0	0.1	0.1		0.2	0.4	0.20
	Green Sunfish		0.1		0.0	0.0	0.0	0.1		0.0	0.0	0.03
	Largemouth Bass		0.1		0.1	0.0	0.0	0.0		0.0	0.4	0.09
	Sunfish Hybrid		0.0		0.0	0.0	0.0	0.0		0.0	0.0	0.00
	Walleye		0.0		0.0	0.0	0.0	0.0		0.0	0.0	0.00
	White Sucker		1.8		3.5	2.0	0.6	0.4		1.0	1.4	1.53
	Yellow Perch		0.1		0.0	0.1	0.0	0.4		0.4	0.2	0.17

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

			Year									
Gear	Species	Index	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
AFS std frame	Black Bullhead	PSD								50		
net		PSD-P								0		
	Black Crappie	PSD								65		
		PSD-P								20		
		Wr								89		
	Bluegill	PSD								78		
		PSD-P								0		
		Wr								99		
	White Sucker	PSD								100		
		PSD-P								80		
AFS std gill net	Black Bullhead	PSD								0		
		PSD-P								0		
	Black Crappie	PSD								33		
		PSD-P								6		
		Wr								96		
	Common Carp	PSD								0		
		PSD-P								0		
	Largemouth Bass	PSD								100		
		PSD-P								100		
		Wr								99		
	White Sucker	PSD								92		
		PSD-P								67		
	Yellow Perch	PSD								37		
		PSD-P								0		
		Wr								84		
boat shocker	Largemouth Bass	PSD										33
(day)		PSD-P										24
		Wr										95
boat shocker	Largemouth Bass	PSD				76	93	78				
(night)		PSD-P				41	35	29				
		Wr				98	89	92				

							Ye	ar				
Gear	Species	Index	2010 20	11 2	2012	2013	2014	2015	2016	2017	2018	2019
frame net (std	Black Bullhead	PSD		80		100	100		50			0
3/4 in)		PSD-P		40		71	67		0			0
		Wr		92		72						
	Black Crappie	PSD		6		94	96	36	41		23	19
		PSD-P		0		0	0	0	9		10	1
		Wr		96		86	91	93	97		93	85
	Bluegill	PSD		18		100	90	84	58		32	53
		PSD-P		0		0	0	2	2		0	0
		Wr		94		83	92	90	107		88	87
	Common Carp	PSD		50		100		100	100		0	100
		PSD-P		0		100		100	100		0	50
		Wr		83		80						
	Largemouth Bass	PSD	1	00		100						100
		PSD-P		0		100						100
		Wr	1	10		95						107
	White Sucker	PSD	1	00		100	100	100	75		100	100
		PSD-P	1	00		100	100	100	75		100	100
		Wr		84		79						
	Yellow Perch	PSD		0			100		25		0	100
		PSD-P		0			0		0		0	0
		Wr		81			82		135		76	92

Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Black Crappie

2015 172 132 191 226 236 (12) (112) (15) (34) 2014 161 138 230 223 (10) (143) 2013 161 205 213 (56) (105) 2011 204 145 226 244 (193) (9) (2) Species: Bluegill Mean Length (expanded sample number) at capture by age Year N 1 2 3 4 5 6 7 8 9 10- 2013 211 171 178 (179) (32) 2011 544 98 133 148 191 173 (164) (243) (125) (5) (7) Species: Largemouth Bass Mean Length (expanded sample number) at capture by age Mean Length (expanded sample number) at capture by age Mean Length (expanded sample number) at capture by age Mean Length (expanded sample number) at capture by age					Mean Len	ıgth (expai	nded sam	ple numb	er) at cap	ture by ag	е	
Company	Year	N	1	2	3	4	5	6	7	8	9	10+
Comparison of	2015	172										
Continue	2014	161										
Mean Length (expanded sample number) at capture by age Year	2013	161										
Year N	2011	204										
Year N 1 2 3 4 5 6 7 8 9 10-10-10-10-10-10-10-10-10-10-10-10-10-1	Species: B	luegill										
2015 205 150 161 183 186 212 (75) (18) (42) (66) (4) 2014 246 108 146 162 177 183 (20) (4) (9) (199) (14) 2013 211 171 178 (179) (32) 2011 544 98 133 148 191 173 (164) (243) (125) (5) (7) Species: Largemouth Bass Mean Length (expanded sample number) at capture by age Year N 1 2 3 4 5 6 7 8 9 10- 2013 65 225 260 342 366 388 417 427 451 454					Mean Len	gth (expai	nded sam	ple numb	er) at cap	ture by ag	е	
(75) (18) (42) (66) (4) 2014 246 108 146 162 177 183 (20) (4) (9) (199) (14) 2013 211 171 178 (179) (32) 2011 544 98 133 148 191 173 (164) (243) (125) (5) (7) Species: Largemouth Bass Mean Length (expanded sample number) at capture by age Year N 1 2 3 4 5 6 7 8 9 10- 2013 65 225 260 342 366 388 417 427 451 454	Year	N	1	2	3	4	5	6	7	8	9	10+
(20) (4) (9) (199) (14) 2013 211	2015	205										
(179) (32) 2011 544 98 133 148 191 173 (164) (243) (125) (5) (7) Species: Largemouth Bass Mean Length (expanded sample number) at capture by age Year N 1 2 3 4 5 6 7 8 9 10- 2013 65 225 260 342 366 388 417 427 451 454	2014	246										
(164) (243) (125) (5) (7) Species: Largemouth Bass Mean Length (expanded sample number) at capture by age Year N 1 2 3 4 5 6 7 8 9 10- 2013 65 225 260 342 366 388 417 427 451 454	2013	211										
Mean Length (expanded sample number) at capture by age Year N 1 2 3 4 5 6 7 8 9 10- 2013 65 225 260 342 366 388 417 427 451 454	2011	544										
Year N 1 2 3 4 5 6 7 8 9 10- 2013 65 225 260 342 366 388 417 427 451 454	Species: L	argemou	th Bass									
2013 65 225 260 342 366 388 417 427 451 454					Mean Len	gth (expai	nded sam	ple numb	er) at cap	ture by ag	е	
	Year	N	1	2	3	4	5	6	7	8	9	10+
	2013	65										454 (1)

Fish Condition


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

					Length	Group	s		
			S-Q		Q-P		P-M		М
Species	Year	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Black Crappie Frame Net	2015	109	96 (1.1)	60	91 (0.7)	0		0	
	2016	13	104 (3.0)	7	87 (1.8)	2	83 (4.0)	0	
	2017	7	98 (3.1)	9	84 (1.0)	4	81 (1.8)	0	
	2018	40	96 (1.5)	7	87 (2.4)	5	77 (2.1)	0	
	2019	98	86 (0.5)	22	79 (0.8)	1	69	0	
Bluegill Frame Net	2015	33	89 (1.2)	168	90 (0.7)	4	89 (5.3)	0	
	2016	67	107 (1.3)	91	108 (1.4)	3	99 (1.1)	0	
	2017	2	106 (12.6)	7	96 (3.6)	0		0	
	2018	17	89 (2.8)	8	88 (1.8)	0		0	
	2019	15	90 (2.9)	17	85 (2.1)	0		0	
Largemouth Bass Electro Fishing	2015	11	89 (4.3)	25	92 (1.5)	12	96 (1.5)	3	89 (8.7)
	2019	92	96 (0.6)	12	92 (9.4)	31	94 (1.5)	2	105
Yellow Perch Gill Net	2017	19	87 (1.5)	11	80 (2.6)	0		0	


Length Frequency Distribution


Length frequency histogram of species sampled by year.


Species: Black Crappie Gear: AFS std frame net




Species: Black Crappie Gear: frame net (std 3/4 in)












Species: Bluegill


Gear: frame net (std 3/4 in)









Species: Largemouth Bass Gear: boat shocker (day)




Species: Largemouth Bass Gear: boat shocker (night)






Species: White Sucker Gear: AFS std gill net




Species: Yellow Perch Gear: AFS std gill net




Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.


Species: Black Bullhead Gear: AFS std gill net




Species: Black Crappie Gear: AFS std frame net




Species: Black Crappie Gear: frame net (std 3/4 in)




Species: Bluegill Gear: AFS std frame net




Species: Bluegill Gear: frame net (std 3/4 in)




Species: Common Carp Gear: AFS std gill net




Species: Largemouth Bass Gear: boat shocker (day)




Species: Largemouth Bass Gear: boat shocker (night)



Species: White Sucker Gear: AFS std gill net



Species: Yellow Perch Gear: AFS std gill net



Fish Stocking

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2011	Yellow Perch	Adult	1,747
2012	Yellow Perch	Adult	1,416
2012	Yellow Perch	Juvenile	7,875
2013	Channel Catfish	Large Fingerling	3,300
2014	Yellow Perch	Small Fingerling	40,820
2015	Largemouth Bass	Fingerling	11,400
2016	Yellow Perch	Adult	5,445