

SOUTH DAKOTA STATEWIDE FISHERIES SURVEY
New Underwood Dam, Pennington County
MCE-Lake-8-000
2019

Lake Information

Name: New Underwood Dam
County: Pennington
Surface Area: 18 Acres

Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
boat shocker (night)	Jun 24, 2019	900 seconds
frame net (std 3/4 in)	Jun 13, 2019	3 net-nights

Common Fish Species Present

Channel Catfish

Bluegill

Black Crappie

Yellow Perch

Largemouth Bass

Green Sunfish

Black Bullhead

Golden Shiner

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- **AFS std gill net** – 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- **std experimental gill net for non-Missouri River waters** - 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- **std experimental gill net for Missouri River reservoirs** – 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$CPUE = \frac{\text{number of fish}}{\text{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{\text{number of fish} \geq \text{quality length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

$$PSD - P = \left(\frac{\text{number of fish} \geq \text{preferred length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (**W**) compared to a length-specific standard weight (**Ws**) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{W_s} \right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (S), quality (Q), preferred (P), memorable (M) and trophy (T). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

* **Methods/Species that ignore stock length**

Gear	Species	Sample Size (n)	Abundance		Stock Density Indices			Condition	
			CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr
boat shocker (night)	Largemouth Bass	59	144.0	59.9	36	12	11	100	1
frame net (std 3/4 in)	Black Bullhead	8	2.7	2.7	100		63	102	4
	Black Crappie	47	15.3	14.8	83	9	9	97	1
	Bluegill	65	21.7	22.2	97		3	108	1
	Golden Shiner	2	0.0	0.0					
	Green Sunfish	53	17.7	14.5	13	7	0	106	1
	Yellow Perch	9	3.0	3.3	89		0	97	4

10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

Gear	Species	CPUE										Avg
		2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	
AFS std frame net	Black Bullhead								0.5			0.50
	Black Crappie								13.8			13.80
	Bluegill								62.8			62.80
	Golden Shiner								0.0			0.00
	Green Sunfish								0.3			0.30
	Yellow Perch								1.3			1.30
boat shocker (night)	Largemouth Bass		51.7			205.1				102.0	144.0	125.70
frame net (std 3/4 in)	Black Bullhead	5.0	6.0		23.3		2.3			1.7	2.7	6.83
	Black Crappie	3.3	1.8		3.3		1.3			1.7	15.3	4.45
	Bluegill	48.0	26.3		149.3		67.0			24.3	21.7	56.10
	Channel Catfish	0.3	0.3		0.0		0.5			0.0	0.0	0.18
	Golden Shiner	0.0	0.0		0.0		0.0			0.0	0.0	0.00
	Green Sunfish	9.0	0.3		0.0		0.8			0.0	17.7	4.63
	Largemouth Bass	0.0	0.0		0.0		0.5			0.0	0.0	0.08
	White Sucker	0.0	0.0		0.3		0.3			0.0	0.0	0.10
	Yellow Perch	0.7	3.5		14.0		55.3			4.0	3.0	13.42

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

Gear	Species	Index	Year												
			2010	2011	2012	2013	2014	2015	2016	2017	2018	2019			
AFS std frame net	Black Bullhead	PSD										100			
		PSD-P											100		
		Wr											89		
	Black Crappie	PSD											44		
		PSD-P											7		
		Wr											96		
	Bluegill	PSD											59		
		PSD-P											8		
		Wr											99		
	Green Sunfish	PSD											100		
		PSD-P											0		
		Wr											106		
	Yellow Perch	PSD											60		
		PSD-P											20		
		Wr											82		
boat shocker (night)	Largemouth Bass	PSD		45				16					41	36	
		PSD-P		5				11					12	11	
		Wr		94				95					96	100	
frame net (std 3/4 in)	Black Bullhead	PSD	60	63		99			100				100	100	
		PSD-P	0	0		14			89				60	63	
		Wr	104	102		112			93				99	102	
	Black Crappie	PSD	80	0		70			100				40	83	
		PSD-P	80	0		0			0				0	9	
		Wr	94	102		98			99				101	97	
	Bluegill	PSD	60	73		96			91				86	97	
		PSD-P	1	4		6			1				3	3	
		Wr	119	110		115			106				106	108	
	Channel Catfish	PSD	100	100					50						
		PSD-P	0	0					50						
		Wr	79						83						
	Green Sunfish	PSD	4	0					100					13	
		PSD-P	0	0					0					0	

Gear	Species	Index	Year									
			2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
frame net (std 3/4 in)	Green Sunfish	Wr	98	110					95			106
		PSD						100				
	Largemouth Bass	PSD-P						100				
		Wr						103				
		PSD	100	0		55		41			75	89
	Yellow Perch	PSD-P	50	0		0		0			0	0
		Wr	92	96		90		95			100	97

Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Bluegill

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2017	248	94 (1)	132 (94)	158 (51)	188 (72)	192 (22)	202 (8)				

Species: Largemouth Bass

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2014	296		205 (15)	195 (49)	222 (120)	269 (73)	323 (35)	441 (4)			
2011	40	203 (2)	209 (4)	301 (30)	349 (2)		410 (2)				

Fish Condition


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

Species	Year	Length Groups							
		S-Q		Q-P		P-M		M	
		N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Black Crappie Frame Net	2015	0		10	99 (0.9)	0		0	
	2017	31	98 (0.9)	20	95 (1.3)	4	88 (0.9)	0	
	2018	3	101 (1.6)	2	101 (2.7)	0		0	
	2019	8	103 (1.0)	34	96 (0.8)	4	90 (1.4)	0	
Bluegill Frame Net	2015	46	108 (0.8)	486	106 (0.3)	4		0	
	2017	102	105 (1.1)	130	93 (0.8)	19	83 (1.5)	0	
	2018	10	112 (3.8)	61	106 (1.2)	2	73 (0.0)	0	
	2019	2	107	61	108 (0.9)	2	95	0	
Largemouth Bass Electro Fishing	2018	20	97 (1.3)	10	96 (1.4)	4	91 (1.9)	0	
	2019	23	101 (1.2)	9	98 (2.1)	4	95 (2.9)	0	



Length Frequency Distribution

Length frequency histogram of species sampled by year.


Species: Black Crappie
Gear: AFS std frame net






Species: Black Crappie
Gear: frame net (std 3/4 in)





Species: Bluegill
Gear: AFS std frame net




Species: Bluegill
 Gear: frame net (std 3/4 in)



Species: Largemouth Bass
 Gear: boat shocker (night)






2019


Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.


Species: Black Crappie
Gear: AFS std frame net




Species: Black Crappie
Gear: frame net (std 3/4 in)




Species: Bluegill
Gear: AFS std frame net



Species: Bluegill
Gear: frame net (std 3/4 in)



Species: Largemouth Bass
Gear: boat shocker (night)



Fish Stocking

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2008	Black Crappie	Adult	300
2008	Bluegill	Adult	40
2008	Bluegill	Fingerling	8,000
2008	Largemouth Bass	Adult	90
2008	Largemouth Bass	Fingerling	1,500
2009	Channel Catfish	Adult	200
2009	Largemouth Bass	Juvenile	300
2010	Channel Catfish	Adult	74
2011	Channel Catfish	Adult	200
2012	Largemouth Bass	Fingerling	1,500
2014	Channel Catfish	Adult	143
2014	Yellow Perch	Adult	325
2015	Channel Catfish	Adult	100
2016	Channel Catfish	Adult	200
2017	Channel Catfish	Adult	137
2018	Channel Catfish	Adult	219
2018	Largemouth Bass	Juvenile	272
2019	Channel Catfish	Adult	200