

# SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Cavour, Beadle County

MJA-Lake-532-000

2019

## Lake Information

**Name:** Cavour **Maximum Depth:** 14 Feet  
**County:** Beadle **Mean Depth:** 4 Feet  
**Legal Description:** T111N- R60W-Sec. 20-22  
**Surface Area:** 528 Acres

## Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
AFS std gill net	Aug 29, 2019	4 net-nights

## **Common Fish Species Present**

Walleye

Black Bullhead

White Sucker

Shortnose Gar

Common Carp

Bigmouth Buffalo

---

## Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- **AFS std gill net** – 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- **std experimental gill net for non-Missouri River waters** - 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- **std experimental gill net for Missouri River reservoirs** – 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$CPUE = \frac{\text{number of fish}}{\text{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left( \frac{\text{number of fish} \geq \text{quality length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

$$PSD - P = \left( \frac{\text{number of fish} \geq \text{preferred length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (*W*) compared to a length-specific standard weight (*Ws*) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left( \frac{W}{W_s} \right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (S), quality (Q), preferred (P), memorable (M) and trophy (T). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

## **Catch Summary of Stock Length Fish**

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

**\* Methods/Species that ignore stock length**

Gear	Species	Sample Size (n)	Abundance		Stock Density Indices			Condition	
			CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr
AFS std gill net	Bigmouth Buffalo	19	0.0	0.0	0		0		
	Black Bullhead	22	4.0	2.1	88		0		
	Common Carp	91	0.0	0.0	0		0		
	Shortnose Gar	3	0.0	0.0					
	White Sucker	5	1.3	0.8	20		0		

## 10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

Gear	Species	CPUE										Avg
		2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	
AFS std gill net	Bigmouth Buffalo									1.2	0.0	0.60
	Black Bullhead									5.5	4.0	4.75
	Black Crappie									0.3	0.0	0.15
	Common Carp									22.3	0.0	11.15
	Freshwater Drum									2.2	0.0	1.10
	Shortnose Gar									0.0	0.0	0.00
	Walleye									0.3	0.0	0.15
	White Sucker									0.0	1.3	0.65
	Yellow Bullhead									0.2	0.0	0.10
frame net (std 3/4 in)	Bigmouth Buffalo	0.0	0.1		0.0	0.0	0.0	0.0		0.4		0.07
	Black Bullhead	23.2	577.3		342.0	159.6	247.2	234.0		22.2		229.36
	Black Crappie	199.8	83.5		20.2	17.2	17.4	25.4		14.0		53.93
	Bluegill	0.4	0.0		0.0	0.0	0.0	0.0		0.0		0.06
	Common Carp	5.1	11.4		4.2	0.2	4.0	8.2		3.6		5.24
	Freshwater Drum	0.0	0.0		0.0	0.0	0.0	0.0		0.0		0.00
	Green Sunfish	0.1	0.6		0.0	0.0	0.4	0.0		0.0		0.16
	Northern Pike	0.0	0.3		0.2	0.4	1.6	1.4		0.4		0.61
	Saugeye	0.0	0.0		0.0	0.0	0.0	0.0		0.0		0.00
	Sunfish Hybrid	0.0	0.0		0.0	0.0	0.0	0.0		0.0		0.00
	Walleye	0.4	0.1		0.0	0.6	1.0	21.6		2.6		3.76
	White Sucker	0.0	0.9		0.8	0.8	1.6	2.2		2.8		1.30
	Yellow Bullhead	0.0	0.7		0.0	1.4	5.2	0.0		1.2		1.21
	Yellow Perch	0.3	0.5		0.0	0.2	0.2	0.6		0.8		0.37
	std exp gill net	Bigmouth Buffalo	0.0	0.0		0.0	0.0	0.7	0.0			
Black Bullhead		32.7	62.7		30.7	170.7	56.0	50.0				67.13
Black Crappie		34.0	15.7		0.3	1.3	6.0	6.3				10.60
Channel Catfish		0.0	0.0		0.0	0.0	0.0	0.0				0.00
Common Carp		12.7	15.7		10.0	11.3	20.7	18.7				14.85
Freshwater Drum		0.0	0.0		0.0	0.0	0.3	0.7				0.17
Northern Pike		0.0	0.7		1.7	1.0	0.3	0.0				0.62
Saugeye		0.0	0.0		0.0	0.0	0.0	0.0				0.00
Walleye		1.3	0.3		2.0	3.7	13.3	3.0				3.93
White Sucker		0.0	0.3		0.7	0.0	0.0	0.7				0.28
Yellow Bullhead		0.0	0.3		0.0	0.0	0.3	0.0				0.10

		CPUE										
Gear	Species	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	Avg
std exp gill net	Yellow Perch	0.0	0.3		2.3	1.7	2.3	0.0				1.10

## 10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

Gear	Species	Index	Year										
			2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	
AFS std gill net	Bigmouth Buffalo	PSD										0	0
		PSD-P										0	0
	Black Bullhead	PSD										18	88
		PSD-P										0	0
	Common Carp	PSD										71	0
		PSD-P										3	0
	Walleye	PSD										0	
		PSD-P										0	
	White Sucker	Wr										92	
		PSD											
		PSD-P											0
frame net (std 3/4 in)	Bigmouth Buffalo	PSD		100								100	
		PSD-P		0								50	
		Wr		74									
	Black Bullhead	PSD	12	6		51	74	38	43			12	
		PSD-P	0	1		0	0	0	0			0	
		Wr	83	84		62							
	Common Carp	PSD	14	47		81	100	65	46			50	
		PSD-P	0	4		10	0	45	15			28	
		Wr	87	87		78							
	Walleye	PSD	75	0			33	80	93			8	
		PSD-P	50	0			0	20	8			8	
		Wr	91	98			88	91	73			89	
	White Sucker	PSD		100		100	100	100	100	100		100	
		PSD-P		100		100	100	88	100			100	
		Wr		98		80							
std exp gill net	Bigmouth Buffalo	PSD							0				
		PSD-P							0				
	Black Bullhead	PSD	3	3		53	65	52	12				
		PSD-P	0	0		0	0	1	0				
	Common Carp	Wr	92	99		71							
		PSD	18	49		70	62	65	50				



Gear	Species	Index	Year									
			2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
std exp gill net	Common Carp	PSD-P	0	0		10	15	3	2			
		Wr	85	87		78						
	Walleye	PSD	25	100		67	45	83	100			
		PSD-P	0	0		0	0	3	0			
		Wr	102	96		79	99	92	74			
	White Sucker	PSD		100		100				100		
		PSD-P		100		100				100		
		Wr		100		82						

## Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Walleye

Year	N	Mean Length (expanded sample number) at capture by age										
		1	2	3	4	5	6	7	8	9	10+	
2010	4	286 (1)		371 (3)								

## **Fish Condition**



Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

Species	Year	Length Groups							
		S-Q		Q-P		P-M		M	
		N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Walleye Gill Net	2015	7	87 (2.2)	32	93 (1.1)	1	91	0	
	2016	0		9	74 (2.8)	0		0	
	2018	2	92 (0.1)	0		0		0	



## Length Frequency Distribution


Length frequency histogram of species sampled by year.

Species: Black Bullhead  
Gear: AFS std gill net





Species: Black Bullhead  
Gear: std exp gill net








Species: Common Carp  
Gear: AFS std gill net






Species: Common Carp  
Gear: std exp gill net






Species: Walleye  
 Gear: std exp gill net




## Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.


Species: Black Bullhead  
Gear: AFS std gill net




Species: Black Bullhead  
Gear: std exp gill net



Species: Common Carp  
Gear: AFS std gill net




Species: Common Carp  
Gear: std exp gill net







Species: Walleye  
Gear: AFS std gill net




Species: Walleye  
Gear: std exp gill net



Species: White Sucker  
Gear: AFS std gill net



Species: White Sucker  
Gear: std exp gill net



## **Fish Stocking**

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2011	Walleye	Small Fingerling	23,340
2012	Walleye	Small Fingerling	46,400
2014	Walleye	Fry	115,000
2015	Walleye	Small Fingerling	27,920
2017	Walleye	Fingerling	44,840
2019	Walleye	Fry	550,000