

**SOUTH DAKOTA STATEWIDE FISHERIES SURVEY**  
**Campbell, Campbell County**  
**WMC-Lake-891-000**  
**2019**

**Lake Information**

<b>Name:</b>	Campbell	<b>Maximum Depth:</b>	22 Feet
<b>County:</b>	Campbell	<b>Mean Depth:</b>	9 Feet
<b>Legal Description:</b>	T126-R77-S11		
<b>Surface Area:</b>	42 Acres		

**Surveys and Investigations**

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
frame net (std 3/4 in)	Jul 09, 2019	4 net-nights
frame net (std 3/4 in)	Jul 10, 2019	5 net-nights

## **Common Fish Species Present**

Largemouth Bass

Black Crappie

Black Bullhead

---

## Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- **AFS std gill net** – 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- **std experimental gill net for non-Missouri River waters** - 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- **std experimental gill net for Missouri River reservoirs** – 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$CPUE = \frac{\text{number of fish}}{\text{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left( \frac{\text{number of fish} \geq \text{quality length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

$$PSD - P = \left( \frac{\text{number of fish} \geq \text{preferred length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (*W*) compared to a length-specific standard weight (*Ws*) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left( \frac{W}{W_s} \right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (S), quality (Q), preferred (P), memorable (M) and trophy (T). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

## **Catch Summary of Stock Length Fish**

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

**\* Methods/Species that ignore stock length**

Gear	Species	Sample Size (n)	Abundance		Stock Density Indices			Condition	
			CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr
frame net (std 3/4 in)	Black Bullhead	723	51.0	19.7	57	3	0	90	1
	Black Crappie	91	4.0	3.5	11		0	126	4

## 10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

Gear	Species	CPUE										Avg
		2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	
AFS std frame net	Black Bullhead								146.2			146.2
	Black Crappie								68.9			68.90
	Yellow Perch								1.0			1.00
boat shocker (night)	Largemouth Bass			0.0		2.0						1.00
frame net (std 3/4 in)	Black Bullhead			7.7		8.7					51.0	22.47
	Black Crappie			0.8		10.0					4.0	4.93
	Bluegill			0.0		0.0					0.0	0.00
	Common Carp			0.0		0.1					0.0	0.03
	Northern Pike			0.0		0.0					0.0	0.00
	White Crappie			0.1		0.0					0.0	0.03
	Yellow Perch			0.1		2.2					0.0	0.77

## 10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

Gear	Species	Index	Year									
			2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
AFS std frame net	Black Bullhead	PSD										0
		PSD-P										0
		Wr										94
	Black Crappie	PSD										7
		PSD-P										0
		Wr										108
boat shocker (night)	Largemouth Bass	PSD			0		100					
		PSD-P			0		0					
		Wr					113					
frame net (std 3/4 in)	Black Bullhead	PSD			26		17					57
		PSD-P			0		0					0
		Wr			100		92					90
	Black Crappie	PSD			88		55					11
		PSD-P			50		1					0
		Wr			110		111					126

## Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Black Crappie

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2017	689			170 (620)	194 (69)						
2014	200		199 (198)			297 (2)					
2012	16		182 (2)	237 (4)	258 (8)	300 (2)					

Species: Largemouth Bass

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2014	4	314 (4)									
2012	14		186 (14)								



## **Fish Condition**


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

Species	Year	Length Groups							
		S-Q		Q-P		P-M		M	
		N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Black Crappie Frame Net	2017	643	109 (0.6)	46	104 (1.6)	0		0	
	2019	32	129 (3.1)	4	107 (1.7)	0		0	



## Length Frequency Distribution

Length frequency histogram of species sampled by year.


Species: Black Crappie  
Gear: AFS std frame net



Species: Black Crappie  
Gear: frame net (std 3/4 in)




Species: Largemouth Bass  
Gear: boat shocker (night)




## Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.


Species: Black Crappie  
Gear: AFS std frame net



Species: Black Crappie  
Gear: frame net (std 3/4 in)



Species: Largemouth Bass  
Gear: boat shocker (night)



## **Fish Stocking**

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2008	Largemouth Bass	Adult	72
2008	Largemouth Bass	Fingerling	4,000
2009	Largemouth Bass	Juvenile	700
2010	Bluegill	Adult	146
2010	Largemouth Bass	Juvenile	211
2010	Yellow Perch	Adult	210
2011	Black Crappie	Adult	110
2011	Channel Catfish	Adult	147
2012	Black Crappie	Adult	90
2012	Largemouth Bass	Juvenile	200
2012	Yellow Perch	Adult	300
2014	Bluegill	Adult	100
2014	Channel Catfish	Juvenile	100
2014	Largemouth Bass	Fingerling	1,260
2014	Largemouth Bass	Large Fingerling	1,000
2015	Bluegill	Adult	300
2015	Largemouth Bass	Juvenile	292
2016	Channel Catfish	Adult	120
2016	Largemouth Bass	Adult	115
2017	Largemouth Bass	Adult	111
2018	Channel Catfish	Adult	112
2019	Channel Catfish	Adult	107
2019	Largemouth Bass	Adult	30
2019	Largemouth Bass	Juvenile	77