

SOUTH DAKOTA BLACK HILLS COLD STREAM SURVEY
Flynn Creek, Custer County
21-0034-00
2009

Table 1. Population and Biomass Estimates for Flynn Creek

| | | |
|--|---------------------------------|-------------------------------------|
| Site Number: 126 | Date Sampled: 05/26/2009 | Conductivity (umhos/cm): 217 |
| Site Midpoint (Latitude, Longitude): 43.68508330, -103.47288268 | Site Length (m): 100 | pH: 8.5 |
| Survey Completed by: South Dakota Game, Fish and Parks | Mean Width (m): 2.4 | Water Temperature (C): 17.3 |
| Stream Classification: None | Number of Pass: 1 | Voltage: 240 |

| Species | Size Class | Total Number Captured | Est. # in site | Upper 95% CI | Lower 95% CI | # per Hectacre | Kg per Hectacre | # per Km | # per Acre | lb. per Acre | # per Mile | Mean Length (mm) | Mean Weight (grams) |
|---------------|------------|-----------------------|----------------|--------------|--------------|----------------|-----------------|----------|------------|--------------|------------|------------------|---------------------|
| Longnose Dace | All | 3 | 3 | 3 | 3 | 126 | 0 | 30 | 51 | 0.30 | 48 | 44.3 | 2.7 |

Stream sites are classified based on the estimates of adult trout over eight inches that exist within the site. Estimates are derived from 3-pass backpack electrofishing surveys at 100-meter sites within streams. While all fish in the site are collected and measured, only trout species are used for classifications. Sites are numbered in accordance for the distance (m) from the stream confluence of a higher order stream/river.

| Based on number of trout in excess of eight inches | |
|--|---|
| Class BR1 | Number of wild Brown Trout exceeds 150 per acre |
| Class BR2 | Number of wild Brook Trout ranges from 25 to 150 per acre |
| Class BR3 | Number of wild Brown Trout is less than 25 per acre |
| Class BK1 | Number of wild Brook Trout exceeds 150 per acre |
| Class BK2 | Number of wild Brook Trout ranges from 25 to 150 per acre |
| Class BK3 | Number of wild Brook Trout is less than 25 per acre |
| Class RB1 | Number of wild Rainbow Trout exceeds 25 per acre |
| Class RB2 | Number of wild Rainbow Trout is less than 25 per acre |

Table 2. Species composition and population estimates on Flynn Creek, 1999-2009

Site 126

| Species/Size | Date | Pop Est. | (95% C.I.) | Stream Classification |
|---------------|--------------|----------|------------|-----------------------|
| Longnose Dace | Aug 22, 2006 | 38 | 37-41 | |
| | May 26, 2009 | 3 | 3-3 | |

Length Frequency Distribution

Length frequency histogram of species sampled by year.

Site: 126

Species: Longnose Dace

