

**SOUTH DAKOTA BLACK HILLS COLD STREAM SURVEY**  
**Gimlet Creek, Pennington County**  
**02-0095-00**  
**2014**

Table 1. Population and Biomass Estimates for Gimlet Creek

<b>Site Number:</b> 12	<b>Date Sampled:</b> 06/05/2014	<b>Conductivity (umhos/cm):</b> 134
<b>Site Midpoint (Latitude, Longitude):</b> 44.11364758, -103.64642536	<b>Site Length (m):</b> 100	<b>pH:</b>
<b>Survey Completed by:</b> South Dakota Game, Fish and Parks	<b>Mean Width (m):</b> 1.7	<b>Water Temperature (C):</b> 9.8
<b>Stream Classification:</b> BK0	<b>Number of Pass:</b> 1	<b>Voltage:</b> 325

Species	Size Class	Total Number Captured	Est. # in site	Upper 95% CI	Lower 95% CI	# per Hectacre	Kg per Hectacre	# per Km	# per Acre	lb. per Acre	# per Mile	Mean Length (mm)	Mean Weight (grams)
Brook Trout	<200mm	7	7	7	7	405	5	70	164	4.81	113	109.7	13.3
Brook Trout	All	7	7	7	7	405	5	70	164	4.81	113	109.7	13.3

<b>Site Number:</b> 32	<b>Date Sampled:</b> 06/05/2014	<b>Conductivity (umhos/cm):</b> 131
<b>Site Midpoint (Latitude, Longitude):</b> 44.12460040, -103.65368981	<b>Site Length (m):</b> 100	<b>pH:</b> 7.7
<b>Survey Completed by:</b> South Dakota Game, Fish and Parks	<b>Mean Width (m):</b> 1.6	<b>Water Temperature (C):</b> 17.0
<b>Stream Classification:</b> BK0, BR2	<b>Number of Pass:</b> 3	<b>Voltage:</b> 325

Species	Size Class	Total Number Captured	Est. # in site	Upper 95% CI	Lower 95% CI	# per Hectacre	Kg per Hectacre	# per Km	# per Acre	lb. per Acre	# per Mile	Mean Length (mm)	Mean Weight (grams)
Brook Trout	<200mm	46	47	50	46	2971	54	470	1203	47.93	756	113.8	18.1
Brook Trout	All	46	47	50	46	2971	54	470	1203	47.93	756	113.8	18.1
Brown Trout	>=200mm	3	3	4	3	190	19	30	77	17.27	48	223.0	102.0
Brown Trout	All	3	3	4	3	190	19	30	77	17.27	48	223.0	102.0

Stream sites are classified based on the estimates of adult trout over eight inches that exist within the site. Estimates are derived from 3-pass backpack electrofishing surveys at 100-meter sites within streams. While all fish in the site are collected and measured, only trout species are used for classifications. Sites are numbered in accordance for the distance (m) from the stream confluence of a higher order stream/river.

Based on number of trout in excess of eight inches	
<b>Class BR1</b>	Number of wild Brown Trout exceeds 150 per acre
<b>Class BR2</b>	Number of wild Brook Trout ranges from 25 to 150 per acre
<b>Class BR3</b>	Number of wild Brown Trout is less than 25 per acre
<b>Class BK1</b>	Number of wild Brook Trout exceeds 150 per acre
<b>Class BK2</b>	Number of wild Brook Trout ranges from 25 to 150 per acre
<b>Class BK3</b>	Number of wild Brook Trout is less than 25 per acre
<b>Class RB1</b>	Number of wild Rainbow Trout exceeds 25 per acre
<b>Class RB2</b>	Number of wild Rainbow Trout is less than 25 per acre

Table 2. Species composition and population estimates on Gimlet Creek, 2004-2014

**Site 12**

Species/Size	Date	Pop Est.	(95% C.I.)	Stream Classification
Brook Trout <200mm	Jun 16, 2008	2	2-2	
	Jun 05, 2014	7	7-7	
Brown Trout <200mm	Jun 16, 2008	1	1-1	
Brown Trout >=200mm	Jun 16, 2008	2	2-2	BR2

**Site 32**


Species/Size	Date	Pop Est.	(95% C.I.)	Stream Classification
Brook Trout <200mm	Jun 05, 2014	47	46-50	
Brown Trout >=200mm	Jun 05, 2014	3	3-4	BR2

## Length Frequency Distribution

Length frequency histogram of species sampled by year.

**Site:** 12

**Species:** Brook Trout



**Site:** 32

**Species:** Brook Trout

