SOUTH DAKOTA BLACK HILLS COLD STREAM SURVEY

Meadow Creek, Lawrence County

09-0072-00

1998

Table 1. Population and Biomass Estimates for Meadow Creek

| Site Number: 12 Site Midpoint (Latitude, Longitude): 44.28582280, -103.56238706 Survey Completed by: South Dakota Game, Fish and Parks Stream Classification: BK2 | | | | | | • | | 1.6 | (um pH: Wat | Conductivity (umhos/cm): pH: Water Temperature (C): Voltage: | | 378 8.6 15.6 200 | |
|--|---------------|-----------------------------|-------------------|-----------------|-----------------|-------------------|--------------------|-------------|-------------------|--|---------------|---------------------------|---------------------------|
| Species | Size Class | Total Number Captured | Est. # in site | Upper 95% Cl | Lower 95% Cl | # per Hectacre | Kg per Hectacre | # per Km | # per Acre | lb. per Acre | # per Mile | Mean Length (mm) | Mean Weight (grams) |
| Brook Trout | <200mm | 11 | 11 | 13 | 11 | 696 | 21 | 110 | 282 | 18.83 | 177 | 103.4 | 30.3 |
| Brook Trout | >=200mm | 1 | 1 | 1 | 1 | 63 | 8 | 10 | 26 | 6.89 | 16 | 226.0 | 122.0 |
| Brook Trout | All | 12 | 12 | 13 | 12 | 759 | 32 | 120 | 308 | 28.31 | 193 | 113.6 | 41.8 |
| Longnose Dace | All | 605 | 645 | 665 | 625 | 40823 | 176 | 6450 | 16533 | 157.06 | 10378 | 79.3 | 4.3 |
| Mountain Sucker | All | 44 | 47 | 53 | 44 | 2975 | 39 | 470 | 1205 | 34.53 | 756 | 103.6 | 13.0 |

Stream sites are classified based on the estimates of adult trout over eight inches that exist within the site. Estimates are derived from 3-pass backpack electrofishing surveys at 100-meter sites within streams. While all fish in the site are collected and measured, only trout species are used for classifications. Sites are numbered in accordance for the distance (m) from the stream confluence of a higher order stream/river.

| Based on number of trout in excess of eight inches | | | | |
|--|---|--|--|--|
| Class BR1 | Number of wild Brown Trout exceeds 150 per acre | | | |
| Class BR2 | Number of wild Brook Trout ranges from 25 to 150 per acre | | | |
| Class BR3 | Number of wild Brown Trout is less than 25 per acre | | | |
| Class BK1 | Number of wild Brook Trout exceeds 150 per acre | | | |
| Class BK2 | Number of wild Brook Trout ranges from 25 to 150 per acre | | | |
| Class BK3 | Number of wild BrookTrout is less than 25 per acre | | | |
| Class RB1 | Number of wild Rainbow Trout exceeds 25 per acre | | | |
| Class RB2 | Number of wild Rainbow Trout is less than 25 per acre | | | |


Table 2. Species composition and population estimates on Meadow Creek, 1988-1998


Site 12

| Species/Size | Date | Pop Est. | (95% C.I.) | Stream Classification |
|---------------------|--------------|----------|------------|-----------------------|
| Brook Trout <200mm | Jun 29, 1998 | 11 | 11-13 | |
| Brook Trout >=200mm | Jun 29, 1998 | 1 | 1-1 | BK2 |
| Longnose Dace | Jun 13, 1995 | 466 | 411-521 | |
| | Jun 29, 1998 | 645 | 625-665 | |
| Mountain Sucker | Jun 13, 1995 | 19 | 17-25 | |
| | Jun 29, 1998 | 47 | 44-53 | |


Length Frequency Distribution

Length frequency histogram of species sampled by year.














8/12/2020 Page 4