

SOUTH DAKOTA BLACK HILLS COLD STREAM SURVEY
Meeker Creek, Custer County
10120109001108
2015

Stream sites are classified based on the estimates of adult trout over eight inches that exist within the site. Estimates are derived from 3-pass backpack electrofishing surveys at 100-meter sites within streams. While all fish in the site are collected and measured, only trout species are used for classifications. Sites are numbered in accordance for the distance (m) from the stream confluence of a higher order stream/river.

Based on number of trout in excess of eight inches	
Class BR1	Number of wild Brown Trout exceeds 150 per acre
Class BR2	Number of wild Brook Trout ranges from 25 to 150 per acre
Class BR3	Number of wild Brown Trout is less than 25 per acre
Class BK1	Number of wild Brook Trout exceeds 150 per acre
Class BK2	Number of wild Brook Trout ranges from 25 to 150 per acre
Class BK3	Number of wild Brook Trout is less than 25 per acre
Class RB1	Number of wild Rainbow Trout exceeds 25 per acre
Class RB2	Number of wild Rainbow Trout is less than 25 per acre

Table 2. Species composition and population estimates on Meeker Creek, 2005-2015

Site 55

Species/Size	Date	Pop Est.	(95% C.I.)	Stream Classification
Brook Stickleback	May 21, 2009	3	3-3	

Length Frequency Distribution

Length frequency histogram of species sampled by year.

Site: 55

Species: Brook Stickleback

