

**SOUTH DAKOTA BLACK HILLS COLD STREAM SURVEY**  
**Strawberry Creek, Lawrence County**  
**09-0034-00**  
**2015**

Table 1. Population and Biomass Estimates for Strawberry Creek

<b>Site Number:</b> 1	<b>Date Sampled:</b> 06/29/2015	<b>Conductivity (umhos/cm):</b> 487
<b>Site Midpoint (Latitude, Longitude):</b> 44.32376938, -103.65155893	<b>Site Length (m):</b> 100	<b>pH:</b> 7.6
<b>Survey Completed by:</b> South Dakota Game, Fish and Parks	<b>Mean Width (m):</b> 2.1	<b>Water Temperature (C):</b> 10.8
<b>Stream Classification:</b> BK0	<b>Number of Pass:</b> 3	<b>Voltage:</b> 300

Species	Size Class	Total Number Captured	Est. # in site	Upper 95% CI	Lower 95% CI	# per Hectacre	Kg per Hectacre	# per Km	# per Acre	lb. per Acre	# per Mile	Mean Length (mm)	Mean Weight (grams)
Brook Trout	<200mm	35	36	40	35	1714	49	360	694	44.10	579	130.3	28.8
Brook Trout	All	35	36	40	35	1714	49	360	694	44.10	579	130.3	28.8

<b>Site Number:</b> 6	<b>Date Sampled:</b> 06/29/2015	<b>Conductivity (umhos/cm):</b> 487
<b>Site Midpoint (Latitude, Longitude):</b> 44.32378728, -103.65699254	<b>Site Length (m):</b> 100	<b>pH:</b> 7.6
<b>Survey Completed by:</b> South Dakota Game, Fish and Parks	<b>Mean Width (m):</b> 1.9	<b>Water Temperature (C):</b> 10.8
<b>Stream Classification:</b> BK2	<b>Number of Pass:</b> 3	<b>Voltage:</b> 300

Species	Size Class	Total Number Captured	Est. # in site	Upper 95% CI	Lower 95% CI	# per Hectacre	Kg per Hectacre	# per Km	# per Acre	lb. per Acre	# per Mile	Mean Length (mm)	Mean Weight (grams)
Brook Trout	<200mm	37	43	54	37	2242	63	430	908	56.19	692	115.9	28.1
Brook Trout	>=200mm	3	3	3	3	156	15	30	63	13.59	48	212.0	97.3
Brook Trout	All	40	45	54	40	2346	82	450	950	72.84	724	123.1	34.8

Stream sites are classified based on the estimates of adult trout over eight inches that exist within the site. Estimates are derived from 3-pass backpack electrofishing surveys at 100-meter sites within streams. While all fish in the site are collected and measured, only trout species are used for classifications. Sites are numbered in accordance for the distance (m) from the stream confluence of a higher order stream/river.

Based on number of trout in excess of eight inches	
<b>Class BR1</b>	Number of wild Brown Trout exceeds 150 per acre
<b>Class BR2</b>	Number of wild Brook Trout ranges from 25 to 150 per acre
<b>Class BR3</b>	Number of wild Brown Trout is less than 25 per acre
<b>Class BK1</b>	Number of wild Brook Trout exceeds 150 per acre
<b>Class BK2</b>	Number of wild Brook Trout ranges from 25 to 150 per acre
<b>Class BK3</b>	Number of wild Brook Trout is less than 25 per acre
<b>Class RB1</b>	Number of wild Rainbow Trout exceeds 25 per acre
<b>Class RB2</b>	Number of wild Rainbow Trout is less than 25 per acre

Table 2. Species composition and population estimates on Strawberry Creek, 2005-2015

**Site 1**

Species/Size	Date	Pop Est.	(95% C.I.)	Stream Classification
Brook Trout <200mm	Sep 11, 2007	102	96-110	
	Jun 29, 2015	36	35-40	

**Site 6**



Species/Size	Date	Pop Est.	(95% C.I.)	Stream Classification
Brook Trout <200mm	Sep 12, 2007	18	18-18	
	Jun 29, 2015	43	37-54	
Brook Trout >=200mm	Jun 29, 2015	3	3-3	BK2

## Length Frequency Distribution

Length frequency histogram of species sampled by year.


**Site: 1**

**Species: Brook Trout**



**Site: 1**

**Species: Longnose Dace**



Site: 6  
Species: Brook Trout

