

SOUTH DAKOTA BLACK HILLS COLD STREAM SURVEY
Tillson Creek, Lawrence County
09-0081-00
1994

Table 1. Population and Biomass Estimates for Tillson Creek

| | | |
|------------------------------------------------------------------------|---------------------------------|-------------------------------------|
| Site Number: 23 | Date Sampled: 08/02/1994 | Conductivity (umhos/cm): 188 |
| Site Midpoint (Latitude, Longitude): 44.17794910, -103.77710190 | Site Length (m): 100 | pH: 8.6 |
| Survey Completed by: South Dakota Game, Fish and Parks | Mean Width (m): 1.6 | Water Temperature (C): 13.0 |
| Stream Classification: BK2, BR2 | Number of Pass: 3 | Voltage: 400 |

| Species | Size Class | Total Number Captured | Est. # in site | Upper 95% CI | Lower 95% CI | # per Hectacre | Kg per Hectacre | # per Km | # per Acre | lb. per Acre | # per Mile | Mean Length (mm) | Mean Weight (grams) |
|-----------------|------------|-----------------------|----------------|--------------|--------------|----------------|-----------------|----------|------------|--------------|------------|------------------|---------------------|
| Brook Trout | <200mm | 59 | 58 | 81 | 59 | 3718 | 80 | 580 | 1506 | 71.74 | 933 | 120.6 | 21.6 |
| Brook Trout | >=200mm | 3 | 2 | 4 | 3 | 128 | 13 | 20 | 52 | 11.83 | 32 | 212.0 | 103.3 |
| Brook Trout | All | 62 | 61 | 85 | 62 | 3910 | 100 | 610 | 1584 | 89.25 | 981 | 125.0 | 25.6 |
| Brown Trout | <200mm | 2 | 2 | 2 | 2 | 128 | 6 | 20 | 52 | 5.67 | 32 | 174.5 | 49.5 |
| Brown Trout | >=200mm | 2 | 2 | 4 | 2 | 128 | 14 | 20 | 52 | 12.19 | 32 | 214.0 | 106.5 |
| Brown Trout | All | 4 | 4 | 5 | 4 | 256 | 20 | 40 | 104 | 17.86 | 64 | 194.3 | 78.0 |
| Longnose Dace | All | 78 | 117 | 169 | 78 | 7500 | 39 | 1170 | 3038 | 34.72 | 1883 | 84.2 | 5.2 |
| Mountain Sucker | All | 5 | 5 | 5 | 5 | 321 | 6 | 50 | 130 | 4.98 | 80 | 116.0 | 17.4 |
| White Sucker | All | 22 | 22 | 22 | 22 | 1410 | 23 | 220 | 571 | 20.15 | 354 | 114.5 | 16.0 |

Stream sites are classified based on the estimates of adult trout over eight inches that exist within the site. Estimates are derived from 3-pass backpack electrofishing surveys at 100-meter sites within streams. While all fish in the site are collected and measured, only trout species are used for classifications. Sites are numbered in accordance for the distance (m) from the stream confluence of a higher order stream/river.

| Based on number of trout in excess of eight inches | |
|----------------------------------------------------|-----------------------------------------------------------|
| Class BR1 | Number of wild Brown Trout exceeds 150 per acre |
| Class BR2 | Number of wild Brook Trout ranges from 25 to 150 per acre |
| Class BR3 | Number of wild Brown Trout is less than 25 per acre |
| Class BK1 | Number of wild Brook Trout exceeds 150 per acre |
| Class BK2 | Number of wild Brook Trout ranges from 25 to 150 per acre |
| Class BK3 | Number of wild Brook Trout is less than 25 per acre |
| Class RB1 | Number of wild Rainbow Trout exceeds 25 per acre |
| Class RB2 | Number of wild Rainbow Trout is less than 25 per acre |

Table 2. Species composition and population estimates on Tillson Creek, 1984-1994

Site 23


| Species/Size | Date | Pop Est. | (95% C.I.) | Stream Classification |
|---------------------|--------------|----------|------------|-----------------------|
| Brook Trout <200mm | Aug 02, 1994 | 58 | 59-81 | |
| Brook Trout >=200mm | Aug 02, 1994 | 2 | 3-4 | BK2 |
| Brown Trout <200mm | Aug 02, 1994 | 2 | 2-2 | |
| Brown Trout >=200mm | Aug 02, 1994 | 2 | 2-4 | BR2 |
| Longnose Dace | Aug 02, 1994 | 117 | 78-169 | |
| Mountain Sucker | Aug 02, 1994 | 5 | 5-5 | |
| White Sucker | Aug 02, 1994 | 22 | 22-22 | |

Length Frequency Distribution

Length frequency histogram of species sampled by year.


Site: 23

Species: Brook Trout




Site: 23

Species: Brown Trout




Site: 23

Species: Longnose Dace



Site: 23
Species: Mountain Sucker



Site: 23
Species: White Sucker

