

# SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Potts, Potter County

LLO-Lake-2378-000

2020

## Lake Information

**Name:** Potts **Maximum Depth:** 15 Feet  
**County:** Potter **Mean Depth:** 6 Feet  
**Legal Description:** T117-R74-S20  
**Surface Area:** 52 Acres

## Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
boat shocker (night)	Sep 23, 2020	3600 seconds
frame net (std 3/4 in)	Jun 29, 2020	10 net-nights

## **Common Fish Species Present**

Largemouth Bass

Bluegill

Black Bullhead

Yellow Perch

---

## Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- **AFS std gill net** – 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- **std experimental gill net for non-Missouri River waters** - 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- **std experimental gill net for Missouri River reservoirs** – 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$CPUE = \frac{\text{number of fish}}{\text{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left( \frac{\text{number of fish} \geq \text{quality length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

$$PSD - P = \left( \frac{\text{number of fish} \geq \text{preferred length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (**W**) compared to a length-specific standard weight (**Ws**) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left( \frac{W}{W_s} \right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (S), quality (Q), preferred (P), memorable (M) and trophy (T). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

## Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

\* **Methods/Species that ignore stock length**

Gear	Species	Sample Size (n)	Abundance		Stock Density Indices			Condition		
			CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
boat shocker (night)	Largemouth Bass	548	157.0	34.7	7	3	4	2	106	1
frame net (std 3/4 in)	Black Bullhead	1752	173.6	63.8	7	1	0		91	1
	Bluegill	40	4.0	2.9	100		93		116	2
	Largemouth Bass	6	0.0	0.0	0		0			
	Yellow Perch	294	29.0	15.2	23	3	2	1	95	1

## **10-Year Catch Per Unit Effort by Gear and Species**

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

\* **Methods/Species that ignore stock length**

Gear	Species	CPUE										Avg
		2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	
boat shocker (night)	Largemouth Bass	87.0				45.0			15.0		157.0	76.00
frame net (std 3/4 in)	Black Bullhead	10.0		9.8		3.6			485.8		173.6	136.5 6
	Black Crappie	0.1		0.0		0.0			0.0		0.0	0.02
	Bluegill	0.0		0.0		0.1			4.6		4.0	1.74
	Largemouth Bass	0.0		0.0		0.0			0.0		0.0	0.00
	Yellow Perch	10.0		0.0		1.7			13.2		29.0	10.78

## 10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

Gear	Species	Index	Year										
			2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	
boat shocker (night)	Largemouth Bass	PSD	60					0			67		7
		PSD-P	8				0			67		4	
		Wr	111				130			110		106	
frame net (std 3/4 in)	Black Bullhead	PSD	0	96			6				1		7
		PSD-P	0	0			0			1		0	
		Wr	79	95			87			82		91	
	Bluegill	PSD					100				100		100
		PSD-P					0				9		93
		Wr					141			118		116	
	Largemouth Bass	PSD											0
		PSD-P											0
	Yellow Perch	PSD	46					47			67		23
		PSD-P	0					6			5		2
		Wr	94					95			88		95

## **Back-Calculated Lengths**

Mean species back-calculated total length (mm) at age, standard error (SE), and sample size (N).

Species: Bluegill

Year Class	Age	N	Mean back-calculated length (SE) at age											
			1	2	3	4	5	6	7	8	9	10		
2018	2	1	84	124										
2014	6	3	92 (3.6)	131 (4.8)	145 (1.3)	162 (3.3)	177 (2.2)	191 (.7)						
2013	7	5	117 (3.4)	149 (2.3)	164 (2.9)	177 (4.6)	188 (4)	201 (2.9)	211 (2.4)					
2012	8	7	103 (3.2)	130 (5.7)	145 (5.7)	158 (5.8)	175 (4.5)	188 (2.9)	199 (2)	206 (1.7)				
2008	12	1	84	97	105	115	124	145	162	176	190	200		
Weighted Mean		17	103	133	148	162	176	190	201	202	190	200		
Year Class	Age	N	11	12	13	14	15	16	17	18	19	20		
2018	2	1												
2014	6	3												
2013	7	5												
2012	8	7												
2008	12	1	207	214										
Weighted Mean		17	207	214										



Species: Largemouth Bass

Year Class	Age	N	Mean back-calculated length (SE) at age											
			1	2	3	4	5	6	7	8	9	10		
2018	2	33	123 (2.4)	174 (3.9)										
2017	3	9	114 (7.4)	207 (17.5)	270 (23.8)									
2015	5	2	132 (11.6)	208 (1.2)	303 (5.2)	373 (7.7)	400 (8.2)							
2014	6	1	117	253	362	395	421	431						
2013	7	4	102 (17.2)	203 (18.8)	280 (28.7)	331 (18)	374 (13.9)	396 (12.3)	416 (9.7)					
Weighted Mean		49	120	185	282	352	388	403	416					
Year Class	Age	N	11	12	13	14	15	16	17	18	19	20		
2018	2	33												
2017	3	9												
2015	5	2												
2014	6	1												
2013	7	4												
Weighted Mean		49												

Species: Yellow Perch

Year Class	Age	N	Mean back-calculated length (SE) at age																	
			1	2	3	4	5	6	7	8	9	10								
2019	1	8	109 (2.9)																	
2018	2	6	104 (1.6)	152 (2.5)																
2017	3	6	104 (4.2)	143 (5.2)	169 (1.4)															
2016	4	5	93 (4.9)	115 (7)	151 (2.2)	174 (2)														
2015	5	7	93 (1.1)	112 (1.6)	146 (5.8)	169 (3.7)	184 (3.2)													
2014	6	6	95 (3.7)	134 (5.9)	161 (3.3)	179 (3.4)	194 (3.2)	207 (2.5)												
2013	7	4	88 (9.2)	119 (5.3)	144 (4.9)	172 (4.2)	199 (1.9)	213 (3.1)	224 (3.2)											
2012	8	2	108 (4.3)	147 (6.1)	177 (6.2)	193 (4.6)	207 (5.3)	221 (6.5)	234 (4.5)	245 (2.8)										
2011	9	1	120	155	176	187	209	224	238	244	257									
Weighted Mean		45	100	131	157	176	194	212	229	245	257									
Year Class	Age	N	11	12	13	14	15	16	17	18	19	20								
2019	1	8																		
2018	2	6																		
2017	3	6																		
2016	4	5																		
2015	5	7																		
2014	6	6																		
2013	7	4																		
2012	8	2																		
2011	9	1																		
Weighted Mean		45																		

## Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Bluegill

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2020	40		156 (1)				202 (5)	214 (13)	209 (20)		221 (1)
2018	46			170 (7)	177 (10)	179 (11)	195 (13)	197 (3)		230 (1)	

Species: Largemouth Bass

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2020	452		195 (414)	232 (31)		414 (2)	436 (1)	431 (4)			
2018	25	112 (9)	112 (1)		283 (3)		396 (1)	381 (6)	359 (3)	415 (2)	
2015	46	248 (46)									
2011	87	215 (4)		302 (76)		435 (2)	459 (3)	445 (1)		435 (2)	

## **Fish Condition**

Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).



Species	Year	Length Groups							
		S-Q		Q-P		P-M		M	
		N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Bluegill Frame Net	2018	0		42	117 (1.5)	4	120 (1.4)	0	
	2020	0		3	132 (8.3)	37	115 (1.1)	0	
Largemouth Bass Electro Fishing	2018	5	104 (2.3)	0		10	113 (2.1)	0	
	2020	146	103 (0.9)	4	123 (2.1)	7	126 (5.5)	0	

## Length Frequency Distribution

Length frequency histogram of species sampled by year.



Species: Bluegill


Gear: frame net (std 3/4 in)



Species: Largemouth Bass

Gear: boat shocker (night)






## Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.


Species: Bluegill

Gear: frame net (std 3/4 in)



Species: Largemouth Bass

Gear: boat shocker (night)



## **Fish Stocking**

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2009	Largemouth Bass	Juvenile	490
2013	Largemouth Bass	Fingerling	1,750
2014	Bluegill	Adult	250
2014	Largemouth Bass	Fingerling	1,260
2014	Largemouth Bass	Large Fingerling	1,000
2015	Bluegill	Adult	450
2015	Largemouth Bass	Juvenile	292
2016	Largemouth Bass	Adult	148