

# SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Albert, Kingsbury County

MBS-Lake-176-000

2020

## Lake Information

<b>Name:</b>	Albert	<b>Maximum Depth:</b>	13 Feet
<b>County:</b>	Kingsbury	<b>Mean Depth:</b>	9 Feet
<b>Legal Description:</b>	T112-R53W-Sec. 1-3, 10-12, 14-15, 22	<b>OHWM Elevation:</b>	1,653
<b>Surface Area:</b>	3,672 Acres	<b>Outlet Elevation:</b>	1,650

## Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
AFS std gill net	Jun 23, 2020	8 net-nights

## **Common Fish Species Present**

Walleye

Channel Catfish

Yellow Perch

White Sucker

Northern Pike

White Bass

Black Bullhead

Common Carp

---

## Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- **AFS std gill net** – 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- **std experimental gill net for non-Missouri River waters** - 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- **std experimental gill net for Missouri River reservoirs** – 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$CPUE = \frac{\text{number of fish}}{\text{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left( \frac{\text{number of fish} \geq \text{quality length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

$$PSD - P = \left( \frac{\text{number of fish} \geq \text{preferred length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (**W**) compared to a length-specific standard weight (**Ws**) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left( \frac{W}{W_s} \right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (S), quality (Q), preferred (P), memorable (M) and trophy (T). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

## Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

\* **Methods/Species that ignore stock length**

Gear	Species	Sample Size (n)	Abundance		Stock Density Indices			Condition		
			CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
AFS std gill net	Black Bullhead	4	0.5	0.4	25		0			
	Channel Catfish	18	2.3	0.6	89		11		110	3
	Common Carp	2	0.3	0.2	100		50			
	Northern Pike	8	1.0	0.7	50		38		90	4
	Walleye	21	2.5	1.1	80		5		90	2
	White Bass	4	0.5	0.5	25		25		95	2
	White Sucker	9	1.1	0.5	78		78			
	Yellow Perch	12	1.5	0.4	75		42	24	110	3

## 10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

\* Methods/Species that ignore stock length

Gear	Species	CPUE										Avg
		2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	
AFS std gill net	Bigmouth Buffalo							4.3	1.3	0.1	0.0	1.43
	Black Bullhead							1.3	1.1	0.3	0.5	0.80
	Black Crappie							0.5	0.3	0.0	0.0	0.20
	Channel Catfish							0.0	0.1	0.0	2.3	0.60
	Common Carp							0.3	1.0	0.1	0.3	0.43
	Northern Pike							0.7	0.3	0.5	1.0	0.63
	Walleye							8.5	17.9	2.1	2.5	7.75
	White Bass							0.5	0.5	0.6	0.5	0.53
	White Sucker							1.2	1.0	1.0	1.1	1.08
	Yellow Perch							14.5	9.3	5.0	1.5	7.58
frame net (std 3/4 in)	Bigmouth Buffalo		30.6									30.60
	Black Bullhead		57.6									57.60
	Channel Catfish		1.0									1.00
	Common Carp		5.6									5.60
	Northern Pike		4.8									4.80
	Smallmouth Bass		0.2									0.20
	Walleye		4.4									4.40
	White Bass		3.2									3.20
	White Sucker		0.8									0.80
	Yellow Bullhead		3.2									3.20
	Yellow Perch		0.2									0.20
std exp gill net	Bigmouth Buffalo	0.0	0.0		0.0	0.0	10.7					2.14
	Black Bullhead	0.0	3.3		1.7	15.3	4.3					4.92
	Channel Catfish	0.0	0.7		0.0	0.0	0.0					0.14
	Common Carp	0.0	1.0		1.0	0.0	0.0					0.40
	Northern Pike	2.3	4.3		2.0	1.0	1.0					2.12
	Orangespotted Sunfish	0.0	0.0		0.0	0.0	0.0					0.00
	Spottail Shiner	0.0	0.0		0.0	0.0	0.0					0.00
	Walleye	19.7	11.7		7.3	17.0	9.0					12.94
	White Bass	0.0	0.7		0.0	0.0	0.7					0.28
	White Sucker	0.0	1.7		1.3	6.3	3.3					2.52
	Yellow Perch	24.3	13.7		22.0	17.3	114.0					38.26

## 10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

Gear	Species	Index	Year											
			2011	2012	2013	2014	2015	2016	2017	2018	2019	2020		
AFS std gill net	Black Bullhead	PSD								88	100	100	25	
		PSD-P								88	67	100	0	
	Channel Catfish	PSD									100		89	
		PSD-P									100		11	
		Wr									98		110	
	Common Carp	PSD									100	100	100	100
		PSD-P									100	100	100	50
	Northern Pike	PSD									100	100	100	50
		PSD-P									75	50	100	38
		Wr									87	96	97	90
	Walleye	PSD									67	91	76	80
		PSD-P									39	14	6	5
		Wr									86	90	86	90
	White Bass	PSD									67	100	100	25
		PSD-P									67	100	100	25
		Wr									97	106	94	95
	White Sucker	PSD									100	100	100	78
		PSD-P									100	100	100	78
	Yellow Perch	PSD									71	38	43	75
		PSD-P									16	22	5	42
		Wr									98	104	110	110
frame net (std 3/4 in)	Black Bullhead	PSD		73										
		PSD-P		3										
		Wr		100										
	Channel Catfish	PSD		100										
		PSD-P		0										
		Wr		99										
	Common Carp	PSD		100										
		PSD-P		75										
		Wr		88										
	Northern Pike	PSD		58										
		PSD-P		25										
		Wr		76										

Gear	Species	Index	Year										
			2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	
frame net (std 3/4 in)	Walleye	PSD		18									
		PSD-P		0									
		Wr		77									
	White Bass	PSD		31									
		PSD-P		19									
		Wr		88									
	White Sucker	PSD		100									
		PSD-P		100									
		Wr		87									
Yellow Perch	PSD		100										
	PSD-P		0										
	Wr		105										
std exp gill net	Black Bullhead	PSD		80		20	63	100					
		PSD-P		0		20	0	38					
		Wr		98									
	Channel Catfish	PSD		100									
		PSD-P		0									
		Wr		119									
	Common Carp	PSD		100		100							
		PSD-P		33		100							
		Wr		106									
	Northern Pike	PSD	14	54		83	100	67					
		PSD-P	0	8		17	33	33					
		Wr		85		91	106	89					
	Walleye	PSD	7	23		91	94	41					
		PSD-P	0	0		0	24	11					
		Wr		85		96	102	96					
	White Bass	PSD		100				0					
		PSD-P		100				0					
		Wr		100				100					
	White Sucker	PSD		100		25	26	90					
		PSD-P		100		25	16	90					
		Wr		104									
Yellow Perch	PSD	40	71		18	96	13						
	PSD-P	22	34		15	25	11						
	Wr		109		106	121	110						





## Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Walleye

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2019	18	173 (1)	320 (1)	411 (3)	395 (8)	467 (5)					
2018	141		371 (9)	432 (97)	468 (13)	499 (3)		554 (3)		640 (12)	623 (4)

Species: Yellow Perch

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2018	76	143 (48)	227 (12)	269 (15)		327 (1)					

## Fish Condition



Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

Species	Year	Length Groups							
		S-Q		Q-P		P-M		M	
		N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Channel Catfish Gill Net	2018	0		0		1	98	0	
	2020	2	120 (0.3)	14	107 (1.8)	2	119 (13.2)	0	
Northern Pike Gill Net	2016	1	82	1	96	1	90	0	
	2017	0		1	82	3	89 (1.4)	0	
	2018	0		1	91	1	101	0	
	2019	0		0		4	97 (9.5)	0	
	2020	4	86 (2.4)	1	99	3	92 (6.3)	0	
Walleye Gill Net	2016	16	95 (1.2)	8	99 (2.2)	3	94 (3.6)	0	
	2017	17	86 (1.6)	14	85 (3.8)	19	86 (1.6)	1	85
	2018	13	92 (2.0)	110	91 (0.6)	9	91 (1.7)	11	86 (2.3)
	2019	4	87 (3.4)	12	86 (1.9)	1	79	0	
	2020	4	90 (2.6)	15	89 (1.3)	1	102	0	
White Bass Gill Net	2016	2	100 (1.1)	0		0		0	
	2017	1	96	0		2	98 (4.2)	0	
	2018	0		0		1	106	3	107 (3.0)
	2019	0		0		3	94 (1.0)	2	95 (4.6)
	2020	3	95 (1.5)	0		1	93	0	
Yellow Perch Gill Net	2016	299	113 (0.8)	4	97	32	100 (1.6)	7	96 (0.4)
	2017	25	110 (1.4)	48	96 (0.9)	11	96 (2.3)	3	94 (3.2)
	2018	46	107 (1.5)	12	105 (2.9)	13	99 (1.4)	3	99 (5.7)
	2019	23	111 (2.1)	15	110 (1.8)	1	99	1	90
	2020	3	116 (3.0)	4	109 (1.3)	4	111 (3.0)	1	95


## Length Frequency Distribution

Length frequency histogram of species sampled by year.


Species: Black Bullhead  
Gear: std exp gill net






Species: Channel Catfish  
Gear: AFS std gill net





Species: Walleye  
Gear: AFS std gill net










Species: Walleye  
Gear: std exp gill net




Species: White Sucker  
 Gear: std exp gill net





Species: Yellow Perch  
 Gear: AFS std gill net






Species: Yellow Perch  
 Gear: std exp gill net




## Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.

Species: Black Bullhead  
Gear: AFS std gill net




Species: Black Bullhead  
Gear: std exp gill net







Species: Channel Catfish  
Gear: AFS std gill net




Species: Channel Catfish  
Gear: std exp gill net




Species: Common Carp  
Gear: AFS std gill net




Species: Common Carp  
Gear: std exp gill net




Species: Northern Pike  
Gear: AFS std gill net




Species: Northern Pike  
Gear: std exp gill net




Species: Walleye  
Gear: AFS std gill net




Species: Walleye  
Gear: std exp gill net




Species: White Bass  
Gear: AFS std gill net




Species: White Bass  
Gear: std exp gill net




Species: White Sucker  
Gear: AFS std gill net




Species: White Sucker  
Gear: std exp gill net



Species: Yellow Perch  
Gear: AFS std gill net



Species: Yellow Perch  
Gear: std exp gill net



## **Fish Stocking**

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2011	Walleye	Fry	3,700,000
2014	Walleye	Fry	1,850,000
2015	Walleye	Fry	1,850,000
2018	Walleye	Fry	3,700,000