

**SOUTH DAKOTA STATEWIDE FISHERIES SURVEY**  
**Oakwood East, Brookings County**  
**MBS-Lake-215-001**  
**2020**

**Lake Information**

<b>Name:</b>	Oakwood East	<b>Maximum Depth:</b>	9 Feet
<b>County:</b>	Brookings	<b>Mean Depth:</b>	6 Feet
<b>Legal Description:</b>	T111N-R51W-Sec. 4-5, 8-9, 16-27	<b>OHWM Elevation:</b>	1,627
<b>Surface Area:</b>	955 Acres	<b>Outlet Elevation:</b>	1,626

**Surveys and Investigations**

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
AFS std gill net	Jul 27, 2020	6 net-nights

## **Common Fish Species Present**

Yellow Perch

Walleye

Common Carp

White Sucker

Northern Pike

Black Bullhead

---

## Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- **AFS std gill net** – 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- **std experimental gill net for non-Missouri River waters** - 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- **std experimental gill net for Missouri River reservoirs** – 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$CPUE = \frac{\text{number of fish}}{\text{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left( \frac{\text{number of fish} \geq \text{quality length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

$$PSD - P = \left( \frac{\text{number of fish} \geq \text{preferred length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (**W**) compared to a length-specific standard weight (**Ws**) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left( \frac{W}{W_s} \right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (S), quality (Q), preferred (P), memorable (M) and trophy (T). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

## **Catch Summary of Stock Length Fish**

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

\* **Methods/Species that ignore stock length**

Gear	Species	Sample Size (n)	Abundance		Stock Density Indices			Condition		
			CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
AFS std gill net	Black Bullhead	6	0.7	0.5	0		0			
	Common Carp	426	3.5	1.9	0		0			
	Northern Pike	13	1.5	0.6	56		22		100	2
	Walleye	24	4.0	1.3	42	16	21	14	97	2
	White Sucker	13	2.2	1.2	77		38		91	
	Yellow Perch	68	11.3	2.1	21	7	3		98	1

## 10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

\* Methods/Species that ignore stock length

Gear	Species	CPUE										Avg
		2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	
AFS std gill net	Bigmouth Buffalo							0.8	0.0	0.0	0.0	0.20
	Black Bullhead							1.7	1.0	0.5	0.7	0.98
	Common Carp							0.0	0.2	0.0	3.5	0.93
	Northern Pike							0.3	0.2	0.0	1.5	0.50
	Walleye							25.3	49.7	1.2	4.0	20.05
	White Sucker							3.8	1.8	1.0	2.2	2.20
	Yellow Perch							33.8	46.5	12.5	11.3	26.03
frame net (std 3/4 in)	Bigmouth Buffalo		2.2									2.20
	Black Bullhead		48.9									48.90
	Black Crappie		0.0									0.00
	Common Carp		3.2									3.20
	Green Sunfish		7.1									7.10
	Northern Pike		3.0									3.00
	Orangespotted Sunfish		0.0									0.00
	Tadpole Madtom		0.0									0.00
	Walleye		1.6									1.60
	White Sucker		7.4									7.40
	Yellow Bullhead		6.2									6.20
Yellow Perch		7.6									7.60	
std exp gill net	Bigmouth Buffalo	0.0	3.7		0.7	0.3	0.3					1.00
	Black Bullhead	2.0	35.3		0.3	0.0	6.7					8.86
	Common Carp	0.0	3.0		1.3	0.7	0.0					1.00
	Green Sunfish	0.0	0.7		0.0	0.0	0.0					0.14
	Northern Pike	4.7	2.7		1.3	4.0	2.0					2.94
	Orangespotted Sunfish	0.0	0.0		0.0	0.0	0.0					0.00
	Walleye	3.3	5.7		25.7	47.3	51.7					26.74
	White Sucker	1.3	12.7		7.7	7.3	2.3					6.26
	Yellow Bullhead	0.3	1.7		0.0	0.0	0.0					0.40
Yellow Perch	13.7	61.3		32.7	5.0	89.7					40.48	

## 10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

Gear	Species	Index	Year										
			2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	
AFS std gill net	Black Bullhead	PSD								50	67	33	0
		PSD-P								0	17	0	0
	Common Carp	PSD									0	0	0
		PSD-P									0	0	0
	Northern Pike	PSD								100	100	0	56
		PSD-P								0	0	0	22
		Wr								86	83		100
	Walleye	PSD								89	25	100	42
		PSD-P								11	11	14	21
		Wr								91	92	97	97
	White Sucker	PSD								100	82	100	77
		PSD-P								100	73	100	38
		Wr											91
	Yellow Perch	PSD								37	18	24	21
		PSD-P								21	3	3	3
Wr									96	100	103	98	
frame net (std 3/4 in)	Black Bullhead	PSD		11									
		PSD-P		0									
		Wr		95									
	Common Carp	PSD		41									
		PSD-P		21									
		Wr		110									
	Northern Pike	PSD		67									
		PSD-P		26									
		Wr		91									
	Walleye	PSD		86									
		PSD-P		7									
		Wr		100									
	White Sucker	PSD		90									
		PSD-P		36									
		Wr		92									
Yellow Perch	PSD		53										
	PSD-P		9										

Gear	Species	Index	Year											
			2011	2012	2013	2014	2015	2016	2017	2018	2019	2020		
frame net (std 3/4 in)	Yellow Perch	Wr		99										
std exp gill net	Black Bullhead	PSD	0	24		0		30						
		PSD-P	0	0		0		0						
		Wr	88	103										
	Common Carp	PSD		0		100	100							
		PSD-P		0		100	100							
		Wr		112										
	Northern Pike	PSD	71	38		100	92	67						
		PSD-P	14	0		75	42	0						
		Wr	97	95		88	87	81						
	Walleye	PSD	10	76		25	25	70						
		PSD-P	10	0		1	0	0						
		Wr	103	95		93	87	78						
	White Sucker	PSD	100	87		70	73	100						
		PSD-P	100	8		26	55	57						
		Wr	108	89										
	Yellow Perch	PSD	5	58		0	73	32						
		PSD-P	0	5		0	20	1						
		Wr	96	102		108	107	87						



## **Fish Condition**


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

Species	Year	Length Groups							
		S-Q		Q-P		P-M		M	
		N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Northern Pike Gill Net	2016	2	91 (12.6)	4	75 (2.7)	0		0	
	2017	0		2	86 (0.8)	0		0	
	2018	0		1	83	0		0	
	2019	0		0		0		0	
	2020	4	100 (2.1)	3	99 (1.8)	2	100 (6.6)	0	
Walleye Gill Net	2016	46	80 (1.1)	109	78 (0.4)	0		0	
	2017	17	91 (0.9)	118	92 (0.6)	17	89 (1.0)	0	
	2018	223	94 (4.1)	43	88 (0.9)	32	89 (1.0)	0	
	2019	0		6	98 (2.8)	1	92	0	
	2020	14	96 (1.5)	5	94 (5.8)	5	103 (2.8)	0	
White Sucker Gill Net	2020	3		5	91	2		3	
Yellow Perch Gill Net	2016	184	89 (0.7)	83	85 (1.0)	2		0	
	2017	128	95 (0.7)	33	97 (1.0)	42	99 (0.9)	0	
	2018	228	101 (0.8)	43	99 (1.5)	8	82	0	
	2019	57	105 (1.1)	16	100 (1.8)	2	97 (1.9)	0	
	2020	54	99 (0.9)	12	96 (1.3)	2	96 (1.8)	0	


## Length Frequency Distribution

Length frequency histogram of species sampled by year.



Species: Black Bullhead  
Gear: AFS std gill net




Species: Black Bullhead  
Gear: std exp gill net




Species: Common Carp  
Gear: AFS std gill net





Species: Northern Pike  
Gear: AFS std gill net





Species: Northern Pike  
Gear: std exp gill net





Species: Walleye  
Gear: AFS std gill net








Species: Walleye  
Gear: std exp gill net




Species: White Sucker  
Gear: AFS std gill net








Species: White Sucker  
 Gear: std exp gill net




Species: Yellow Perch  
 Gear: AFS std gill net






2019




2020

Species: Yellow Perch  
Gear: std exp gill net



2015




2016


## Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.


Species: Black Bullhead  
Gear: AFS std gill net




Species: Black Bullhead  
Gear: std exp gill net



Species: Common Carp  
Gear: AFS std gill net




Species: Common Carp  
Gear: std exp gill net







Species: Northern Pike  
Gear: AFS std gill net




Species: Northern Pike  
Gear: std exp gill net




Species: Walleye  
Gear: AFS std gill net




Species: Walleye  
Gear: std exp gill net




Species: White Sucker  
Gear: AFS std gill net




Species: White Sucker  
Gear: std exp gill net



Species: Yellow Perch  
Gear: AFS std gill net



Species: Yellow Perch  
Gear: std exp gill net



## **Fish Stocking**

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2010	Walleye	Fry	1,000,000
2012	Walleye	Fry	500,000
2013	Walleye	Fry	650,000
2014	Walleye	Fry	453,750
2019	Walleye		933,000
2019	Yellow Perch	Juvenile	47,000