

Creel and Lake Information

Lake Information

Lake Name: Enemy Swim
Lake Code: UBS-Lake-196-000
Surface Area: 884.65 Hectares
County: Day
Start Date: 1/3/2020
End Date: 12/31/2020
Months Included: January, February, March, May, June, July, August, December

Table 1. Percent of angling parties interviewed primarily targeting a species (or any species) by month and overall for Enemy Swim during 2019 - 2020. Any Species (ANY), Black Crappie (BLC), Bluegill (BLG), Northern Pike (NOP), Walleye (WAE), Yellow Perch (YEP).

Year	Month	# of Interviews	Percent of Anglers Targeting					
			ANY	BLC	BLG	NOP	WAE	YEP
2019	January	8	37.5	0.0	50.0	12.5	0.0	0.0
	February	11	18.2	0.0	63.6	18.2	0.0	0.0
	March	9	0.0	11.1	55.6	22.2	0.0	0.0
	May	3	33.3	0.0	0.0	0.0	66.7	0.0
	June	11	36.4	0.0	18.2	0.0	9.1	0.0
	July	13	15.4	0.0	7.7	7.7	53.8	0.0
	August	6	50.0	0.0	16.7	0.0	33.3	0.0
	December	19	10.5	10.5	57.9	15.8	5.3	0.0
	Overall	80	21.3	3.8	38.8	11.3	16.3	0.0
2020	January	18	22.2	0.0	16.7	50.0	0.0	11.1
	February	24	29.2	0.0	41.7	20.8	0.0	8.3
	March	15	13.3	0.0	60.0	20.0	0.0	6.7
	December	10	20.0	0.0	70.0	0.0	0.0	10.0
	Overall	67	22.4	0.0	43.3	25.4	0.0	9.0

Table 2. Monthly and overall number of interviews, angler hours, angler days, mean trip length, mean party size, percent of parties interviewed that were South Dakota residents, percent of angling activity from a boat or shack and percent of angling activity that occurred on weekends and holidays at Enemy Swim (80% confidence interval).

Year	Month	# Interviews	Angler Hours	Angler Days	Trip Length (h)	Party Size	% SD Residents	% Boat or Shack	% Weekend/Holidays
2018-2019	December	14	1,091 (544)	472	2.31 (0.67)	1.47 (0.20)	57.14	86	57
	January	8	1,109 (340)	291	3.81	1.32	87.50	100	75
	February	11	1,892 (984)	563	3.36	1.70	9.09	64	82
	March	9	1,693 (545)	508	3.33 (0.28)	2.51 (0.10)	66.67	56	67
	Overall	42	5,785 (1,295)	1,808	3.20 (0.19)	1.75 (0.06)	52.38	76	69
2019	May	3	3,118 (3,587)	962	3.24	2.00	100.00	100	33
	June	11	5,708 (1,841)	1,735	3.29	2.17 (0.38)	54.55	91	55
	July	13	6,722 (2,191)	1,500	4.48	2.25 (0.36)	30.77	92	62
	August	5	2,412	693	3.48	3.00	66.67	100	17
	Overall	32	17,960 (4,589)	4,934	3.64	2.31 (0.14)	51.52	94	48
2019-2020	December	19	1,004 (492)	407	2.47	2.07 (0.16)	73.68	53	84
	January	18	1,893 (1,129)	295	6.41	1.86 (0.08)	22.22	78	28
	February	24	1,436 (505)	435	3.30 (1.31)	2.01 (0.74)	58.33	79	42
	March	15	5,364 (3,020)	1,812	2.96 (0.97)	1.39 (0.42)	46.67	40	53
	Overall	76	9,697 (3,301)	2,436	3.98 (0.39)	1.83 (0.21)	51.32	64	51

Table 3. Estimated fishing pressure (angler hours) by month during the summer season 2020 daylight period on Enemy Swim (80% confidence interval).

Lake	Month
	Total
Enemy Swim (Day)	0

Table 4. Estimated number of fish harvested, by species and month, for the summer season 2020 daylight period on Enemy Swim (80% confidence interval).

Species	Month
	Total
Black Crappie	0
Bluegill	0
Northern Pike	0
Walleye	0
Yellow Perch	0
Total	0

Table 5a. Estimated monthly and overall angler catch rate per hour fished (C/h) for all anglers by species at Enemy Swim (80% confidence interval). Abbreviations for species are as follows: Black Crappie (BLC), Bluegill (BLG), Northern Pike (NOP), Walleye (WAE), Yellow Perch (YEP) and other species (Other).

Year	Month	BLC	BLG	NOP	WAE	YEP
		C/h	C/h	C/h	C/h	C/h
2018-2019	December	0.01 (0.01)	2.98 (1.67)	0.03 (0.03)	0.00	0.69
	January	0.00	2.84 (2.51)	0.07	0.00	2.75 (1.52)
	February	0.19 (0.10)	5.92 (2.46)	0.07 (0.07)	0.00	0.50 (0.46)
	March	0.51 (0.30)	2.70 (1.35)	0.00	0.01 (0.02)	2.42 (0.86)
	Overall	0.22 (0.09)	3.83 (0.90)	0.04 (0.02)	<0.01 (<0.01)	1.53 (0.42)
2019	May	0.00	0.00	0.00	0.16	0.00
	June	0.00	0.08 (0.09)	0.03	0.00	0.01 (0.01)
	July	0.01 (<0.01)	0.03 (0.03)	0.01 (0.02)	0.18 (0.15)	0.01 (0.01)
	August	0.00	0.62	0.00	0.02	0.00
	Overall	<0.01 (<0.01)	0.12 (0.04)	0.01 (<0.01)	0.10 (0.06)	0.01 (<0.01)
2019-2020	December	0.44 (0.56)	2.96	0.02 (0.01)	0.01 (0.02)	0.29 (0.43)
	January	0.00	0.83 (0.78)	0.04 (0.05)	0.08 (0.05)	0.85 (0.51)
	February	0.00	0.54 (0.88)	0.01 (0.02)	0.00	0.61 (0.22)
	March	<0.01	1.40 (0.89)	0.01 (0.01)	0.00	1.67 (1.47)
	Overall	0.05 (0.06)	1.32 (0.56)	0.02 (0.01)	0.02 (<0.01)	1.21 (0.69)

Table 5b. Estimated monthly and overall angler harvest rate per hour fished (H/h) for all anglers by species at Enemy Swim (80% confidence interval). Abbreviations for species are as follows: Black Crappie (BLC), Bluegill (BLG), Northern Pike (NOP), Walleye (WAE), Yellow Perch (YEP) and other species (Other).

Year	Month	BLC	BLG	NOP	WAE	YEP
		H/h	H/h	H/h	H/h	H/h
2018- 2019	December	0.01 (0.01)	0.29 (0.16)	0.02 (0.02)	0.00	0.03 (0.03)
	January	0.00	0.23 (0.30)	0.02	0.00	0.06 (0.10)
	February	0.00	0.20 (0.12)	0.07 (0.07)	0.00	0.05 (0.02)
	March	0.05 (0.01)	0.04 (0.01)	0.00	0.01 (0.02)	0.00
	Overall	0.02 (<0.01)	0.18 (0.07)	0.03 (0.02)	<0.01 (<0.01)	0.03 (0.02)
2019	May	0.00	0.00	0.00	0.08	0.00
	June	0.00	0.01 (0.01)	0.00	0.00	0.00
	July	0.01 (<0.01)	0.00	0.00	0.13 (0.13)	0.00
	August	0.00	0.30	0.00	0.02	0.00
	Overall	<0.01 (<0.01)	0.04 (0.01)	0.00	0.07 (0.05)	0.00
2019- 2020	December	0.26 (0.43)	0.53 (0.35)	0.02 (0.01)	0.00	0.00
	January	0.00	0.21 (0.26)	0.04 (0.05)	0.07 (0.04)	0.04 (0.06)
	February	0.00	0.00	0.01 (0.02)	0.00	0.00
	March	<0.01	0.06 (0.04)	0.01 (0.01)	0.00	0.03 (0.02)
	Overall	0.03 (0.04)	0.13 (0.07)	0.02 (0.01)	0.01 (<0.01)	0.03 (0.01)

Table 6a. Estimated monthly and overall angler catch rate per hour fished (C/h) only for anglers primarily targeting a species at Enemy Swim. (80% confidence interval). Abbreviations for species are as follows: Black Crappie (BLC), Bluegill (BLG), Northern Pike (NOP), Walleye (WAE), Yellow Perch (YEP) and Other Species (Other).

Year	Month	BLC	BLG	NOP	WAE	YEP
		C/h	C/h	C/h	C/h	C/h
2018- 2019	December	0.00	3.52 (0.37)	0.04 (0.12)	0.00	0.00
	January	0.00	0.92	0.30	0.00	0.00
	February	0.00	13.99	0.32	0.00	0.00
	March	0.00	3.93 (2.45)	0.00	0.00	0.00
	Overall	0.00	5.38 (0.63)	0.16 (0.05)	0.00	0.00
2019	May	0.00	0.00	0.00	0.19	0.00
	June	0.00	0.94	0.00	0.00	0.00
	July	0.00	0.00	0.00	0.34 (0.61)	0.00
	August	0.00	0.88	0.00	0.00	0.00
	Overall	0.00	0.78	0.00	0.16 (0.20)	0.00
2019- 2020	December	5.42	3.14 (1.68)	0.16 (0.22)	0.00	0.00
	January	0.00	1.51 (0.59)	0.26	0.00	1.25
	February	0.00	1.09 (1.26)	0.14	0.00	0.00
	March	0.00	5.17 (0.72)	0.03	0.00	0.41
	Overall	5.42	2.78 (0.63)	0.13 (0.02)	0.00	0.56

Table 6b. Estimated monthly and overall angler harvest rate per hour fished (H/h) only for anglers primarily targeting a species at Enemy Swim. (80% confidence interval). Abbreviations for species are as follows: Black Crappie (BLC), Bluegill (BLG), Northern Pike (NOP), Walleye (WAE), Yellow Perch (YEP) and Other Species (Other).

Year	Month	BLC	BLG	NOP	WAE	YEP
		H/h	H/h	H/h	H/h	H/h
2018-2019	December	0.00	0.34	0.04 (0.07)	0.00	0.00
	January	0.00	0.03	0.30	0.00	0.00
	February	0.00	0.11	0.32	0.00	0.00
	March	0.00	0.06	0.00	0.00	0.00
	Overall	0.00	0.14	0.16 (0.03)	0.00	0.00
2019	May	0.00	0.00	0.00	0.09	0.00
	June	0.00	0.16	0.00	0.00	0.00
	July	0.00	0.00	0.00	0.26 (0.41)	0.00
	August	0.00	0.29	0.00	0.00	0.00
	Overall	0.00	0.18	0.00	0.11 (0.13)	0.00
2019-2020	December	4.27	1.03 (0.71)	0.16 (0.22)	0.00	0.00
	January	0.00	0.45 (0.57)	0.26	0.00	0.00
	February	0.00	0.00	0.14	0.00	0.00
	March	0.00	0.60	0.03	0.00	0.00
	Overall	4.27	0.56 (0.26)	0.13 (0.02)	0.00	0.00

Table 7a. Estimated monthly and overall angler catch (C) for all anglers by species at Enemy Swim (80% confidence interval). Abbreviations for species are as follows: Black Crappie (BLC), Bluegill (BLG), Northern Pike (NOP), Walleye (WAE), Yellow Perch (YEP) and other species (Other).

Year	Month	BLC	BLG	NOP	WAE	YEP
		C	C	C	C	C
2018- 2019	December	16 (14)	3,254 (1,390)	36 (23)	0	753 (177)
	January	0	3,154 (2,610)	78 (36)	0	3,043 (1,407)
	February	364 (36)	11,194 (684)	126 (109)	0	942 (715)
	March	870 (421)	4,570 (1,755)	0	21 (42)	4,096 (507)
	Overall	1,250 (423)	22,173 (3,506)	240 (117)	21 (42)	8,835 (1,667)
	2019	May	0	0	0	500
	June	0	435 (467)	145	0	72 (78)
	July	38 (39)	189 (171)	75 (124)	1,242 (974)	75 (77)
	August	0	1,488	0	43	0
	Overall	38 (39)	2,112 (497)	220 (124)	1,785 (974)	148 (110)
2019- 2020	December	442 (524)	2,977 (1,693)	19 (20)	14 (22)	291 (411)
	January	0	1,575 (779)	84 (77)	153 (21)	1,607 (212)
	February	0	772 (790)	15 (27)	0	871 (794)
	March	13	7,488 (3,694)	64 (61)	0	8,961 (3,126)
	Overall	455 (524)	12,812 (4,212)	182 (104)	168 (31)	11,730 (3,258)

Table 7b. Estimated monthly and overall angler harvest (H) for all anglers by species at Enemy Swim (80% confidence interval). Abbreviations for species are as follows: Black Crappie (BLC), Bluegill (BLG), Northern Pike (NOP), Walleye (WAE), Yellow Perch (YEP) and other species (Other).

Year	Month	BLC	BLG	NOP	WAE	YEP
		H	H	H	H	H
2018- 2019	December	16 (14)	318 (163)	24 (12)	0	28 (16)
	January	0	253 (324)	26 (55)	0	67 (108)
	February	0	377 (36)	126 (109)	0	90
	March	77	69	0	21 (42)	0
	Overall	93 (14)	1,016 (364)	176 (123)	21 (42)	185 (109)
2019	May	0	0	0	250	0
	June	0	72 (78)	0	0	0
	July	38 (39)	0	0	902 (811)	0
	August	0	723	0	43	0
	Overall	38 (39)	795 (78)	0	1,195 (811)	0
2019- 2020	December	265 (412)	534 (370)	19 (20)	0	0
	January	0	400 (187)	84 (77)	131 (17)	79 (55)
	February	0	0	15 (27)	0	0
	March	13	340	64 (61)	0	176
	Overall	278 (412)	1,275 (414)	182 (104)	131 (17)	255 (55)

Table 8. Estimated number and rate (number per hour) of Black Crappie (BLC), Bluegill (BLG), Northern Pike (NOP), Walleye (WAE), Yellow Perch (YEP) caught and harvested by anglers during the summer and winter fisheries on Enemy Swim, (80% confidence interval).

Black Crappie

Season	Year	Catch	Harvest	Catch Rate	Harvest Rate
Summer	2011	136	136	0.01	0.01
Summer	2013	2,018 (655)	1,254 (513)	0.04 (0.02)	0.03 (0.02)
Summer	2014	2,238 (1,008)	1,883 (951)	0.04 (0.03)	0.03 (0.03)
Summer	2015	117 (70)	117 (70)	0.01 (<0.01)	0.01 (<0.01)
Summer	2016	1,650 (277)	381 (284)	0.04 (0.01)	0.01 (<0.01)
Summer	2017	42 (42)	27 (35)	<0.01 (<0.01)	<0.01 (<0.01)
Summer	2019	38 (39)	38 (39)	<0.01 (<0.01)	<0.01 (<0.01)
Winter	2013-2014	811	690	0.04	0.03
Winter	2014-2015	2,530	2,255	0.11	0.10
Winter	2015-2016	984	804	0.05	0.04
Winter	2016-2017	1,432	208	0.12	0.02
Winter	2018-2019	1,250 (423)	93 (14)	0.22 (0.09)	0.02 (<0.01)
Winter	2019-2020	455 (524)	278 (412)	0.05 (0.06)	0.03 (0.04)

Bluegill

Season	Year	Catch	Harvest	Catch Rate	Harvest Rate
Summer	2011	10,089 (4,497)	2,976 (442)	0.53 (0.27)	0.16 (0.05)
Summer	2013	59,846 (13,046)	33,680 (6,987)	1.31 (0.29)	0.74 (0.18)
Summer	2014	32,097 (11,679)	22,492 (7,689)	0.59 (0.41)	0.41 (0.27)
Summer	2015	8,718 (2,360)	5,668 (1,409)	0.45 (0.22)	0.29 (0.15)
Summer	2016	27,971 (12,260)	16,519 (6,459)	0.71 (0.44)	0.42 (0.23)
Summer	2017	6,892 (4,505)	1,194 (554)	0.37 (0.24)	0.06 (0.03)
Summer	2019	2,112 (497)	795 (78)	0.12 (0.04)	0.04 (0.01)
Winter	2010-2011	187	94	0.46	0.23
Winter	2013-2014	30,661	17,149	1.48	0.83
Winter	2014-2015	13,148	8,352	0.56	0.36
Winter	2015-2016	30,060	10,252	1.51	0.51
Winter	2016-2017	46,056	7,268	3.98	0.63
Winter	2018-2019	22,173 (3,506)	1,016 (364)	3.83 (0.90)	0.18 (0.07)
Winter	2019-2020	12,812 (4,212)	1,275 (414)	1.32 (0.56)	0.13 (0.07)

Northern Pike

Season	Year	Catch	Harvest	Catch Rate	Harvest Rate
Summer	2011	1,541 (88)	139 (74)	0.08 (0.02)	0.01 (<0.01)
Summer	2013	1,698 (772)	50 (65)	0.04 (0.02)	<0.01 (<0.01)
Summer	2014	2,590 (869)	490 (502)	0.05 (0.02)	0.01 (<0.01)
Summer	2015	1,168 (918)	104 (67)	0.06 (0.04)	0.01 (<0.01)
Summer	2016	1,083 (426)	153 (119)	0.03 (0.01)	<0.01 (<0.01)
Summer	2017	1,445 (919)	127 (86)	0.08 (0.05)	0.01 (<0.01)
Summer	2019	220 (124)	0	0.01 (<0.01)	0.00
Winter	2010-2011	16	16	0.04	0.04
Winter	2013-2014	204	96	0.01	<0.01
Winter	2014-2015	249	53	0.01	<0.01
Winter	2015-2016	601	524	0.03	0.03
Winter	2016-2017	1,048	285	0.09	0.02
Winter	2018-2019	240 (117)	176 (123)	0.04 (0.02)	0.03 (0.02)
Winter	2019-2020	182 (104)	182 (104)	0.02 (0.01)	0.02 (0.01)

Walleye

Season	Year	Catch	Harvest	Catch Rate	Harvest Rate
Summer	2011	2,924 (309)	0	0.15 (0.03)	0.00
Summer	2013	11,166 (2,151)	1,862 (262)	0.25 (0.08)	0.04 (<0.01)
Summer	2014	26,675 (9,103)	3,236 (964)	0.49 (0.26)	0.06 (0.02)
Summer	2015	4,012 (2,223)	554 (274)	0.21 (0.22)	0.03 (0.03)
Summer	2016	5,747 (2,282)	1,990 (717)	0.15 (0.08)	0.05 (0.03)
Summer	2017	2,347 (765)	1,364 (515)	0.13 (0.05)	0.07 (0.03)
Summer	2019	1,785 (974)	1,195 (811)	0.10 (0.06)	0.07 (0.05)
Winter	2013-2014	721	65	0.03	<0.01
Winter	2014-2015	1,883	435	0.08	0.02
Winter	2015-2016	463	53	0.02	<0.01
Winter	2016-2017	108	72	0.01	0.01
Winter	2018-2019	21 (42)	21 (42)	<0.01 (<0.01)	<0.01 (<0.01)
Winter	2019-2020	168 (31)	131 (17)	0.02 (<0.01)	0.01 (<0.01)

Yellow Perch

Season	Year	Catch	Harvest	Catch Rate	Harvest Rate
Summer	2011	2,904 (836)	68	0.15 (0.01)	<0.01 (<0.01)
Summer	2013	3,204 (1,313)	239 (64)	0.07 (0.04)	0.01 (<0.01)
Summer	2014	1,721 (937)	455 (459)	0.03 (0.03)	0.01 (0.01)
Summer	2015	951 (204)	61 (65)	0.05 (0.02)	<0.01 (<0.01)
Summer	2016	5,142 (2,747)	26 (24)	0.13 (0.08)	<0.01 (<0.01)
Summer	2017	6,215 (981)	25 (32)	0.34 (0.07)	<0.01 (<0.01)
Summer	2019	148 (110)	0	0.01 (<0.01)	0.00
Winter	2013-2014	3,289	57	0.16	<0.01
Winter	2014-2015	1,852	98	0.08	<0.01
Winter	2015-2016	12,550	156	0.63	0.01
Winter	2016-2017	11,372	121	0.98	0.01
Winter	2018-2019	8,835 (1,667)	185 (109)	1.53 (0.42)	0.03 (0.02)
Winter	2019-2020	11,730 (3,258)	255 (55)	1.21 (0.69)	0.03 (0.01)

Table 9. The number of fish harvested with each species by length group (mm) for the winter and summer fisheries at Enemy Swim. Abbreviations for species are as follows: Black Crappie (BLC), Bluegill (BLG), Northern Pike (NOP), Walleye (WAE), Yellow Perch (YEP).

Winter 2018-2019					
Length Group (mm)	BLC (N=3)	BLG (N=46)	NOP (N=7)	WAE (N=0)	YEP (N=7)
150		1			
160		8			
170		7			
180		13			1
190		8			1
200		4			
210		3			3
220		1			
230					
240		1			
250	1				2
260	2				
530			1		
540					
550					
560					
570					
580			1		
590					
600					
610			1		
620					
630					
640			1		
650					
660					
670					
680					
690			1		
700			1		
710					
720					
730					
740					
750					
760					
770					
780					
790					
800					
810					
820					
830					
840					
850					
860					
870					
880					
890			1		

Summer 2019

Length Group (mm)	BLC (N=1)	BLG (N=11)	NOP (N=0)	WAE (N=16)	YEP (N=0)
150		1			
160		2			
170					
180		2			
190		1			
200		2			
210					
220					
230					
240					
250		2			
260					
270		1			
320	1				
370				1	
380				1	
390				1	
400				2	
410				3	
420					
430				1	
440					
450					
460				3	
470				1	
480				2	
490					
500					
510					
520					
530					
540					
550					
560					
570					
580				1	

Winter 2019-2020

Length Group (mm)	BLC (N=15)	BLG (N=77)	NOP (N=14)	WAE (N=8)	YEP (N=14)
150		2			
160		3			
170		10			
180		13			1
190		19			1
200		10			3
210	1	13			6
220		4			2
230		1			
240	1	1			
250	2				1
260	2				
270		1		1	
280					
290					
300					
310					
320	1				
330	2			1	
340	4			1	
350	1				
360	1				
370					
380					
390					
400				1	
410					
420					
430					
440				1	
450					
460					
470					
480				2	
490					
500				1	
570			1		
580			1		
590			2		
600			1		
610			1		
620			1		
630			1		
640			1		
650					
660					
670					
680			1		
690			2		
700					
710					
720			1		
730			1		

Figure 1a. Length frequency of angler-harvested Black Crappie measured by the creel clerk during creel surveys on Enemy Swim.







Figure 1b. Length frequency of angler-harvested Bluegill measured by the creel clerk during creel surveys on Enemy Swim.



Winter 2018-2019
N = 46
Mean TL = 181 mm



Summer 2019
N = 11
Mean TL = 199 mm



Winter 2019-2020
N = 77
Mean TL = 191 mm

Figure 1c. Length frequency of angler-harvested Northern Pike measured by the creel clerk during creel surveys on Enemy Swim.




Figure 1d. Length frequency of angler-harvested Walleye measured by the creel clerk during creel surveys on Enemy Swim.




Figure 1e. Length frequency of angler-harvested Yellow Perch measured by the creel clerk during creel surveys on Enemy Swim.




Figure 2. South Dakota county of residence for winter and summer resident anglers fishing Enemy Swim, South Dakota between January 2019 and December 2020.




Table 10. Number of parties interviewed that harvested (or caught) a specified number of fish totaled over selected strata for a species.

Angler Harvest	Counts				
	BLC (n=33)	BLG (n=33)	NOP (n=33)	WAE (n=33)	YEP (n=33)
0.00	32	29	33	28	33
0.33	1	0	0	1	0
0.50	0	1	0	1	0
0.67	0	1	0	0	0
1.25	0	1	0	0	0
2.00	0	0	0	2	0
2.50	0	1	0	0	0
6.00	0	0	0	1	0

Winter 2018-2019

Counts

Angler Harvest	BLC (n=42)	BLG (n=42)	NOP (n=42)	WAE (n=42)	YEP (n=42)
0.00	39	28	37	41	36
0.17	0	1	0	0	0
0.33	0	1	0	0	0
0.50	1	2	1	0	1
1.00	1	1	2	1	2
1.50	0	0	1	0	0
2.00	1	2	1	0	2
2.50	0	1	0	0	0
4.50	0	1	0	0	0
5.00	0	1	0	0	1
10.00	0	2	0	0	0
15.00	0	2	0	0	0

Winter 2019-
2020

Counts

Angler Harvest	BLC (n=76)	BLG (n=76)	NOP (n=76)	WAE (n=76)	YEP (n=76)
0.00	73	61	64	70	71
0.20	0	0	0	1	0
0.25	0	0	0	1	0
0.50	0	1	1	2	0
0.67	0	1	0	0	0
1.00	1	1	4	1	0
1.50	0	0	0	1	1
2.00	0	1	3	0	1
3.00	0	1	2	0	1
4.00	0	1	2	0	1
5.00	0	1	0	0	1
7.00	1	0	0	0	0
8.00	0	2	0	0	0
10.00	0	1	0	0	0
11.00	0	1	0	0	0
11.50	0	1	0	0	0
12.00	0	2	0	0	0
15.00	1	1	0	0	0

Table 11 The number (N) and percent of total interviews in supplemental-response categories by question asked during interviews at Enemy Swim.

Question: Considering all factors, how satisfied are you with your fishing trip today

Response	Summer 2019 (n = 31)
Very Satisfied	42%
Moderately Satisfied	16%
Slightly Satisfied	13%
Neutral/No Opinion	19%
Slightly Dissatisfied	6%
Moderately Dissatisfied	3%
Very Dissatisfied	0%