

## Dry 2 Survey Summary

Dry 2, located 2.5 miles west and 2.5 miles north of Willow Lake, is managed as a walleye and yellow perch fishery, but other fish species (e.g., northern pike) are present and contribute to the fishery.

- **Walleye.** Walleye numbers were considerably lower in 2021 than in 2018. At 5.8 per gill net, relative abundance was considered low to moderate for Dry 2. Sampled walleyes ranged in length from 10.2 to 27.2 inches, 63% were  $\geq 15.0$  inches and 37% were  $\geq 20.0$  inches. Fourteen year classes contributed to the catch, most (13 of 14) were represented by eight or fewer individuals. Walleyes from the naturally produced 2019 (age-2) cohort were the most abundant accounting for 24 of the 70 fish in the sample. The 2021 sample suggests good walleye growth with mean length at capture values of 14.6 inches at age 2 and 18.7 inches at age 3.
- **Yellow Perch.** Yellow perch were the most abundant species in the 2021 gill net catch. The mean gill net CPUE was 9.2 and suggested moderate relative abundance. Sampled yellow perch ranged in length from 5.1 to 13.8 inches, 91% were  $\geq 8.0$  inches and 75% were  $\geq 10.0$  inches. Five year classes (2013 and 2017 – 2020) were present. Individuals from the 2018 (age 3) cohort, which had a mean length at capture of 10.4 inches, were the most numerous accounting for three-fourths (75%) of yellow perch in the sample.

For more detailed results see the computer-generated South Dakota Statewide Fisheries Survey for Dry 2 (below).

# SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Dry 2, Clark County

MBS-Lake-115-003

2021

## Lake Information

**Name:** Dry 2 **Maximum Depth:** 14 Feet  
**County:** Clark  
**Surface Area:** 9,268 Acres

## Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
AFS std gill net	Jul 27, 2021	6 net-nights
AFS std gill net	Jul 29, 2021	6 net-nights

## **Common Fish Species Present**

Yellow Perch

Walleye

Northern Pike

Common Carp

---

## Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- **AFS std gill net** – 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- **std experimental gill net for non-Missouri River waters** - 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- **std experimental gill net for Missouri River reservoirs** – 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$CPUE = \frac{\text{number of fish}}{\text{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left( \frac{\text{number of fish} \geq \text{quality length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

$$PSD - P = \left( \frac{\text{number of fish} \geq \text{preferred length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (*W*) compared to a length-specific standard weight (*Ws*) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left( \frac{W}{W_s} \right) \times 100$$

Confidence intervals (**CI**) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (**S**), quality (**Q**), preferred (**P**), memorable (**M**) and trophy (**T**). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

## Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

Gear	Species	Sample Size (n)	Abundance		Stock Density Indices			Condition		
			CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
AFS std gill net	Common Carp	78	6.5	1.1	97		31	8	117	1
	Northern Pike	31	2.6	0.8	81	11	19	11	83	3
	Walleye	70	5.8	0.9	63	8	37	8	83	1
	Yellow Perch	110	9.2	1.4	91	4	75	6	106	1

## **10-Year Catch Per Unit Effort by Gear and Species**

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

Gear	Species	CPUE										Avg
		2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
AFS std gill net	Common Carp							0.0			6.5	3.25
	Northern Pike							0.7			2.6	1.65
	Walleye							16.0			5.8	10.90
	Yellow Perch							24.7			9.2	16.95
std exp gill net	Northern Pike				1.8							1.80
	Walleye				22.0							22.00
	Yellow Perch				16.5							16.50

## 10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

Gear	Species	Index	Year											
			2012	2013	2014	2015	2016	2017	2018	2019	2020	2021		
AFS std gill net	Walleye	PSD								74			63	
		PSD-P								19			37	
		Wr								91			83	
	Yellow Perch	PSD									21			91
		PSD-P									10			75
		Wr									115			106
std exp gill net	Walleye	PSD				52								
		PSD-P				5								
		Wr				85								
	Yellow Perch	PSD				90								
		PSD-P				35								
		Wr				108								



## Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Walleye

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2021	70	267 (8)	371 (24)	475 (3)	481 (5)	495 (2)	519 (4)	526 (3)	571 (4)	610 (2)	600 (15)
2018	200	264 (49)	377 (18)	417 (21)	467 (49)	481 (15)	539 (2)	500 (28)		551 (5)	588 (14)
2015	134	258 (22)	339 (31)	384 (18)	415 (49)		474 (4)	473 (1)	523 (8)	545 (1)	

Species: Yellow Perch

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2021	110	155 (10)	239 (6)	265 (83)	306 (10)				350 (1)		
2018	296	157 (233)	242 (50)	297 (7)		313 (3)		337 (3)			
2015	99	149 (3)	216 (60)	257 (1)	288 (32)	294 (2)	314 (1)				

## **Fish Condition**



Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

Species	Year	Length Groups							
		S-Q		Q-P		P-M		M	
		N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Walleye Gill Net	2018	49	89 (0.6)	107	93 (0.7)	32	89 (1.2)	4	82 (3.4)
	2021	26	86 (1.0)	18	86 (1.0)	18	80 (1.7)	8	72 (3.3)
Yellow Perch Gill Net	2018	233	115 (0.7)	33	117 (1.7)	21	118 (1.4)	9	111 (3.0)
	2021	10	108 (2.5)	18	106 (2.0)	72	106 (1.0)	10	103 (3.5)



## Length Frequency Distribution

Length frequency histogram of species sampled by year.

Species: Walleye  
Gear: AFS std gill net




Species: Yellow Perch  
Gear: AFS std gill net




## Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.


Species: Walleye  
Gear: AFS std gill net




Species: Walleye  
Gear: std exp gill net



Species: Yellow Perch  
Gear: AFS std gill net



Species: Yellow Perch  
Gear: std exp gill net



## **Fish Stocking**

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2011	Walleye	Fry	3,500,000
2013	Walleye	Fry	3,500,000
2015	Walleye	Fry	3,500,000
2018	Walleye	Fry	3,500,000
2021	Walleye	Fry	3,500,000