

## South Red Iron Survey Summary

South Red Iron, located 8.0 miles southeast of Lake City, is primarily managed as a northern pike, walleye, and yellow perch fishery; other fish species are present and contribute to the fishery.

- **Northern pike.** Northern pike numbers were considerably higher in 2021 than in 2018. At 8.3 per gill net, relative abundance was high. Sampled northern pike ranged in length from 16.1 to 32.3 inches, 66% were  $\geq 21.0$  inches and 8% were  $\geq 28.0$  inches.
- **Walleye.** Walleyes were not abundant in 2021 (2.4 per gill net). Those sampled ranged in length from 10.6 to 26.0 inches, 62% were  $\leq 15.0$  inches and 45% were  $\geq 20.0$  inches. Eleven year classes (2003, 2005, 2009, 2012 – 2019) were present, most (10 of 11) were represented by a low number of individuals (i.e.,  $\leq 4$ ). The oldest walleye sampled was from the 2003 (age-18) cohort. Individuals from the 2019 (age-2) cohort, which coincided with a fry stocking, were the most abundant accounting for 46% of walleye in the sample. Growth of the 2019 year class has been slower to age 2 than previous cohorts. In 2021, the mean length at capture at age 2 was 12.5 inches compared to 12.8 inches in 2015 and 13.5 inches in 2012.
- **Yellow perch.** Yellow perch numbers were higher in 2021 than in 2018. At 12.3 per gill net, relative abundance was moderate to high for South Red Iron Lake. Sampled yellow perch ranged in length from 5.1 to 10.6 inches, 3% were  $\geq 8.0$  inches and only 1% were  $\geq 10.0$  inches. Individuals from four consecutive cohorts (2016 – 2019) contributed to the catch, those from the 2018 (age-3) year class, which had a mean length at capture of 6.1 inches, were the most abundant accounting for 85% of yellow perch in the sample.

For more detailed results see the computer-generated South Dakota Statewide Fisheries Survey for Red Iron South (Marshall; below).

**SOUTH DAKOTA STATEWIDE FISHERIES SURVEY**  
**Red Iron South, Marshall County**  
**UJA-Lake-917-002**  
**2021**

**Lake Information**

<b>Name:</b>	Red Iron South	<b>Maximum Depth:</b>	15 Feet
<b>County:</b>	Marshall	<b>Mean Depth:</b>	8 Feet
<b>Surface Area:</b>	661 Acres		

**Surveys and Investigations**

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
AFS std gill net	Jun 15, 2021	4 net-nights
AFS std gill net	Jun 16, 2021	4 net-nights
AFS std gill net	Jun 17, 2021	4 net-nights

## **Common Fish Species Present**

Walleye

Smallmouth Bass

Northern Pike

Largemouth Bass

Yellow Perch

Black Bullhead

White Sucker

Black Crappie

Bluegill

Common Carp

---

## Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- **AFS std gill net** – 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- **std experimental gill net for non-Missouri River waters** - 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- **std experimental gill net for Missouri River reservoirs** – 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$CPUE = \frac{\text{number of fish}}{\text{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left( \frac{\text{number of fish} \geq \text{quality length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

$$PSD - P = \left( \frac{\text{number of fish} \geq \text{preferred length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (*W*) compared to a length-specific standard weight (*Ws*) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left( \frac{W}{W_s} \right) \times 100$$

Confidence intervals (**CI**) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (**S**), quality (**Q**), preferred (**P**), memorable (**M**) and trophy (**T**). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

## Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

Gear	Species	Sample Size (n)	Abundance		Stock Density Indices			Condition		
			CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
AFS std gill net	Black Bullhead	223	15.8	4.4	5	2	4	2	92	1
	Black Crappie	21	1.8	0.5	71	16	0		108	4
	Bluegill	5	0.4	0.3	80		0		106	4
	Common Carp	1	0.1	0.1	100		100		98	
	Northern Pike	100	8.3	1.8	66	7	8	4	90	1
	Walleye	29	2.4	0.9	62	14	45	14	93	2
	White Sucker	47	3.9	1.0	100		70	10	101	1
	Yellow Perch	147	12.3	3.5	3		1		92	1

## 10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

Gear	Species	CPUE										Avg
		2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
AFS std gill net	Black Bullhead							3.1			15.8	9.45
	Black Crappie							0.5			1.8	1.15
	Bluegill							0.5			0.4	0.45
	Common Carp							0.3			0.1	0.20
	Largemouth Bass							0.1			0.0	0.05
	Northern Pike							1.8			8.3	5.05
	Smallmouth Bass							0.9			0.0	0.45
	Walleye							2.2			2.4	2.30
	White Sucker							0.8			3.9	2.35
	Yellow Perch							5.5		12.3	8.90	
frame net (std 3/4 in)	Black Bullhead	15.9			49.9							32.90
	Black Crappie	1.3			0.5							0.90
	Bluegill	3.6			0.5							2.05
	Common Carp	0.1			0.0							0.05
	Northern Pike	1.3			0.4							0.85
	Smallmouth Bass	0.7			0.4							0.55
	Walleye	0.1			0.0							0.05
	White Sucker	0.1			0.5							0.30
	Yellow Perch	1.6			0.1							0.85
std exp gill net	Black Bullhead	26.5			96.3							61.40
	Black Crappie	1.2			1.5							1.35
	Northern Pike	12.0			3.0							7.50
	Smallmouth Bass	1.5			1.8							1.65
	Walleye	5.5			4.3							4.90
	White Sucker	6.8			2.0							4.40
		Yellow Perch	24.5			3.5						

## 10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

Gear	Species	Index	Year											
			2012	2013	2014	2015	2016	2017	2018	2019	2020	2021		
AFS std gill net	Northern Pike	PSD								70			66	
		PSD-P								0			8	
		Wr								86			90	
	Walleye	PSD									100			62
		PSD-P									58			45
		Wr									91			93
	Yellow Perch	PSD									2			3
		PSD-P									0			1
		Wr									99			92
std exp gill net	Northern Pike	PSD	49				67							
		PSD-P	3				17							
		Wr	88				79							
	Walleye	PSD	33				65							
		PSD-P	6				8							
		Wr	94				91							
	Yellow Perch	PSD	3				33							
		PSD-P	1				0							
		Wr	103				101							



## Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Walleye

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2021	28		317 (13)	460 (1)	486 (1)	524 (1)	510 (2)	524 (1)	592 (4)	575 (2)	611 (3)
2018	25	206 (1)		431 (3)	463 (2)	517 (6)	525 (5)	567 (3)	558 (3)	695 (2)	
2015	28	138 (2)	325 (8)	407 (11)	460 (1)	499 (5)	502 (1)				
2012	34	245 (2)	342 (22)	460 (7)	521 (3)						

Species: Yellow Perch

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2021	147		151 (16)	156 (125)	217 (5)	270 (1)					
2018	61		160 (59)	200 (2)							
2015	23	98 (1)	140 (14)	171 (1)	225 (2)	227 (5)					
2012	157	92 (3)	144 (107)	176 (41)	201 (6)						

## Fish Condition

Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).



Species	Year	Length Groups							
		S-Q		Q-P		P-M		M	
		N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Northern Pike Gill Net	2018	6	89 (1.3)	14	85 (1.3)	0		0	
	2021	34	92 (1.0)	58	89 (0.6)	8	93 (3.0)	0	
Walleye Gill Net	2018	0		10	90 (1.2)	12	93 (1.2)	2	84 (9.3)
	2021	11	85 (2.1)	5	104 (4.6)	12	95 (1.4)	1	97
Yellow Perch Gill Net	2018	60	99 (0.7)	1	83	0		0	
	2021	143	93 (0.6)	3	90 (0.3)	1	87	0	

## Length Frequency Distribution

Length frequency histogram of species sampled by year.



Species: Northern Pike

Gear: AFS std gill net





Species: Walleye

Gear: AFS std gill net




Species: Yellow Perch  
Gear: AFS std gill net




## Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.


Species: Northern Pike  
Gear: AFS std gill net




Species: Northern Pike  
Gear: std exp gill net




Species: Walleye  
Gear: AFS std gill net




Species: Walleye  
Gear: std exp gill net



Species: Yellow Perch  
Gear: AFS std gill net



Species: Yellow Perch  
Gear: std exp gill net



## **Fish Stocking**

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2010	Walleye	Large Fingerling	16,687
2012	Walleye	Large Fingerling	7,380
2014	Walleye	Large Fingerling	11,224
2017	Walleye	Fry	300,000
2019	Walleye	Fry	300,000
2021	Walleye	Fry	300,000