

SOUTH DAKOTA STATEWIDE FISHERIES SURVEY
Oakwood East, Brookings County
MBS-Lake-215-001
2021

Lake Information

Name:	Oakwood East	Maximum Depth:	9 Feet
County:	Brookings	Mean Depth:	6 Feet
Legal Description:	T111N-R51W-Sec. 4-5, 8-9, 16-27	OHWM Elevation:	1,627
Surface Area:	955 Acres	Outlet Elevation:	1,626

Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
AFS std gill net	Aug 17, 2021	6 net-nights

Common Fish Species Present

Yellow Perch

Walleye

Black Bullhead

Northern Pike

Bigmouth Buffalo

White Sucker

Common Carp

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- **AFS std gill net** – 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- **std experimental gill net for non-Missouri River waters** - 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- **std experimental gill net for Missouri River reservoirs** – 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$CPUE = \frac{\text{number of fish}}{\text{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{\text{number of fish} \geq \text{quality length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

$$PSD - P = \left(\frac{\text{number of fish} \geq \text{preferred length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (**W**) compared to a length-specific standard weight (**Ws**) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{W_s} \right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (S), quality (Q), preferred (P), memorable (M) and trophy (T). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

* **Methods/Species that ignore stock length**

Gear	Species	Sample Size (n)	Abundance		Stock Density Indices			Condition		
			CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
AFS std gill net	Bigmouth Buffalo	10	1.7	1.2	40		30			
	Black Bullhead	73	12.0	4.9	1		0			
	Common Carp	35	0.2	0.2	0		0			
	Northern Pike	14	2.3	1.0	64	22	0		92	2
	Walleye	61	3.7	2.0	91		5		98	1
	White Sucker	8	1.3	1.1	63		50			
	Yellow Perch	97	16.2	3.2	31	7	6	4	100	1

10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

* Methods/Species that ignore stock length

Gear	Species	CPUE										Avg
		2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
AFS std gill net	Bigmouth Buffalo						0.8	0.0	0.0	0.0	1.7	0.50
	Black Bullhead						1.7	1.0	0.5	0.7	12.0	3.18
	Common Carp						0.0	0.2	0.0	3.5	0.2	0.78
	Northern Pike						0.3	0.2	0.0	1.5	2.3	0.86
	Walleye						25.3	49.7	1.2	4.0	3.7	16.78
	White Sucker						3.8	1.8	1.0	2.2	1.3	2.02
	Yellow Perch						33.8	46.5	12.5	11.3	16.2	24.06
frame net (std 3/4 in)	Bigmouth Buffalo	2.2										2.20
	Black Bullhead	48.9										48.90
	Black Crappie	0.0										0.00
	Common Carp	3.2										3.20
	Green Sunfish	7.1										7.10
	Northern Pike	3.0										3.00
	Orangespotted Sunfish	0.0										0.00
	Tadpole Madtom	0.0										0.00
	Walleye	1.6										1.60
	White Sucker	7.4										7.40
	Yellow Bullhead	6.2										6.20
Yellow Perch	7.6										7.60	
std exp gill net	Bigmouth Buffalo	3.7		0.7	0.3	0.3						1.25
	Black Bullhead	35.3		0.3	0.0	6.7						10.58
	Common Carp	3.0		1.3	0.7	0.0						1.25
	Green Sunfish	0.7		0.0	0.0	0.0						0.18
	Northern Pike	2.7		1.3	4.0	2.0						2.50
	Orangespotted Sunfish	0.0		0.0	0.0	0.0						0.00
	Walleye	5.7		25.7	47.3	51.7						32.60
	White Sucker	12.7		7.7	7.3	2.3						7.50
	Yellow Bullhead	1.7		0.0	0.0	0.0						0.43
Yellow Perch	61.3		32.7	5.0	89.7						47.18	

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

Gear	Species	Index	Year										
			2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
AFS std gill net	Bigmouth Buffalo	PSD							0	0	0		40
		PSD-P							0	0	0		30
	Black Bullhead	PSD							50	67	33	0	1
		PSD-P							0	17	0	0	0
	Common Carp	PSD								0	0	0	0
		PSD-P								0	0	0	0
	Northern Pike	PSD							100	100	0	56	64
		PSD-P							0	0	0	22	0
		Wr							86	83		100	92
	Walleye	PSD							89	25	100	42	91
		PSD-P							11	11	14	21	5
		Wr							91	92	97	97	98
	White Sucker	PSD							100	82	100	77	63
		PSD-P							100	73	100	38	50
		Wr										91	
	Yellow Perch	PSD							37	18	24	21	31
		PSD-P							21	3	3	3	6
		Wr							96	100	103	98	100
frame net (std 3/4 in)	Bigmouth Buffalo	PSD	75										
		PSD-P	30										
		Wr	88										
	Black Bullhead	PSD	11										
		PSD-P	0										
		Wr	95										
	Common Carp	PSD	41										
		PSD-P	21										
		Wr	110										
	Northern Pike	PSD	67										
		PSD-P	26										
		Wr	91										
Walleye	PSD	86											
	PSD-P	7											
	Wr	100											

Gear	Species	Index	Year									
			2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
frame net (std 3/4 in)	White Sucker	PSD	90									
		PSD-P	36									
		Wr	92									
	Yellow Perch	PSD	53									
		PSD-P	9									
		Wr	99									
std exp gill net	Bigmouth Buffalo	PSD	9		100	100	0					
		PSD-P	0		50	0	0					
		Wr	96									
	Black Bullhead	PSD	24		0		30					
		PSD-P	0		0		0					
		Wr	103									
	Common Carp	PSD	0		100	100						
		PSD-P	0		100	100						
		Wr	112									
	Northern Pike	PSD	38		100	92	67					
		PSD-P	0		75	42	0					
		Wr	95		88	87	81					
	Walleye	PSD	76		25	25	70					
		PSD-P	0		1	0	0					
		Wr	95		93	87	78					
	White Sucker	PSD	87		70	73	100					
		PSD-P	8		26	55	57					
		Wr	89									
	Yellow Perch	PSD	58		0	73	32					
		PSD-P	5		0	20	1					
		Wr	102		108	107	87					

Fish Condition



Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

Species	Year	Length Groups							
		S-Q		Q-P		P-M		M	
		N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Northern Pike Gill Net	2017	0		2	86 (0.8)	0		0	
	2018	0		1	83	0		0	
	2019	0		0		0		0	
	2020	4	100 (2.1)	3	99 (1.8)	2	100 (6.6)	0	
	2021	5	94 (4.2)	9	91 (1.7)	0		0	
Walleye Gill Net	2017	17	91 (0.9)	118	92 (0.6)	17	89 (1.0)	0	
	2018	223	94 (4.1)	43	88 (0.9)	32	89 (1.0)	0	
	2019	0		6	98 (2.8)	1	92	0	
	2020	14	96 (1.5)	5	94 (5.8)	5	103 (2.8)	0	
	2021	2	98 (3.4)	19	98 (1.2)	0		1	91
White Sucker Gill Net	2020	3		5	91	2		3	
Yellow Perch Gill Net	2017	128	95 (0.7)	33	97 (1.0)	42	99 (0.9)	0	
	2018	228	101 (0.8)	43	99 (1.5)	8	82	0	
	2019	57	105 (1.1)	16	100 (1.8)	2	97 (1.9)	0	
	2020	54	99 (0.9)	12	96 (1.3)	2	96 (1.8)	0	
	2021	67	101 (1.0)	24	98 (0.9)	5	97 (1.4)	1	


Length Frequency Distribution

Length frequency histogram of species sampled by year.


Species: Black Bullhead
Gear: AFS std gill net





Species: Black Bullhead
Gear: std exp gill net





Species: Common Carp
Gear: AFS std gill net










Species: Northern Pike
Gear: AFS std gill net




Species: Walleye
Gear: AFS std gill net










Species: Walleye
 Gear: std exp gill net






Species: White Sucker
 Gear: AFS std gill net




Species: Yellow Perch
 Gear: AFS std gill net






Species: Yellow Perch
 Gear: std exp gill net




Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.


Species: Black Bullhead
Gear: AFS std gill net




Species: Black Bullhead
Gear: std exp gill net




Species: Common Carp
Gear: AFS std gill net




Species: Common Carp
Gear: std exp gill net




Species: Northern Pike
Gear: AFS std gill net




Species: Northern Pike
Gear: std exp gill net




Species: Walleye
Gear: AFS std gill net




Species: Walleye
Gear: std exp gill net




Species: White Sucker
Gear: AFS std gill net




Species: White Sucker
Gear: std exp gill net



Species: Yellow Perch
Gear: AFS std gill net



Species: Yellow Perch
Gear: std exp gill net



Fish Stocking

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2010	Walleye	Fry	1,000,000
2012	Walleye	Fry	500,000
2013	Walleye	Fry	650,000
2014	Walleye	Fry	453,750
2019	Walleye		933,000
2019	Yellow Perch	Juvenile	47,000
2021	Walleye	Fry	1,933,000