

**SOUTH DAKOTA STATEWIDE FISHERIES SURVEY**  
**Hwy 81 Southeast, Brookings County**  
**MBS-Lake-233-003**  
**2021**

**Lake Information**

**Name:** Hwy 81 Southeast  
**County:** Brookings  
**Surface Area:** 141 Acres

**Surveys and Investigations**

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
AFS std gill net	Jul 21, 2021	3 net-nights

## **Common Fish Species Present**

Walleye

Yellow Perch

White Bass

Common Carp

Black Bullhead

Black Crappie

Northern Pike

---

## Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- **AFS std gill net** – 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- **std experimental gill net for non-Missouri River waters** - 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- **std experimental gill net for Missouri River reservoirs** – 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$CPUE = \frac{\text{number of fish}}{\text{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left( \frac{\text{number of fish} \geq \text{quality length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

$$PSD - P = \left( \frac{\text{number of fish} \geq \text{preferred length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (**W**) compared to a length-specific standard weight (**Ws**) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left( \frac{W}{W_s} \right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (S), quality (Q), preferred (P), memorable (M) and trophy (T). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

## Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

\* **Methods/Species that ignore stock length**

Gear	Species	Sample Size (n)	Abundance		Stock Density Indices			Condition	
			CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr
AFS std gill net	Black Bullhead	7	2.3	1.3	71		43		
	Black Crappie	4	1.3	1.3	100		75	97	7
	Common Carp	9	3.0	2.9	100		33		
	Northern Pike	2	0.7	1.3	50		0	69	1
	Walleye	3	1.0	1.1	100		67	73	8
	White Bass	12	4.0	4.7	92		83	82	2
	Yellow Perch	9	3.0	0.0	78		0	93	1

## 10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

\* Methods/Species that ignore stock length

Gear	Species	CPUE										Avg
		2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
AFS std gill net	Black Bullhead										2.3	2.30
	Black Crappie										1.3	1.30
	Common Carp										3.0	3.00
	Northern Pike										0.7	0.70
	Walleye										1.0	1.00
	White Bass										4.0	4.00
	Yellow Perch										3.0	3.00
frame net (std 3/4 in)	Black Bullhead	361.6										361.6 0
	Black Crappie	0.2										0.20
	Bluegill	0.1										0.10
	Common Carp	6.3										6.30
	Green Sunfish	0.3										0.30
	Northern Pike	0.0										0.00
	Orangespotted Sunfish	0.0										0.00
	Smallmouth Bass	0.1										0.10
	Walleye	0.7										0.70
	White Bass	1.0										1.00
	White Sucker	0.0										0.00
	Yellow Bullhead	12.5										12.50
Yellow Perch	16.0										16.00	
std exp gill net	Black Bullhead	47.5		34.7	18.3	1.3						25.45
	Black Crappie	0.5		0.0	0.0	0.0						0.13
	Common Carp	0.5		1.0	0.3	1.7						0.88
	Northern Pike	1.0		0.0	0.0	0.0						0.25
	Smallmouth Bass	0.0		0.0	0.0	0.3						0.08
	Walleye	10.0		5.0	28.0	8.3						12.83
	White Bass	1.0		0.3	10.3	7.3						4.73
	White Sucker	0.0		0.3	0.7	0.0						0.25
	Yellow Bullhead	0.5		1.0	1.7	1.3						1.13
Yellow Perch	162.0		38.0	32.3	35.0						66.83	

## 10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

Gear	Species	Index	Year													
			2012	2013	2014	2015	2016	2017	2018	2019	2020	2021				
AFS std gill net	Black Bullhead	PSD													71	
		PSD-P														43
		Wr														97
	Black Crappie	PSD														100
		PSD-P														75
		Wr														97
	Common Carp	PSD														100
		PSD-P														33
		Wr														97
	Northern Pike	PSD														50
		PSD-P														0
		Wr														69
	Walleye	PSD														100
		PSD-P														67
		Wr														73
	White Bass	PSD														92
		PSD-P														83
		Wr														82
Yellow Perch	PSD														78	
	PSD-P														0	
	Wr														93	
frame net (std 3/4 in)	Black Bullhead	PSD	27													
		PSD-P	2													
		Wr	86													
	Black Crappie	PSD	0													
		PSD-P	0													
		Wr	105													
	Common Carp	PSD	67													
		PSD-P	0													
		Wr	110													
	Walleye	PSD	57													
		PSD-P	0													
		Wr	83													
White Bass	PSD	100														
	PSD-P	50														

Gear	Species	Index	Year									
			2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
frame net (std 3/4 in)	White Bass	Wr	87									
	Yellow Perch	PSD	14									
		PSD-P	1									
		Wr	85									
std exp gill net	Black Bullhead	PSD	31		55	60	100					
		PSD-P	2		6	7	75					
		Wr	98									
	Black Crappie	PSD	0									
		PSD-P	0									
		Wr	116									
	Common Carp	PSD	0		67	100	100					
		PSD-P	0		67	0	20					
		Wr	123									
	Northern Pike	PSD	50									
		PSD-P	0									
		Wr	90									
	Walleye	PSD	85		80	21	52					
		PSD-P	15		40	13	12					
		Wr	93		90	91	91					
	White Bass	PSD	100		100	100	91					
		PSD-P	50		100	74	50					
		Wr	90		88	99	93					
	Yellow Perch	PSD	40		23	88	19					
		PSD-P	7		6	16	15					
		Wr	92		97	94	112					



## Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Walleye

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2012	26	242 (9)	395 (1)	474 (13)	552 (1)	555 (2)					

Species: Yellow Perch

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2015	96	179 (11)	231 (72)	263 (9)	282 (4)						
2014	114	166 (89)	233 (12)	245 (12)	265 (1)						
2012	324	166 (200)	231 (121)	284 (3)							

## **Fish Condition**


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

Species	Year	Length Groups							
		S-Q		Q-P		P-M		M	
		N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Northern Pike Gill Net	2021	1	70	1	69	0		0	
Walleye Gill Net	2021	0		1	77	1	80	1	61
White Bass Gill Net	2021	1	82	1	81	2	85 (3.2)	8	81 (1.8)
Yellow Perch Gill Net	2021	2	92 (2.9)	7	93 (1.0)	0		0	


## Length Frequency Distribution

Length frequency histogram of species sampled by year.


Species: Walleye  
Gear: std exp gill net




Species: White Bass  
Gear: AFS std gill net



Species: White Bass  
Gear: std exp gill net



Species: Yellow Perch  
Gear: std exp gill net




## Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.


Species: Black Bullhead

Gear: AFS std gill net




Species: Black Bullhead


Gear: std exp gill net




Species: Black Crappie  
Gear: frame net (std 3/4 in)




Species: Common Carp  
Gear: AFS std gill net




Species: Common Carp  
Gear: std exp gill net




Species: Northern Pike  
Gear: AFS std gill net




Species: Northern Pike  
Gear: std exp gill net




Species: Walleye  
Gear: AFS std gill net



Species: Walleye  
Gear: std exp gill net




Species: White Bass  
Gear: AFS std gill net







Species: White Bass  
Gear: std exp gill net



Species: Yellow Perch  
Gear: AFS std gill net



Species: Yellow Perch  
Gear: std exp gill net



## **Fish Stocking**

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2021	Walleye	Juvenile	15,840