

SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Byre, Lyman County

MED-Lake-25-000

2021

Lake Information

Name: Byre **Maximum Depth:** 17 Feet
County: Lyman **Mean Depth:** 7 Feet
Legal Description: T75-R105-S4
Surface Area: 121 Acres

Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
fall night EF-WAE	Sep 15, 2021	3600 seconds

Common Fish Species Present

Walleye

Largemouth Bass

Black Crappie

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- **AFS std gill net** – 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- **std experimental gill net for non-Missouri River waters** - 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- **std experimental gill net for Missouri River reservoirs** – 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$CPUE = \frac{\text{number of fish}}{\text{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{\text{number of fish} \geq \text{quality length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

$$PSD - P = \left(\frac{\text{number of fish} \geq \text{preferred length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (**W**) compared to a length-specific standard weight (**Ws**) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{W_s} \right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (S), quality (Q), preferred (P), memorable (M) and trophy (T). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

* Methods/Species that ignore stock length

Gear	Species	CPUE										Avg
		2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	
AFS std gill net	Black Bullhead								1.0			1.00
	Black Crappie								2.0			2.00
	Northern Pike								2.0			2.00
	Walleye								2.0			2.00
	Yellow Perch								0.5			0.50
boat shocker (night)	Largemouth Bass		11.0			7.0			20.0			12.67
fall night EF-WAE*	Walleye								23.0		8.0	15.50
frame net (std 3/4 in)	Black Bullhead		1.4			45.8			3.2			16.80
	Black Crappie		9.8			26.0			91.9			42.57
	Bluegill		0.0			0.9			15.3			5.40
	Common Carp		1.8			0.9			0.5			1.07
	Green Sunfish		0.0			0.2			0.0			0.07
	Largemouth Bass		0.4			0.0			0.0			0.13
	Northern Pike		0.3			1.1			0.3			0.57
	Orangespotted Sunfish		0.0			0.0			0.0			0.00
	Walleye		1.9			1.3			2.5			1.90
	Yellow Perch		0.0			0.2			0.8			0.33
std exp gill net	Black Bullhead		0.5			35.5						18.00
	Black Crappie		0.0			7.0						3.50
	Common Carp		1.0			2.0						1.50
	Northern Pike		0.0			0.5						0.25
	Walleye		1.0			8.0						4.50

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

Gear	Species	Index	Year									
			2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
AFS std gill net	Black Crappie	PSD										100
		PSD-P										25
		Wr										81
	Walleye	PSD										75
		PSD-P										0
		Wr										82
boat shocker (night)	Largemouth Bass	PSD		91			100					15
		PSD-P		91			43					10
		Wr		121			116					119
	Walleye	PSD		32			24					
		PSD-P		5			0					
		Wr		91			89					
frame net (std 3/4 in)	Black Crappie	PSD		89			69					71
		PSD-P		1			0					1
		Wr		88			101					92
	Largemouth Bass	PSD		100								0
		PSD-P		50								0
		Wr		109								
	Walleye	PSD		89			62					44
		PSD-P		47			54					24
		Wr		83			75					101
std exp gill net	Black Crappie	PSD					86					
		PSD-P					0					
		Wr					93					
	Walleye	PSD		50			13					
		PSD-P		0			6					
		Wr		77			78					

Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Black Crappie

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2019	1242	120 (97)	125 (276)	191 (194)	208 (650)	235 (13)	250 (13)				
2016	259			203 (235)	218 (25)						
2013	99	100 (1)	163 (3)	193 (7)	211 (17)	220 (45)	224 (26)				

Species: Largemouth Bass

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2019	20	217 (9)	235 (8)	256 (1)		423 (1)		461 (1)			
2016	7	183 (1)		342 (3)	335 (1)		441 (1)		460 (1)		
2013	63	165 (54)	272 (1)			416 (7)			521 (1)		

Species: Walleye

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2016	16		303 (8)	309 (6)	508 (2)						
2013	2		324 (1)	404 (1)							

Fish Condition



Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

Species	Year	Length Groups							
		S-Q		Q-P		P-M		M	
		N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Black Crappie Frame Net	2019	262	97 (1.2)	644	90 (0.5)	13	84	0	
Largemouth Bass Electro Fishing	2019	17	118 (2.2)	1	135	2	115 (5.4)	0	
Walleye Gill Net	2019	1	76	3	84 (2.0)	0		0	



Length Frequency Distribution

Length frequency histogram of species sampled by year.


Species: Black Crappie
Gear: frame net (std 3/4 in)



Species: Largemouth Bass
Gear: boat shocker (night)




Species: Walleye
Gear: std exp gill net




Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.


Species: Black Crappie
Gear: frame net (std 3/4 in)




Species: Largemouth Bass
Gear: boat shocker (night)



Species: Walleye
Gear: AFS std gill net



Species: Walleye
Gear: std exp gill net



Fish Stocking

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2010	Walleye	Small Fingerling	14,980
2013	Walleye	Large Fingerling	1,144
2015	Walleye	Large Fingerling	900
2017	Walleye	Large Fingerling	2,960
2019	Walleye	Small Fingerling	9,185
2021	Walleye	Juvenile	10,866