SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Trail City, Corson County ULO-Lake-765-000 2021

Lake Information

Name: Trail City Maximum Depth: 16 Feet

County: Corson Mean Depth: 5 Feet

Legal Description: T18-R27-S35

Surface Area: 81 Acres

Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
frame net (std 3/4 in)	Jul 07, 2021	5 net-nights
frame net (std 3/4 in)	Jul 08, 2021	5 net-nights

Common Fish Species Present

Northern Pike

Yellow Perch

White Crappie

Black Crappie

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- AFS std gill net 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- std experimental gill net for non-Missouri River waters 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- std experimental gill net for Missouri River reservoirs 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$\mathit{CPUE} = \frac{\mathit{number of fish}}{\mathit{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{number\ of\ fish \ge quality\ length}{number\ of\ fish \ge stock\ length}\right) \times 100$$

$$\textit{PSD} - \textit{P} = \left(\frac{number\ of\ fish\ \geq preferred\ length}{number\ of\ fish\ \geq stock\ length}\right) \ge 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{Ws}\right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (S), quality (Q), preferred (P), memorable (M) and trophy (T). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

	St	ock	Qu	ality	Preferred		Mem	orable	Tro	ophy
Species Name	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

* Methods/Species that ignore stock length

			Abun	dance	Stock Density Indices					ndition
Gear	Species	Sample Size (n)	CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
frame net (std 3/4	Black Crappie	2	0.2	0.2	100		100		108	1
in)	Northern Pike	11	1.1	0.4	91		0		83	3
	White Crappie	3	0.3	0.3	100		100		90	2

10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

* Methods/Species that ignore stock length

							CPUE					
Gear	Species	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	Avg
AFS std gill net	Black Crappie							0.3				0.30
	Northern Pike							2.0				2.00
	Yellow Perch							0.8				0.80
frame net (std	Black Crappie		0.0			0.0		1.3			0.2	0.38
3/4 in)	Bluegill		0.0			0.0		0.1			0.0	0.03
	Northern Pike		5.4			1.6		1.1			1.1	2.30
	White Crappie		0.0			0.0		0.0			0.3	0.08
	Yellow Perch		0.0			0.0		0.4			0.0	0.10
std exp gill net	Northern Pike		15.0			5.0						10.00
	Yellow Perch		1.0			0.0						0.50

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

							Ye	ar				
Gear	Species	Inde	ex 2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
AFS std gill net	Black Crappie	PSI)						100			
		PSI)-P						100			
		Wr							98			
	Northern Pike	PSI)						63			
		PSI)-P						25			
		Wr							86			
	Yellow Perch	PSI)						100			
		PSI)-P						33			
		Wr							120			
frame net (std	Black Crappie	PSI)						100			100
3/4 in)		PSI)-P						100			100
		Wr							88			108
	Northern Pike	PSI)	72			75		64			91
		PSI)-P	0			0		0			0
		Wr		71			83		70			83
	White Crappie	PSI)									100
		PSI)-P									100
		Wr										90
	Yellow Perch	PSI)						100			
		PSI)-P						50			
		Wr							112			
std exp gill net	Northern Pike	PSI)	73			100					
		PSI)-P	0			0					
		Wr		76			86					
	Yellow Perch	PSI)	100								
		PSI)-P	50								
		Wr		108								

Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Yellow Perch

				Mean Ler	ngth (expa	nded san	nple numbe	er) at capt	ure by age	Э	
Year	N	1	2	3	4	5	6	7	8	9	10+
2018	3			205 (1)	205 (1)		258 (1)				
2013	2					259 (2)					

Fish Condition


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

		Length Groups							
			S-Q		Q-P		P-M		M
Species	Year	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Black Crappie Frame Net	2018	0		0		0		13	88 (1.9)
	2021	0		0		1	108	1	109
Northern Pike Gill Net	2018	3	99 (3.6)	3	72 (2.1)	1	82	1	93
White Crappie Frame Net	2021	0		0		0		3	90 (1.9)
Yellow Perch Gill Net	2018	0		2	117 (7.2)	1	125	0	


Length Frequency Distribution

Length frequency histogram of species sampled by year.

Species: Black Crappie Gear: frame net (std 3/4 in)




Species: Northern Pike Gear: std exp gill net




Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.


Species: Black Crappie Gear: frame net (std 3/4 in)




Species: Northern Pike Gear: AFS std gill net




Species: Northern Pike Gear: std exp gill net




Species: White Crappie Gear: frame net (std 3/4 in)



Species: Yellow Perch Gear: AFS std gill net



Species: Yellow Perch Gear: std exp gill net



Fish Stocking

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2012	Yellow Perch	Adult	400
2014	Yellow Perch	Adult	250
2015	Yellow Perch	Adult	578
2015	Yellow Perch	Fingerling	3,180
2017	Black Crappie	Adult	200
2018	Black Crappie	Adult	170
2019	Black Crappie	Adult	50
2020	Black Crappie	Adult	41