# SOUTH DAKOTA STATEWIDE FISHERIES SURVEY
## Scott, Minnehaha County
## LBS-Lake-65-000
## 2022

## Lake Information

**Name:** Scott

**County:** Minnehaha

**Legal Description:** T102-R51-Sec. 7-8

**Surface Area:** 115 Acres

**Maximum Depth:** 11 Feet

**Mean Depth:** 4 Feet

## Surveys and Investigations

Survey methods used by gear type, date, and effort.

| Gear | Date | Effort |
|---|---|---|
| AFS std gill net | Aug 18, 2022 | 3 net-nights |

## Common Fish Species Present

Yellow Perch

Walleye

Black Bullhead

Northern Pike

Common Carp

Black Crappie

## Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- **AFS std gill net** – 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- **std experimental gill net for non-Missouri River waters** - 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- s**td experimental gill net for Missouri River reservoirs** – 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$CPUE = \frac{number\ of\ fish}{effort}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{number\ of\ fish \geq quality\ length}{number\ of\ fish \geq stock\ length}\right) \text{x } 100$$

$$PSD - P = \left(\frac{number\ of\ fish \geq preferred\ length}{number\ of\ fish \geq stock\ length}\right) \text{x } 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{Ws}\right) \text{x } 100$$

Confidence intervals (**CI**) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (**S**), quality (**Q**), preferred (**P**), memorable (**M**) and trophy (**T**). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

| | Stock | | Quality | | Preferred | | Memorable | | Trophy | |
|---|---|---|---|---|---|---|---|---|---|---|
| Species Name | (in) | (cm) | (in) | (cm) | (in) | (cm) | (in) | (cm) | (in) | (cm) |
| Black Bullhead | 6 | 15 | 9 | 23 | 12 | 30 | 15 | 38 | 18 | 46 |
| Black Crappie | 5 | 13 | 8 | 20 | 10 | 25 | 12 | 30 | 15 | 38 |
| Bluegill | 3 | 8 | 6 | 15 | 8 | 20 | 10 | 25 | 12 | 30 |
| Brown Trout | 8 | 20 | 12 | 30 | 16 | 40 | 20 | 50 | 18 | 46 |
| Channel Catfish | 11 | 28 | 16 | 41 | 24 | 61 | 28 | 71 | 36 | 91 |
| Freshwater Drum | 8 | 20 | 12 | 30 | 15 | 38 | 20 | 51 | 25 | 63 |
| Lake Trout | 12 | 30 | 20 | 50 | 26 | 65 | 31 | 80 | 39 | 100 |
| Largemouth Bass | 8 | 20 | 12 | 30 | 15 | 38 | 20 | 51 | 25 | 63 |
| Muskellunge | 20 | 51 | 30 | 76 | 38 | 97 | 42 | 107 | 50 | 127 |
| Northern Pike | 14 | 35 | 21 | 53 | 28 | 71 | 34 | 86 | 44 | 112 |
| Pumpkinseed | 3 | 8 | 6 | 15 | 8 | 20 | 10 | 25 | 12 | 30 |
| Rainbow Trout | 10 | 25 | 16 | 40 | 20 | 50 | 26 | 65 | 31 | 80 |
| Rudd | 6 | 15 | 10 | 25 | 12 | 30 | 15 | 38 | 19 | 48 |
| Sauger | 8 | 20 | 12 | 30 | 15 | 38 | 20 | 51 | 25 | 63 |
| Smallmouth Bass | 7 | 18 | 11 | 28 | 14 | 35 | 17 | 43 | 20 | 51 |
| Walleye | 10 | 25 | 15 | 38 | 20 | 51 | 25 | 63 | 30 | 76 |
| White Bass | 6 | 15 | 9 | 23 | 12 | 30 | 15 | 38 | 18 | 46 |
| White Crappie | 5 | 13 | 8 | 20 | 10 | 25 | 12 | 30 | 15 | 38 |
| Yellow Bullhead | 4 | 10 | 7 | 18 | 9 | 23 | 11 | 28 | 14 | 36 |
| Yellow Perch | 5 | 13 | 8 | 20 | 10 | 25 | 12 | 30 | 15 | 38 |

## Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

*** Methods/Species that ignore stock length**

| | | | Abundance | | Stock Density Indices | | | | Condition | |
|---|---|---|---|---|---|---|---|---|---|---|
| Gear | Species | Sample Size (n) | CPUE | CI-80 | PSD | CI-80 | PSD-P | CI-80 | Wr | CI-80 |
| AFS std gill net | Black Bullhead | 302 | 91.7 | 54.5 | 1 | | 0 | | | |
| | Black Crappie | 1 | 0.3 | 0.6 | 100 | | 100 | | 121 | |
| | Common Carp | 1 | 0.3 | 0.6 | 100 | | 100 | | | |
| | Northern Pike | 3 | 1.0 | 1.1 | 100 | | 0 | | 82 | 5 |
| | Walleye | 17 | 5.0 | 3.3 | 100 | | 40 | 21 | 94 | 2 |
| | Yellow Perch | 11 | 3.7 | 3.8 | 18 | | 0 | | 99 | 4 |

## 10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

*** Methods/Species that ignore stock length**

| | | CPUE | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Gear | Species | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 | 2021 | 2022 | Avg |
| AFS std frame net | Black Bullhead | | | | | 271.4 | | | | | | 271.40 |
| | Black Crappie | | | | | 1.6 | | | | | | 1.60 |
| | Bluegill | | | | | 0.0 | | | | | | 0.00 |
| | Green Sunfish | | | | | 0.8 | | | | | | 0.80 |
| | Orangespotted Sunfish | | | | | 0.0 | | | | | | 0.00 |
| | Sunfish Hybrid | | | | | 5.2 | | | | | | 5.20 |
| | Walleye | | | | | 0.8 | | | | | | 0.80 |
| | Yellow Perch | | | | | 2.0 | | | | | | 2.00 |
| AFS std gill net | Black Bullhead | | | | | 102.3 | 92.8 | 63.7 | | | 91.7 | 87.63 |
| | Black Crappie | | | | | 0.5 | 0.0 | 0.3 | | | 0.3 | 0.28 |
| | Common Carp | | | | | 0.0 | 0.0 | 0.0 | | | 0.3 | 0.08 |
| | Northern Pike | | | | | 0.3 | 0.0 | 1.3 | | | 1.0 | 0.65 |
| | O. Spotted X Gr. Sunfish Hybrid | | | | | 0.0 | 0.0 | 0.0 | | | 0.0 | 0.00 |
| | Orangespotted Sunfish | | | | | 0.0 | 0.0 | 0.0 | | | 0.0 | 0.00 |
| | Pumpkinseed | | | | | 0.8 | 0.0 | 0.0 | | | 0.0 | 0.20 |
| | Walleye | | | | | 7.5 | 6.8 | 2.7 | | | 5.0 | 5.50 |
| | Yellow Perch | | | | | 5.0 | 6.5 | 0.7 | | | 3.7 | 3.98 |
| frame net (std 3/4 in) | Black Bullhead | 496.0 | | | 448.6 | | | | | | | 472.30 |
| | Green Sunfish | 0.0 | | | 1.4 | | | | | | | 0.70 |
| | Yellow Perch | 0.4 | | | 0.6 | | | | | | | 0.50 |
| std exp gill net | Black Bullhead | 273.0 | 217.0 | 88.7 | 327.0 | | | | | | | 226.43 |
| | Black Crappie | 0.0 | 0.3 | 0.7 | 0.0 | | | | | | | 0.25 |
| | Green Sunfish | 0.0 | 0.0 | 0.0 | 0.0 | | | | | | | 0.00 |
| | Northern Pike | 0.7 | 0.0 | 0.0 | 0.3 | | | | | | | 0.25 |
| | Walleye | 2.7 | 4.3 | 5.7 | 0.0 | | | | | | | 3.18 |
| | Yellow Perch | 7.7 | 9.0 | 26.0 | 31.7 | | | | | | | 18.60 |

## 10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

| | | | Year | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Gear | Species | Index | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 | 2021 | 2022 |
| AFS std frame net | Black Bullhead | PSD | | | | | 3 | | | | | |
| | | PSD-P | | | | | 0 | | | | | |
| | Black Crappie | PSD | | | | | 63 | | | | | |
| | | PSD-P | | | | | 0 | | | | | |
| | | Wr | | | | | 104 | | | | | |
| | Walleye | PSD | | | | | 75 | | | | | |
| | | PSD-P | | | | | 50 | | | | | |
| | | Wr | | | | | 80 | | | | | |
| | Yellow Perch | PSD | | | | | 50 | | | | | |
| | | PSD-P | | | | | 0 | | | | | |
| | | Wr | | | | | 91 | | | | | |
| AFS std gill net | Black Bullhead | PSD | | | | | 3 | 4 | 6 | | | 1 |
| | | PSD-P | | | | | 0 | 0 | 0 | | | 0 |
| | Black Crappie | PSD | | | | | 100 | | 0 | | | 100 |
| | | PSD-P | | | | | 0 | | 0 | | | 100 |
| | | Wr | | | | | 102 | | 101 | | | 121 |
| | Common Carp | PSD | | | | | | | 0 | | | 100 |
| | | PSD-P | | | | | | | 0 | | | 100 |
| | Northern Pike | PSD | | | | | 100 | | 25 | | | 100 |
| | | PSD-P | | | | | 0 | | 0 | | | 0 |
| | | Wr | | | | | 92 | | 89 | | | 82 |
| | Walleye | PSD | | | | | 90 | 67 | 100 | | | 100 |
| | | PSD-P | | | | | 60 | 19 | 25 | | | 40 |
| | | Wr | | | | | 95 | 98 | 93 | | | 94 |
| | Yellow Perch | PSD | | | | | 55 | 54 | 100 | | | 18 |
| | | PSD-P | | | | | 5 | 0 | 0 | | | 0 |
| | | Wr | | | | | 101 | 96 | 105 | | | 99 |
| frame net (std 3/4 in) | Black Bullhead | PSD | 33 | | | 1 | | | | | | |
| | | PSD-P | 0 | | | 0 | | | | | | |
| | | Wr | 90 | | | | | | | | | |
| | Yellow Perch | PSD | 50 | | | 0 | | | | | | |
| | | PSD-P | 0 | | | 0 | | | | | | |

| | | | Year | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Gear | Species | Index | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 | 2021 | 2022 |
| frame net (std 3/4 in) | Yellow Perch | Wr | 109 | | | 106 | | | | | | |
| std exp gill net | Black Bullhead | PSD | 40 | 24 | 12 | 1 | | | | | | |
| | | PSD-P | 0 | 0 | 0 | 0 | | | | | | |
| | | Wr | 94 | | | | | | | | | |
| | Black Crappie | PSD | | 0 | 0 | | | | | | | |
| | | PSD-P | | 0 | 0 | | | | | | | |
| | | Wr | | 127 | 115 | | | | | | | |
| | Northern Pike | PSD | 100 | | | 0 | | | | | | |
| | | PSD-P | 0 | | | 0 | | | | | | |
| | | Wr | 96 | | | 97 | | | | | | |
| | Walleye | PSD | 50 | 62 | 88 | | | | | | | |
| | | PSD-P | 13 | 8 | 0 | | | | | | | |
| | | Wr | 92 | 92 | 91 | | | | | | | |
| | Yellow Perch | PSD | 43 | 0 | 0 | 2 | | | | | | |
| | | PSD-P | 17 | 0 | 0 | 0 | | | | | | |
| | | Wr | 107 | 103 | 104 | 101 | | | | | | |

## Fish Condition

Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

| | | Length Groups | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| | | S-Q | | Q-P | | P-M | | M | |
| Species | Year | N | Wr (SE) | N | Wr (SE) | N | Wr (SE) | N | Wr (SE) |
| Northern Pike Gill Net | 2019 | 3 | 87 (1.0) | 1 | 93 | 0 | | 0 | |
| | 2022 | 0 | | 3 | 82 (4.1) | 0 | | 0 | |
| Walleye Gill Net | 2018 | 9 | 101 (1.9) | 13 | 96 (1.8) | 5 | 97 (3.2) | 0 | |
| | 2019 | 0 | | 6 | 95 (3.2) | 2 | 88 (0.0) | 0 | |
| | 2022 | 0 | | 9 | 94 (2.3) | 4 | 94 (1.2) | 2 | 89 (5.6) |
| Yellow Perch Gill Net | 2018 | 12 | 98 (2.1) | 14 | 95 (1.5) | 0 | | 0 | |
| | 2019 | 0 | | 2 | 105 (3.0) | 0 | | 0 | |
| | 2022 | 9 | 98 (3.2) | 2 | 104 (5.9) | 0 | | 0 | |

## Length Frequency Distribution

Length frequency histogram of species sampled by year.

Species: Black Bullhead
Gear: AFS std gill net




2017




2018




2019




2022

Species: Walleye
Gear: AFS std gill net




2017




2018




2022

Species: Yellow Perch
Gear: AFS std gill net




2017




2018


5
4
3
2
1
0
Number
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
Total Length (cm)

2022

## Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.

Species: Black Bullhead
Gear: AFS std gill net




Species: Black Bullhead
Gear: std exp gill net



Species: Black Crappie
Gear: AFS std frame net




Species: Common Carp
Gear: AFS std gill net



Species: Northern Pike
Gear: AFS std gill net




Species: Northern Pike
Gear: std exp gill net



Species: Walleye
Gear: AFS std gill net




Species: Walleye
Gear: std exp gill net



Species: Yellow Perch
Gear: AFS std gill net




Species: Yellow Perch
Gear: std exp gill net



## Fish Stocking

Number of fish stocked by year, species, and size.

| Year | Species | Size | Number |
|---|---|---|---|
| 2011 | Walleye | Small Fingerling | 12,480 |
| 2011 | Yellow Perch | Small Fingerling | 57,680 |
| 2012 | Yellow Perch | Fingerling | 96,640 |
| 2012 | Yellow Perch | Juvenile | 19,891 |
| 2012 | Yellow Perch | Large Fingerling | 2,470 |
| 2013 | Yellow Perch | Adult | 3,516 |
| 2014 | Walleye | Fry | 108,000 |
| 2014 | Yellow Perch | Adult | 3,570 |
| 2015 | Walleye | Juvenile | 212 |
| 2015 | Walleye | Small Fingerling | 7,560 |
| 2015 | Yellow Perch | Adult | 6,147 |
| 2015 | Yellow Perch | Fingerling | 11,060 |
| 2016 | Walleye | Juvenile | 1,390 |
| 2016 | Walleye | Small Fingerling | 7,560 |
| 2016 | Yellow Perch | Adult | 2,063 |
| 2016 | Yellow Perch | Juvenile | 3,630 |
| 2017 | Walleye | Large Fingerling | 1,280 |
| 2017 | Yellow Perch | Adult | 11,428 |
| 2018 | Walleye | Large Fingerling | 919 |
| 2018 | Yellow Perch | Adult | 5,379 |
| 2019 | Walleye | Juvenile | 140 |
| 2019 | Walleye | Small Fingerling | 9,150 |
| 2020 | Yellow Perch | Adult | 1,068 |
| 2021 | Walleye | Adult | 1,134 |
| 2021 | Walleye | Juvenile | 1,980 |
| 2021 | Yellow Perch | Adult | 2,441 |
| 2022 | Black Crappie | Adult | 1,041 |
| 2022 | Black Crappie | Juvenile | 939 |