# SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Henry, Kingsbury County LKT-Lake-55-003

2022

#### Lake Information

Name:	Henry	Maximum Depth:	8 Feet
County:	Kingsbury	Mean Depth:	4 Feet
Legal Description:	T110-R56-Sec. 13, 18, 19, 24		
Surface Area:	2,539 Acres		

#### **Surveys and Investigations**

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
AFS std gill net	Jun 28, 2022	6 net-nights

# **Common Fish Species Present**

Walleye

Yellow Perch

Northern Pike

Common Carp

Black Crappie

White Sucker

Black Bullhead

#### **Terminology**

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- **AFS std gill net** 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- std experimental gill net for non-Missouri River waters 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- std experimental gill net for Missouri River reservoirs 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$CPUE = \frac{number \, off ish}{effort}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{number \ of fish \ge quality \ length}{number \ of \ fish \ge stock \ length}\right) \ge 100$$

$$PSD - P = \left(\frac{number \ offish \ge preferred \ length}{number \ of \ fish \ge stock \ length}\right) \ge 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{Ws}\right) \ge 100$$

Confidence intervals (**CI**) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (**S**), quality (**Q**), preferred (**P**), memorable (**M**) and trophy (**T**). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

	St	ock	Qu	ality	Preferred		Mem	orable	Trophy	
Species Name	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

# Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80). \* Methods/Species that ignore stock length

			Abun	Abundance		ock Der	nsity Indic	es	Cor	ndition
Gear	Species	Sample Size (n)	CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
AFS std gill net	Black Bullhead	1	0.2	0.2	100		100			
	Black Crappie	3	0.5	0.3	100		100		103	3
	Common Carp	4	0.7	0.6	100		100			
	Northern Pike	5	0.8	0.5	60		40		87	8
	Walleye	13	2.2	0.5	100		38		81	4
	White Sucker	1	0.2	0.2	100		100			
	Yellow Perch	7	1.2	0.7	57		29		101	5

# 10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

\* Methods/Species that ignore stock length

							CPUE					
Gear	Species	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	Avg
AFS std gill net	Bigmouth Buffalo					0.0	0.0	0.3			0.0	0.08
	Black Bullhead					0.2	0.0	0.7			0.2	0.28
	Black Crappie					0.0	0.2	2.0			0.5	0.68
	Common Carp					3.0	0.8	4.5			0.7	2.25
	Northern Pike					0.5	1.0	1.5			0.8	0.95
	Walleye					3.3	1.7	9.0			2.2	4.05
	White Bass					0.2	2.0	0.0			0.0	0.55
	White Sucker					0.0	0.0	0.3			0.2	0.13
	Yellow Perch					0.2	3.0	6.0			1.2	2.60
std exp gill net	Bigmouth Buffalo		0.0	0.0	0.0							0.00
	Black Bullhead		7.3	6.0	1.7							5.00
	Black Crappie		3.3	2.3	0.3							1.97
	Common Carp		0.3	0.7	1.3							0.77
	Northern Pike		4.7	3.0	2.0							3.23
	Walleye		13.0	20.3	11.0							14.77
	White Bass		0.3	0.7	0.0							0.33
	Yellow Perch		13.0	24.0	8.0							15.00

----

# **10-Year Size Structure and Condition Statistics by Gear and Species**

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

		Year										
Gear	Species	Index	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
AFS std gill net	Black Bullhead	PSD					100		50			100
		PSD-P					100		25			100
	Black Crappie	PSD						100	8			100
		PSD-P						100	8			100
		Wr						93	130			103
	Common Carp	PSD					78	100	15			100
		PSD-P					17	60	11			100
	Northern Pike	PSD					100	83	78			60
		PSD-P					33	33	11			40
		Wr					83	81	88			87
	Walleye	PSD					50	80	67			100
		PSD-P					15	30	13			38
		Wr					82	89	91			81
	White Sucker	PSD							100			100
		PSD-P							100			100
	Yellow Perch	PSD					100	6	25			57
		PSD-P					100	0	14			29
		Wr					96	104	116			101
std exp gill net	Black Bullhead	PSD		14	78	20						
		PSD-P		0	0	20						
	Black Crappie	PSD		20	29	0						
		PSD-P		20	0	0						
		Wr		122	106	133						
	Common Carp	PSD		100	100	75						
		PSD-P		100	50	75						
	Northern Pike	PSD		86	56	83						
		PSD-P		21	44	0						
		Wr		85	84	77						
	Walleye	PSD		54	61	73						
		PSD-P		5	18	12						
		Wr		88	76	84						
	Yellow Perch	PSD		36	33	42						
		PSD-P		23	10	21						

						Year							
Gear	Species	Index	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	
std exp gill net	Yellow Perch	Wr		112	98	103							

# Length at Capture

Mean length at capture by age across years sampled, sample size (N).

# Species: Walleye

				Mean Leng	gth (expa	anded sam	ole numbe	er) at capt	ure by age	;	
Year	Ν	1	2	3	4	5	6	7	8	9	10+
2018	15	233 (5)	345 (2)	405 (4)		674 (1)			510 (2)		657 (1)
Species: Y	ellow Pe	erch		Mean Len	gth (expa	anded sam	ole numbe	er) at capt	ure by age	;	
Year	N	1	2	3	4	5	6	7	8	9	10+
2018	17	148 (16)	248 (1)								


# Fish Condition


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

					Length	Group	S		
			S-Q		Q-P		P-M		М
Species	Year	N	Wr (SE)	Ν	Wr (SE)	Ν	Wr (SE)	Ν	Wr (SE)
Northern Pike Gill Net	2018	1	98	3	81 (4.4)	2	71 (1.7)	0	
	2019	2	91 (2.0)	6	87 (4.1)	1	83	0	
	2022	2	73 (0.1)	1	88	2	101 (4.4)	0	
Walleye Gill Net	2018	2	90 (1.1)	5	90 (4.1)	1	96	2	82 (6.7)
	2019	18	87 (3.3)	29	93 (1.1)	5	94 (2.6)	2	89 (13.5)
	2022	0		8	83 (1.6)	4	85 (2.6)	1	48
Yellow Perch Gill Net	2018	17	104 (2.4)	1	101	0		0	
	2019	27	115 (4.0)	4	113 (13.5)	4	127 (6.6)	1	116
	2022	3	93 (6.3)	2	101 (7.2)	1	112	1	111


#### **Length Frequency Distribution**


Length frequency histogram of species sampled by year.






Species: Walleye Gear: AFS std gill net






Species: Yellow Perch Gear: AFS std gill net




#### **Historic Fish Sizes and Relative Abundance**


Size distribution per net by color for species sampled by year.

#### Species: Black Bullhead Gear: AFS std gill net





Species: Black Bullhead Gear: std exp gill net







Species: Common Carp Gear: std exp gill net









Species: Northern Pike Gear: std exp gill net




Species: Walleye Gear: AFS std gill net




Species: Walleye Gear: std exp gill net




Species: White Sucker Gear: AFS std gill net



Species: Yellow Perch Gear: AFS std gill net





# Fish Stocking

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2012	Walleye	Fry	1,200,000
2013	Walleye	Fry	1,161,000
2014	Walleye	Fry	1,160,000
2015	Walleye	Fry	1,200,000
2018	Walleye	Fry	1,200,000
2019	Walleye	Fry	1,200,000
2021	Walleye	Fry	1,200,000