

SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Lynn, Day County

MUD-Lake-351-001

2022

Lake Information

Name: Lynn

County: Day

Surface Area: 255 Acres

Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
AFS std gill net	Sep 07, 2022	4 net-nights
AFS std gill net	Sep 08, 2022	4 net-nights
AFS std gill net	Sep 09, 2022	4 net-nights
frame net (std 3/4 in)	Sep 07, 2022	6 net-nights
frame net (std 3/4 in)	Sep 08, 2022	6 net-nights
frame net (std 3/4 in)	Sep 09, 2022	6 net-nights

Common Fish Species Present

Bluegill

Yellow Perch

Walleye

Smallmouth Bass

Northern Pike

Rock Bass

Black Crappie

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- **AFS std gill net** – 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- **std experimental gill net for non-Missouri River waters** - 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- **std experimental gill net for Missouri River reservoirs** – 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$CPUE = \frac{\text{number of fish}}{\text{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{\text{number of fish} \geq \text{quality length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

$$PSD - P = \left(\frac{\text{number of fish} \geq \text{preferred length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (**W**) compared to a length-specific standard weight (**Ws**) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{W_s} \right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (S), quality (Q), preferred (P), memorable (M) and trophy (T). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

* **Methods/Species that ignore stock length**

Gear	Species	Sample Size (n)	Abundance		Stock Density Indices			Condition		
			CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
AFS std gill net	Northern Pike	9	0.8	0.3	100		22		87	2
	Rock Bass	2	0.2	0.2	50		0		110	2
	Smallmouth Bass	2	0.2	0.2	0		0		104	4
	Walleye	75	6.1	0.9	58	8	18	7	87	1
	Yellow Perch	127	10.5	3.2	6	3	1		114	1
frame net (std 3/4 in)	Black Crappie	4	0.1	0.1	0		0		123	
	Bluegill	214	11.9	5.4	12	3	2		120	1
	Northern Pike	6	0.2	0.1	100		0		89	3
	Smallmouth Bass	20	1.0	0.5	72	18	72	18	107	3
	Walleye	35	1.9	0.8	80	11	46	13	85	2
	Yellow Perch	61	3.4	3.2	0		0		105	1

10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

* **Methods/Species that ignore stock length**

Gear	Species	CPUE										Avg
		2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	
AFS std gill net	Black Bullhead									0.1	0.0	0.05
	Black Crappie									0.1	0.0	0.05
	Bluegill									0.2	0.0	0.10
	Northern Pike									0.8	0.8	0.80
	Rock Bass									0.7	0.2	0.45
	Smallmouth Bass									0.0	0.2	0.10
	Walleye									4.6	6.1	5.35
	Yellow Perch									6.5	10.5	8.50
frame net (std 3/4 in)	Black Crappie										0.1	0.10
	Bluegill										11.9	11.90
	Northern Pike										0.2	0.20
	Smallmouth Bass										1.0	1.00
	Walleye										1.9	1.90
	Yellow Perch										3.4	3.40

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

Gear	Species	Index	Year											
			2013	2014	2015	2016	2017	2018	2019	2020	2021	2022		
AFS std gill net	Black Crappie	PSD										0		
		PSD-P										0		
		Wr										110		
	Bluegill	PSD										0		
		PSD-P										0		
		Wr										106		
	Northern Pike	PSD										100	100	
		PSD-P										0	22	
		Wr										91	87	
	Rock Bass	PSD										25	50	
		PSD-P										0	0	
		Wr										106	110	
	Smallmouth Bass	PSD											0	
		PSD-P											0	
		Wr											104	
	Walleye	PSD										67	58	
		PSD-P										24	18	
		Wr										85	87	
Yellow Perch	PSD										19	6		
	PSD-P										14	1		
	Wr										96	114		
frame net (std 3/4 in)	Black Crappie	PSD											0	
		PSD-P											0	
		Wr											123	
	Bluegill	PSD												12
		PSD-P												2
		Wr												120
	Northern Pike	PSD												100
		PSD-P												0
		Wr												89
	Smallmouth Bass	PSD												72
		PSD-P												72
		Wr												107

Gear	Species	Index	Year										
			2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	
frame net (std 3/4 in)	Walleye	PSD											80
		PSD-P											46
		Wr											85
	Yellow Perch	PSD											0
		PSD-P											0
		Wr											105

Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Bluegill

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2022	213	119 (193)	182 (20)								

Species: Walleye

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2022	75	275 (24)	389 (18)	449 (2)	486 (25)	524 (1)	578 (2)		543 (1)		663 (2)
2021	55	318 (17)	415 (5)	438 (15)	462 (4)	494 (1)	578 (1)	583 (1)			629 (11)

Species: Yellow Perch

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2022	126	174 (120)	225 (6)								
2021	78	169 (67)		280 (10)	311 (1)						

Fish Condition


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

Species	Year	Length Groups							
		S-Q		Q-P		P-M		M	
		N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Black Crappie Frame Net	2022	1	123	0		0		0	
Bluegill Frame Net	2022	189	120 (0.7)	21	125 (1.7)	4	100 (26.8)	0	
Northern Pike Gill Net	2021	0		9	91 (2.5)	0		0	
	2022	0		7	88 (2.0)	2	81	0	
Walleye Gill Net	2021	18	83 (2.4)	24	85 (1.4)	8	87 (1.9)	5	91 (2.7)
	2022	31	86 (1.2)	29	88 (1.2)	12	88 (1.6)	1	90
Yellow Perch Gill Net	2021	63	96 (0.9)	4	96 (3.3)	10	97 (1.5)	1	89
	2022	119	114 (0.8)	6	109 (2.8)	1	105	0	



Length Frequency Distribution

Length frequency histogram of species sampled by year.


Species: Bluegill
Gear: frame net (std 3/4 in)




Species: Walleye
Gear: AFS std gill net



Species: Yellow Perch
Gear: AFS std gill net







Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.


Species: Black Crappie
Gear: frame net (std 3/4 in)




Species: Bluegill
Gear: frame net (std 3/4 in)



Species: Northern Pike
Gear: AFS std gill net



Species: Walleye
Gear: AFS std gill net



Species: Yellow Perch
Gear: AFS std gill net

