

SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Berry, Gregory County

PON-Lake-89-000

2022

Lake Information

Name: Berry **Maximum Depth:** 17 Feet
County: Gregory **Mean Depth:** 9 Feet
Legal Description: T96-R70-S32
Surface Area: 18 Acres

Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
boat shocker (night)	Oct 17, 2022	1487 seconds
frame net (std 3/4 in)	Dec 19, 2022	1 net-nights
frame net (std 3/4 in)	Jun 27, 2022	7 net-nights

Common Fish Species Present

Largemouth Bass

Bluegill

Black Crappie

Yellow Perch

Black Bullhead

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- **AFS std gill net** – 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- **std experimental gill net for non-Missouri River waters** - 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- **std experimental gill net for Missouri River reservoirs** – 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$CPUE = \frac{\text{number of fish}}{\text{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{\text{number of fish} \geq \text{quality length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

$$PSD - P = \left(\frac{\text{number of fish} \geq \text{preferred length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (*W*) compared to a length-specific standard weight (*Ws*) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{W_s} \right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (S), quality (Q), preferred (P), memorable (M) and trophy (T). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

* **Methods/Species that ignore stock length**

Gear	Species	Sample Size (n)	Abundance		Stock Density Indices			Condition		
			CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
boat shocker (night)	Largemouth Bass	19	28.7	21.3	60		50	28	115	4
frame net (std 3/4 in)	Black Bullhead	22	2.8	1.2	100		68	16	94	2
	Black Crappie	100	12.3	9.1	100		32	7	105	1
	Bluegill	120	15.0	5.7	43	6	2		111	2
	Largemouth Bass	5	0.6	0.6	40		20		119	30
	Yellow Perch	32	4.0	3.4	81	11	9		90	2

10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

* Methods/Species that ignore stock length

Gear	Species	CPUE										Avg
		2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	
boat shocker (night)	Black Crappie				0.0			0.0			0.0	0.00
	Bluegill				0.0			0.0			0.0	0.00
	Largemouth Bass				120.0			65.6			28.7	71.43
	Yellow Perch				0.0			0.0			0.0	0.00
frame net (std 3/4 in)	Black Bullhead				0.3			10.5			2.8	4.53
	Black Crappie				1.6			3.3			12.3	5.73
	Bluegill				27.3			18.0			15.0	20.10
	Largemouth Bass				0.1			0.0			0.6	0.23
	Yellow Perch				4.9			2.6			4.0	3.83

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

Gear	Species	Index	Year									
			2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
boat shocker (night)	Largemouth Bass	PSD				27				37		60
		PSD-P				9				29		50
		Wr				105				120		115
frame net (std 3/4 in)	Black Bullhead	PSD				33				56		100
		PSD-P				33				15		68
		Wr				96				97		94
	Black Crappie	PSD				100				31		100
		PSD-P				75				0		32
		Wr				96				116		105
	Bluegill	PSD				82				56		43
		PSD-P				30				1		2
		Wr				110				113		111
	Largemouth Bass	PSD				100				0		40
		PSD-P				100				0		20
		Wr				98						119
Yellow Perch	PSD				43				48		81	
	PSD-P				4				5		9	
	Wr				91				92		90	

Back-Calculated Lengths

Mean species back-calculated total length (mm) at age, standard error (SE), and sample size (N).

Species: Black Crappie

Year Class	Age	N	Mean back-calculated length (SE) at age																	
			1	2	3	4	5	6	7	8	9	10								
2021	1	1	81																	
2019	3	10	71 (2.4)	148 (5.8)	201 (4.8)															
2018	4	9	76 (2)	134 (11.5)	181 (8.6)	220 (4.8)														
2017	5	9	73 (2.5)	125 (7.2)	171 (5.2)	211 (6.5)	237 (3.9)													
2016	6	2	81 (.7)	136 (13)	178 (11.5)	211 (0)	231 (3.2)	248 (4.7)												
Weighted Mean		31	74	136	184	215	236	248												
Year Class	Age	N	11	12	13	14	15	16	17	18	19	20								
2021	1	1																		
2019	3	10																		
2018	4	9																		
2017	5	9																		
2016	6	2																		
Weighted Mean		31																		

Species: Bluegill

Year Class	Age	N	Mean back-calculated length (SE) at age											
			1	2	3	4	5	6	7	8	9	10		
2021	1	2	77 (15.2)											
2020	2	22	57 (3.6)	109 (4.5)										
2019	3	16	57 (3.2)	113 (5.8)	154 (4.6)									
2018	4	3	52 (3.4)	108 (8.9)	144 (4.6)	170 (2.6)								
2017	5	2	54 (8.5)	83 (9.8)	124 (7.4)	160 (11.2)	190 (7.8)							
Weighted Mean		45	57	109	150	166	190							

Year Class	Age	N	11	12	13	14	15	16	17	18	19	20
2021	1	2										
2020	2	22										
2019	3	16										
2018	4	3										
2017	5	2										
Weighted Mean		45										

Species: Largemouth Bass

Year Class	Age	N	Mean back-calculated length (SE) at age										
			1	2	3	4	5	6	7	8	9	10	
2022	0	1											
2021	1	1	92										
2020	2	6	81 (5.7)	151 (9.6)									
2019	3	2	78 (3.1)	121 (2.3)	183 (17.4)								
2018	4	1	77	112	167	302							
2016	6	3	86 (8.5)	160 (30.2)	260 (47.7)	331 (40.6)	397 (23)	440 (16)					
Weighted Mean		14	82	145	219	324	397	440					

Year Class	Age	N	Mean back-calculated length (SE) at age										
			11	12	13	14	15	16	17	18	19	20	
2022	0	1											
2021	1	1											
2020	2	6											
2019	3	2											
2018	4	1											
2016	6	3											
Weighted Mean		14											

Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Black Crappie

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2022	99	124 (1)		230 (29)	236 (36)	254 (30)	259 (4)				
2016	16		218 (1)	259 (14)	262 (1)						

Species: Bluegill

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2022	96	111 (3)	138 (48)	166 (40)	186 (4)	213 (2)					
2019	144	105 (7)	126 (48)	160 (85)	150 (3)	232 (1)					
2016	273	89 (4)	146 (68)	175 (61)	189 (47)	201 (54)	213 (6)	209 (23)		213 (12)	

Species: Largemouth Bass

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2022	14	131 (1)	193 (6)	236 (2)	366 (1)		478 (4)				
2019	66	198 (36)	232 (13)	302 (2)	297 (9)	410 (3)			484 (2)		533 (1)
2016	133	177 (46)	265 (48)	293 (31)		435 (1)		498 (1)		477 (3)	497 (4)

Fish Condition



Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

Species	Year	Length Groups							
		S-Q		Q-P		P-M		M	
		N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Black Crappie Frame Net	2019	18	120 (2.5)	8	107 (2.9)	0		0	
	2022	0		67	105 (0.5)	31	104 (0.9)	0	
Bluegill Frame Net	2019	63	117 (1.9)	80	111 (1.2)	1	110	0	
	2022	69	113 (2.4)	49	109 (1.2)	2	107 (3.1)	0	
Largemouth Bass Electro Fishing	2019	24	122 (2.0)	3	122 (1.3)	10	115 (3.4)	1	120
	2022	4	115 (4.2)	1	103	4	117 (6.7)	1	118



Length Frequency Distribution

Length frequency histogram of species sampled by year.



Species: Black Crappie
Gear: frame net (std 3/4 in)



Species: Bluegill
Gear: frame net (std 3/4 in)




Species: Largemouth Bass
Gear: boat shocker (night)




Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.

Species: Black Crappie
Gear: frame net (std 3/4 in)



Species: Bluegill
Gear: frame net (std 3/4 in)



Species: Largemouth Bass
Gear: boat shocker (night)

