# SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

**Drywood North, Roberts County** 

UMN-Lake-476-000

2022

#### Lake Information

Name:	Drywood North	Maximum Depth:	15 Feet
County:	Roberts		
Surface Area:	2,694 Acres		

#### Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort	
AFS std gill net	May 24, 2022	6 net-nights	
AFS std gill net	May 25, 2022	6 net-nights	

# **Common Fish Species Present**

Yellow Perch

Walleye

Northern Pike

Black Bullhead

Bluegill

Largemouth Bass

#### **Terminology**

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- **AFS std gill net** 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- std experimental gill net for non-Missouri River waters 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- std experimental gill net for Missouri River reservoirs 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$CPUE = \frac{number \, off ish}{effort}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{number \ of fish \ge quality \ length}{number \ of \ fish \ge stock \ length}\right) \ge 100$$

$$PSD - P = \left(\frac{number \ offish \ge preferred \ length}{number \ of \ fish \ge stock \ length}\right) \ge 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{Ws}\right) \ge 100$$

Confidence intervals (**CI**) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (**S**), quality (**Q**), preferred (**P**), memorable (**M**) and trophy (**T**). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

	St	Stock Quality Preferred		erred	Mem	orable	Trophy			
Species Name	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

#### Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80). \* Methods/Species that ignore stock length

			Abun	Abundance			Stock Density Indices			
Gear	Species	Sample Size (n)	CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
AFS std gill net	Black Bullhead	49	4.1	0.9	92		22	9	102	2
	Bluegill	2	0.2	0.2	100		100		124	2
	Largemouth Bass	1	0.1	0.1	100		0		130	
	Northern Pike	95	7.9	1.3	88	5	19	6	83	1
	Walleye	101	8.4	2.0	88	5	24	6	89	1
	Yellow Perch	24	2.0	0.5	75	14	46	16	95	2

#### 10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

\* Methods/Species that ignore stock length

							CPUE					
Gear	Species	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	Avg
AFS std gill n	et Black Bullhead										4.1	4.10
	Bluegill										0.2	0.20
	Largemouth Bass										0.1	0.10
	Northern Pike										7.9	7.90
	Walleye										8.4	8.40
	Yellow Perch										2.0	2.00

#### **10-Year Size Structure and Condition Statistics by Gear and Species**

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

							Ye	ar				
Gear	Species	Index	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
AFS std gill net	Black Bullhead	PSD										92
		PSD-P										22
		Wr										102
	Bluegill	PSD										100
		PSD-P										100
		Wr										124
	Largemouth Bass	PSD										100
		PSD-P										0
		Wr										130
	Northern Pike	PSD										88
		PSD-P										19
		Wr										83
	Walleye	PSD										88
		PSD-P										24
		Wr										89
	Yellow Perch	PSD										75
		PSD-P										46
		Wr										95

#### Length at Capture

Mean length at capture by age across years sampled, sample size (N).

#### Species: Walleye

				Mean Len	gth (expa	nded sam	ple numbe	er) at capt	ture by ag	е	
Year	Ν	1	2	3	4	5	6	7	8	9	10+
2022	101		361 (16)	446 (21)	469 (36)	532 (10)	573 (3)		663 (2)	624 (1)	662 (11)
Species: Y	ellow Per	ch		Mean Len	gth (expa	nded sam	ple numbe	er) at cap	ture by ag	e	
Year	N	1	2	3	4	5	6	7	8	9	10+
2022	24		141 (1)	210 (13)	270 (10)						


### Fish Condition


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

		Length Groups									
			S-Q		Q-P		P-M		М		
Species	Year	N	Wr (SE)	Ν	Wr (SE)	Ν	Wr (SE)	Ν	Wr (SE)		
Black Bullhead Gill Net	2022	4	91 (5.9)	34	101 (1.6)	11	109 (3.5)	0			
Northern Pike Gill Net	2022	11	82 (1.4)	66	81 (0.8)	14	89 (2.0)	4	89 (5.0)		
Walleye Gill Net	2022	12	90 (0.7)	65	89 (0.8)	14	89 (2.1)	10	92 (2.5)		
Yellow Perch Gill Net	2022	6	102 (2.6)	7	96 (3.4)	11	91 (1.3)	0			


#### **Length Frequency Distribution**

Length frequency histogram of species sampled by year.







Species: Northern Pike Gear: AFS std gill net



Species: Walleye Gear: AFS std gill net




Species: Yellow Perch Gear: AFS std gill net




#### **Historic Fish Sizes and Relative Abundance**

Size distribution per net by color for species sampled by year.


#### Species: Black Bullhead Gear: AFS std gill net




Species: Northern Pike Gear: AFS std gill net



Species: Walleye Gear: AFS std gill net



Species: Yellow Perch Gear: AFS std gill net



# Fish Stocking

Number of fish stocked by year, species, and size.

2012 Walleye	Fry	600,000
2014 Walleye	Fry	600,000
2016 Walleye	Fry	600,000
2018 Walleye	Fry	600,000