

SOUTH DAKOTA STATEWIDE FISHERIES SURVEY
Campbell, Campbell County
WMC-Lake-891-000
2022

Lake Information

Name:	Campbell	Maximum Depth:	22 Feet
County:	Campbell	Mean Depth:	9 Feet
Legal Description:	T126-R77-S11		
Surface Area:	42 Acres		

Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
boat shocker (night)	Oct 05, 2022	3600 seconds
frame net (std 3/4 in)	Jun 02, 2022	5 net-nights

Common Fish Species Present

Largemouth Bass

Black Crappie

Black Bullhead

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- **AFS std gill net** – 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- **std experimental gill net for non-Missouri River waters** - 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- **std experimental gill net for Missouri River reservoirs** – 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$CPUE = \frac{\text{number of fish}}{\text{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{\text{number of fish} \geq \text{quality length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

$$PSD - P = \left(\frac{\text{number of fish} \geq \text{preferred length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (**W**) compared to a length-specific standard weight (**Ws**) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{W_s} \right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (S), quality (Q), preferred (P), memorable (M) and trophy (T). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

* **Methods/Species that ignore stock length**

Gear	Species	Sample Size (n)	Abundance		Stock Density Indices			Condition		
			CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
boat shocker (night)	Largemouth Bass	85	6.0	3.2	100		0		111	5
frame net (std 3/4 in)	Black Bullhead	211	42.2	23.7	17	4	0		90	2
	Black Crappie	231	46.2	14.2	42	4	3	2	105	1
	Largemouth Bass	2	0.4	0.6	50		0		109	2

10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

* **Methods/Species that ignore stock length**

Gear	Species	CPUE										Avg
		2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	
AFS std frame net	Black Bullhead					146.2						146.2
	Black Crappie					68.9						68.90
	Yellow Perch					1.0						1.00
boat shocker (night)	Largemouth Bass		2.0								6.0	4.00
frame net (std 3/4 in)	Black Bullhead		8.7					51.0			42.2	33.97
	Black Crappie		10.0					4.0			46.2	20.07
	Common Carp		0.1					0.0			0.0	0.03
	Largemouth Bass		0.0					0.0			0.4	0.13
	White Crappie		0.0					0.0			0.0	0.00
	Yellow Perch		2.2					0.0			0.0	0.73

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

Gear	Species	Index	Year										
			2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	
AFS std frame net	Black Bullhead	PSD						0					
		PSD-P						0					
		Wr						94					
	Black Crappie	PSD						7					
		PSD-P						0					
		Wr						108					
boat shocker (night)	Largemouth Bass	PSD		100									100
		PSD-P		0									0
		Wr		113									111
frame net (std 3/4 in)	Black Bullhead	PSD		17						57			17
		PSD-P		0						0			0
		Wr		92							90		90
	Black Crappie	PSD		55							11		42
		PSD-P		1							0		3
		Wr		111							126		105
	Largemouth Bass	PSD											50
		PSD-P											0
		Wr											109

Back-Calculated Lengths

Mean species back-calculated total length (mm) at age, standard error (SE), and sample size (N).

Species: Black Crappie

Year Class	Age	N	Mean back-calculated length (SE) at age											
			1	2	3	4	5	6	7	8	9	10		
2019	3	8	107 (3.6)	139 (3.5)	170 (3.5)									
2018	4	2	112 (4.2)	130 (1.8)	151 (.4)	175 (3.8)								
2017	5	19	104 (2.9)	129 (3.1)	152 (3.5)	172 (3.8)	196 (4.4)							
2016	6	5	94 (6.6)	125 (11.8)	159 (15.1)	184 (15.3)	207 (16.7)	233 (13.9)						
Weighted Mean		34	104	131	157	175	198	233						
Year Class	Age	N	11	12	13	14	15	16	17	18	19	20		
2019	3	8												
2018	4	2												
2017	5	19												
2016	6	5												
Weighted Mean		34												

Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Black Crappie

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2022	231			184 (45)	189 (18)	201 (150)	234 (18)				
2017	689			170 (620)	194 (69)						
2014	200		199 (198)			297 (2)					

Species: Largemouth Bass

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2014	4	314 (4)									

Fish Condition


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

Species	Year	Length Groups							
		S-Q		Q-P		P-M		M	
		N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Black Crappie Frame Net	2019	32	129 (3.1)	4	107 (1.7)	0		0	
	2022	134	106 (1.4)	90	103 (1.6)	7	102 (2.0)	0	
Largemouth Bass Electro Fishing	2022	0		6	111 (3.9)	0		0	



Length Frequency Distribution

Length frequency histogram of species sampled by year.


Species: Black Crappie
Gear: AFS std frame net



Species: Black Crappie
Gear: frame net (std 3/4 in)




Species: Largemouth Bass
Gear: boat shocker (night)




Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.


Species: Black Crappie
Gear: AFS std frame net



Species: Black Crappie
Gear: frame net (std 3/4 in)



Species: Largemouth Bass
Gear: boat shocker (night)



Fish Stocking

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2011	Black Crappie	Adult	110
2011	Channel Catfish	Adult	147
2012	Black Crappie	Adult	90
2012	Largemouth Bass	Juvenile	200
2012	Yellow Perch	Adult	300
2014	Bluegill	Adult	100
2014	Channel Catfish	Juvenile	100
2014	Largemouth Bass	Fingerling	1,260
2014	Largemouth Bass	Large Fingerling	1,000
2015	Bluegill	Adult	300
2015	Largemouth Bass	Juvenile	292
2016	Channel Catfish	Adult	120
2016	Largemouth Bass	Adult	115
2017	Largemouth Bass	Adult	111
2018	Channel Catfish	Adult	112
2019	Channel Catfish	Adult	107
2019	Largemouth Bass	Adult	30
2019	Largemouth Bass	Juvenile	77
2020	Channel Catfish	Adult	103
2020	Largemouth Bass	Juvenile	100
2021	Channel Catfish	Adult	138
2022	Largemouth Bass	Juvenile	700