#### SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Lehrman Slough, McCook County LJA-Lake-725-000 2023

#### **Lake Information**

Name: Lehrman Slough

County: McCook

Surface Area: 200 Acres

#### **Surveys and Investigations**

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
AFS std gill net	Jun 07, 2023	4 net-nights

## **Common Fish Species Present**

Yellow Perch

#### **Terminology**

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- **AFS std gill net** 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- std experimental gill net for non-Missouri River waters 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- std experimental gill net for Missouri River reservoirs 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$\mathit{CPUE} = \frac{\mathit{number of fish}}{\mathit{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{number\ of\ fish \ge quality\ length}{number\ of\ fish \ge stock\ length}\right) \times 100$$

$$\textit{PSD} - \textit{P} = \left(\frac{number\ of\ fish\ \geq preferred\ length}{number\ of\ fish\ \geq stock\ length}\right) \ge 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{Ws}\right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (S), quality (Q), preferred (P), memorable (M) and trophy (T). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

	St	ock	Qu	ality	Pref	erred	Mem	orable	Tro	ophy
Species Name	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

### 10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

\* Methods/Species that ignore stock length

							CPUE					
Gear	Species	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	Avg
AFS std gill net	Black Bullhead								17.0			17.00
	Common Carp								10.5			10.50
	Green Sunfish								0.3			0.30
	Walleye								1.8			1.80
	Yellow Perch								61.5			61.50
std exp gill net	Black Bullhead			63.3								63.30
	Common Carp			3.3								3.30
	Orangespotted Sunfish			0.0								0.00
	Walleye			0.3								0.30
	White Sucker			0.0								0.00
	Yellow Perch			27.7								27.70

### 10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

		Year										
Gear	Species	Index	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023
AFS std gill net	Yellow Perch	PSD	1				1			51		
		PSD-P								22		
		Wr								74		
std exp gill net	Yellow Perch	PSD			4							
		PSD-P			2							
		Wr			108							

### **Length at Capture**

Mean length at capture by age across years sampled, sample size (N).

Species: Yellow Perch

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2016	35	159 (32)		265 (3)							

### **Fish Condition**


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

			Length Groups									
		S-Q		Q-P		P-M		М				
Species	Year	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)			
Yellow Perch Gill Net	2021	121	64 (1.3)	71	78 (1.3)	54	83 (0.5)	0				

### **Length Frequency Distribution**

Length frequency histogram of species sampled by year.


Species: Yellow Perch Gear: AFS std gill net




#### **Historic Fish Sizes and Relative Abundance**

Size distribution per net by color for species sampled by year.

Species: Yellow Perch Gear: AFS std gill net



Species: Yellow Perch Gear: std exp gill net



# Fish Stocking

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2015	Yellow Perch	Small Fingerling	98,510
2019	Yellow Perch	Fry	86,500
2021	Walleye	Fry	200,000
2023	Walleye	Fry	100,000
2023	Yellow Perch	Juvenile	87,500