Hwy 81 East Lake Survey Summary

Hwy 81 East Lake, located 5 miles south of Arlington, SD, is managed as a walleye and yellow perch fishery; other fish species (e.g., black crappie, northern pike, smallmouth bass, and white bass) are also present.

- Walleye. Walleye abundance increased to 5.7 fish per gill net in 2023, resulting in one of the highest catch rates in the region. Relative abundance was higher than the previous four sample years and the long term mean (3.1 fish per net). Netted fish ranged in length from 10.6 to 26.8 inches with most (74%) measuring >15 inches. A large proportion (50%) also measured >20 inches. The sample was comprised of at least 7 different year classes of fish, but a few stood out amongst others. Age 4 and age 5 fish accounted for almost half of all fish sampled (18 and 24%, respectively). Growth of these two cohorts (produced in 2018 and 2019) is excellent with fish averaging 19.1 inches in length by age 4 and 20.3 inches by age 5. A mean condition score of 91 also indicates that these fish are healthy.
- Yellow Perch. Gill netting efforts produced the highest yellow perch catch rate in the region (21.8 fish per net in 2023). Relative abundance was considerably higher than the previous several sample years and the long term mean (9.1 fish per net). Sampled fish ranged in length from 5.1 to 13.0 inches with about half (58%) measuring >8 inches. A mean relative weight score of 94 indicates they were in good condition. Highway 81 East Lake is worth a look for any angler targeting yellow perch in the southeast region.

For more detailed results see the computer-generated South Dakota Statewide Fisheries Survey for Hwy 81 East Lake (below).

SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Hwy 81 Southeast, Brookings County MBS-Lake-233-003 2023

Lake Information

Name: Hwy 81 Southeast

County: Brookings

Surface Area: 141 Acres

Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
AFS std gill net	Aug 08, 2023	4 net-nights

Common Fish Species Present

Walleye

Yellow Perch

Yellow Bullhead

Common Carp

Smallmouth Bass

Northern Pike

Black Bullhead

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- AFS std gill net 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- std experimental gill net for non-Missouri River waters 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- std experimental gill net for Missouri River reservoirs 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$\mathit{CPUE} = \frac{\mathit{number of fish}}{\mathit{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{number\ of\ fish \ge quality\ length}{number\ of\ fish \ge stock\ length}\right) \times 100$$

$$\textit{PSD} - \textit{P} = \left(\frac{number\ of\ fish\ \geq preferred\ length}{number\ of\ fish\ \geq stock\ length}\right) \ge 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{Ws}\right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (S), quality (Q), preferred (P), memorable (M) and trophy (T). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

	St	ock	Qu	ality	Pref	erred	Mem	orable	Tro	ophy
Species Name	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

* Methods/Species that ignore stock length

			Abundance		St	Stock Density Indices				dition
Gear	Species	Sample Size (n)	CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
AFS std gill net	Black Bullhead	1	0.3	0.4	100		0			
	Common Carp	2	0.5	0.5	100		100			
	Northern Pike	1	0.3	0.4	100		0		77	
	Smallmouth Bass	1	0.3	0.4	0		0		104	
	Walleye	11	2.5	1.1	90		10		87	2
	Yellow Bullhead	4	1.0	1.2	100		100			
	Yellow Perch	38	9.5	1.6	39	12	5		96	3

10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

* Methods/Species that ignore stock length

							CPUE					
Gear	Species	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	Avg
AFS std gill net	Black Bullhead					,			2.3		0.3	1.30
	Black Crappie								1.3		0.0	0.65
	Common Carp								3.0		0.5	1.75
	Northern Pike								0.7		0.3	0.50
	Smallmouth Bass								0.0		0.3	0.15
	Walleye								1.0		2.5	1.75
	White Bass								4.0		0.0	2.00
	Yellow Bullhead								0.0		1.0	0.50
	Yellow Perch								3.0		9.5	6.25
std exp gill net	Black Bullhead	34.7	18.3	1.3								18.10
	Black Crappie	0.0	0.0	0.0								0.00
	Common Carp	1.0	0.3	1.7								1.00
	Northern Pike	0.0	0.0	0.0								0.00
	Smallmouth Bass	0.0	0.0	0.3								0.10
	Walleye	5.0	28.0	8.3								13.77
	White Bass	0.3	10.3	7.3								5.97
	White Sucker	0.3	0.7	0.0								0.33
	Yellow Bullhead	1.0	1.7	1.3								1.33
	Yellow Perch	38.0	32.3	35.0								35.10

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

							Ye	ar				
Gear	Species	Index	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023
AFS std gill net	Black Bullhead	PSD								71		100
		PSD-P								43		0
	Common Carp	PSD								100		100
		PSD-P								33		100
	Northern Pike	PSD								50		100
		PSD-P								0		0
		Wr								69		77
	Smallmouth Bass	PSD										0
		PSD-P										0
		Wr										104
	Walleye	PSD								100		90
		PSD-P								67		10
		Wr								73		87
	Yellow Bullhead	PSD										100
		PSD-P										100
	Yellow Perch	PSD								78		39
		PSD-P								0		5
		Wr								93		96
std exp gill net	Black Bullhead	PSD	55	60	100							
		PSD-P	6	7	75							
	Common Carp	PSD	67	100	100							
		PSD-P	67	0	20							
	Smallmouth Bass	PSD			0							
		PSD-P			0							
		Wr			111							
	Walleye	PSD	80	21	52							
		PSD-P	40	13	12							
		Wr	90	91	91							
	Yellow Bullhead	PSD	100	100	75							
		PSD-P	100	60	75							
	Yellow Perch	PSD	23	88	19							
		PSD-P	6	16	15							

			Year									
Gear	Species	Index	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023
std exp gill net	Yellow Perch	Wr	97	94	112	,		,				

Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Yellow Perch

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2015	96	179 (11)	231 (72)	263 (9)	282 (4)						
2014	114	166 (89)	233 (12)	245 (12)	265 (1)						

Fish Condition


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

			Length Groups										
			S-Q		Q-P		P-M		М				
Species	Year	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)				
Northern Pike	2021	1	70	1	69	0		0					
Gill Net	2023	0		1	77	0		0					
Walleye	2021	0		1	77	1	80	1	61				
Gill Net	2023	1	88	8	87 (1.6)	1	90	0					
Yellow Perch Gill Net	2021	2	92 (2.9)	7	93 (1.0)	0		0					
	2023	23	97 (2.6)	13	98 (1.5)	2	63 (27.3)	0					


Length Frequency Distribution

Length frequency histogram of species sampled by year.

Species: Walleye Gear: AFS std gill net




Species: Yellow Perch Gear: AFS std gill net




Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.


Species: Black Bullhead Gear: AFS std gill net




Species: Black Bullhead Gear: std exp gill net




Species: Common Carp Gear: AFS std gill net




Species: Common Carp Gear: std exp gill net




Species: Northern Pike Gear: AFS std gill net




Species: Walleye Gear: AFS std gill net




Species: Walleye Gear: std exp gill net



Species: Yellow Perch Gear: AFS std gill net



Species: Yellow Perch Gear: std exp gill net



Fish Stocking

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2021	Walleye	Juvenile	15,840
2022	Walleye	Juvenile	11,645
2023	Walleye	Juvenile	12,707