SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Clear, Hamlin County

UBS-Lake-175-001

2023

Lake Information

Name:	Clear	Maximum Depth:	13 Feet
County:	Hamlin		
Surface Area:	771 Acres		

Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort	
AFS std gill net	Jun 27, 2023	4 net-nights	
AFS std gill net	Jun 28, 2023	4 net-nights	
AFS std gill net	Jun 29, 2023	4 net-nights	

Common Fish Species Present

Walleye

Smallmouth Bass

Northern Pike

Yellow Perch

White Crappie

Common Carp

Black Bullhead

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- **AFS std gill net** 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- std experimental gill net for non-Missouri River waters 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- std experimental gill net for Missouri River reservoirs 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$\textit{CPUE} = \frac{\textit{number of fish}}{\textit{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{number \ of fish \ge quality \ length}{number \ of \ fish \ge stock \ length}\right) \ge 100$$

$$PSD - P = \left(\frac{number \ offish \ge preferred \ length}{number \ of \ fish \ge stock \ length}\right) \ge 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{Ws}\right) \ge 100$$

Confidence intervals (**CI**) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (**S**), quality (**Q**), preferred (**P**), memorable (**M**) and trophy (**T**). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

	St	ock	Qu	ality	Pref	ferred	Mem	orable	Trophy	
Species Name	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80). * Methods/Species that ignore stock length

			Abundance		St	ock Der	nsity Indic	es	Condition	
Gear	Species	Sample Size (n)	CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
AFS std gill net	Black Bullhead	2	0.2	0.2	50		50		102	5
	Common Carp	4	0.3	0.3	100		75		93	5
	Smallmouth Bass	1	0.1	0.1	0		0		114	
	Walleye	158	13.2	1.8	96	2	7	3	95	1
	White Crappie	42	2.2	0.9	8		8		107	2
	Yellow Perch	18	1.2	0.5	0		0		110	3

10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

* Methods/Species that ignore stock length

							CPUE					
Gear	Species	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	Avg
AFS std gill net	Black Bullhead				2.3			0.8			0.2	1.10
	Black Crappie				0.0			0.1			0.0	0.03
	Common Carp				2.6			0.2			0.3	1.03
	Northern Pike				0.0			0.1			0.0	0.03
	Smallmouth Bass				0.0			0.0			0.1	0.03
	Walleye				16.5			7.7			13.2	12.47
	White Crappie				0.0			0.5			2.2	0.90
	White Sucker				0.6			0.8			0.0	0.47
	Yellow Perch				3.6			14.0			1.2	6.27
std exp gill net	Black Bullhead	4.3										4.30
	Black Crappie	0.2										0.20
	Common Carp	4.2										4.20
	Northern Pike	0.3										0.30
	Walleye	25.7										25.70
	White Sucker	1.2										1.20
	Yellow Perch	20.7										20.70

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

		Year										
Gear	Species	Index	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023
AFS std gill net	Black Bullhead	PSD				85			89			50
		PSD-P				48			67			50
		Wr				101			111			102
	Common Carp	PSD				94			100			100
		PSD-P				94			100			75
		Wr				99			96			93
	Northern Pike	PSD							100			
		PSD-P							0			
		Wr							110			
	Smallmouth Bass	PSD										0
		PSD-P										0
		Wr										114
	Walleye	PSD				17			98			96
		PSD-P				11			11			7
		Wr				84			92			95
	White Crappie	PSD							33			8
		PSD-P							33			8
		Wr							106			107
	Yellow Perch	PSD				95			41			0
		PSD-P				37			2			0
		Wr				105			115			110
std exp gill net	Black Bullhead	PSD	100									
1.0		PSD-P	65									
		Wr	93									
	Common Carp	PSD	100									
	·	PSD-P	32									
		Wr	99									
	Northern Pike	PSD	100									
		PSD-P	100									
		Wr	81									
	Walleye	PSD	3									
		PSD-P	3									
		Wr	80									

			Year									
Gear	Species	Index	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023
std exp gill net	Yellow Perch	PSD	98									
		PSD-P	35									
		Wr	101									

Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Walleye

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2023	158	314 (6)	414 (33)	447 (3)	469 (103)	491 (3)	625 (2)	607 (1)	587 (3)		671 (4)
2020	121	196 (30)	358 (1)	419 (63)	475 (2)	474 (20)		634 (2)		632 (2)	702 (1)
2017	199	250 (4)	321 (163)	463 (1)	498 (10)	444 (1)	519 (18)	571 (1)	624 (1)		
2014	155			322 (150)		544 (5)					

Species: Yellow Perch

			ĺ	ple numb	er) at capt	ure by age	Э				
Year	N	1	2	3	4	5	6	7	8	9	10+
2023	18	135 (18)									
2020	170	143 (51)	201 (116)		299 (2)	277 (1)					
2017	43		220 (24)	252 (5)		304 (2)	304 (10)	301 (3)			
2014	124			242 (117)	284 (7)						


Fish Condition


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

					Length	Group	S		
			S-Q		Q-P		P-M		М
Species	Year	N	Wr (SE)	Ν	Wr (SE)	Ν	Wr (SE)	Ν	Wr (SE)
Black Bullhead Gill Net	2020	1	110	2	113 (8.6)	6	110 (7.0)	0	
	2023	1	106	0		1	98	0	
Common Carp	2020	0		0		1	94	1	97
Gill Net	2023	0		1	88	1	89	2	98 (5.2)
Northern Pike Gill Net	2020	0		1	110	0		0	
Walleye Gill Net	2020	2	90 (5.7)	80	92 (0.6)	6	89 (2.2)	4	89 (2.1)
	2023	6	100 (1.9)	141	95 (0.9)	6	89 (1.5)	5	94 (2.9)
Yellow Perch Gill Net	2020	99	115 (0.7)	66	115 (0.8)	2	102 (5.1)	1	99
	2023	14	110 (2.2)	0		0		0	


Length Frequency Distribution

Length frequency histogram of species sampled by year.






Species: Yellow Perch Gear: AFS std gill net




Historic Fish Sizes and Relative Abundance


Size distribution per net by color for species sampled by year.

Species: Black Bullhead Gear: AFS std gill net




Species: Black Bullhead Gear: std exp gill net







Species: Common Carp Gear: std exp gill net




Species: Northern Pike Gear: std exp gill net




Species: Walleye Gear: AFS std gill net




Species: Walleye Gear: std exp gill net



Species: Yellow Perch Gear: AFS std gill net





Fish Stocking

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2012	Smallmouth Bass	Fingerling	34,970
2013	Walleye	Fry	300,000
2015	Walleye	Fry	300,000
2017	Walleye	Fry	300,000
2019	Walleye	Fry	300,000
2021	Walleye	Fry	300,000
2023	Walleye	Fry	300,000