SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Diamond, Minnehaha County LBS-Lake-223-800 2023

Lake Information

Name: Diamond Maximum Depth: 12 Feet

County: Minnehaha Mean Depth: 8 Feet

Legal Description: T104N-R52W-Sec. 5

Surface Area: 295 Acres

Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
AFS std gill net	Jun 08, 2023	6 net-nights

Common Fish Species Present

Yellow Perch

Walleye

Black Bullhead

Bigmouth Buffalo

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- **AFS std gill net** 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- std experimental gill net for non-Missouri River waters 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- std experimental gill net for Missouri River reservoirs 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$\mathit{CPUE} = \frac{\mathit{number of fish}}{\mathit{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{number\ of\ fish \ge quality\ length}{number\ of\ fish \ge stock\ length}\right) \times 100$$

$$\textit{PSD} - \textit{P} = \left(\frac{number\ of\ fish\ \geq preferred\ length}{number\ of\ fish\ \geq stock\ length}\right) \ge 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{Ws}\right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (S), quality (Q), preferred (P), memorable (M) and trophy (T). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

	St	ock	Qu	ality	Preferred		Mem	orable	Tro	ophy
Species Name	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

* Methods/Species that ignore stock length

			Abun	dance	St	ock Der	sity Indic	es	Cor	ndition
Gear	Species	Sample Size (n)	CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
AFS std gill net	Bigmouth Buffalo	2	0.3	0.5	100		0			
	Black Bullhead	7	1.2	1.0	0		0			
	Yellow Perch	57	9.3	4.9	4		0		83	2

10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

* Methods/Species that ignore stock length

							CPUE					
Gear	Species	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	Avg
AFS std frame	Black Bullhead				2.4		,					2.40
net	Common Carp				1.2							1.20
	Walleye				3.0							3.00
	White Sucker				0.2							0.20
AFS std gill net	Bigmouth Buffalo				0.0	0.0	0.0		2.5		0.3	0.56
	Black Bullhead				13.7	8.5	3.3		35.3		1.2	12.40
	Common Carp				2.8	4.7	1.8		2.3		0.0	2.32
	Sunfish Hybrid				0.0	0.0	0.0		0.2		0.0	0.04
	Walleye				2.2	4.8	12.2		3.7		0.0	4.58
	White Sucker				0.0	0.0	0.0		0.3		0.0	0.06
	Yellow Bullhead				0.0	0.0	0.0		0.2		0.0	0.04
	Yellow Perch				8.0	6.0	1.5		12.5		9.3	6.02
std exp gill net	Black Bullhead	148.7	76.0	50.3								91.67
	Common Carp	14.3	7.7	12.0								11.33
	Sunfish Hybrid	0.0	0.7	0.3								0.33
	Walleye	0.0	4.3	6.7								3.67
	Yellow Perch	22.3	7.0	14.3								14.53

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

							Ye	ar				
Gear	Species	Index	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023
AFS std frame	Black Bullhead	PSD				100						
net		PSD-P				17						
	Walleye	PSD				33						
		PSD-P				13						
		Wr				80						
AFS std gill net	Bigmouth Buffalo	PSD								0		100
		PSD-P								0		0
	Black Bullhead	PSD				99	53	15		46		0
		PSD-P				4	14	0		0		0
	Walleye	PSD				46	86	88		68		
		PSD-P				8	10	18		18		
		Wr				84	98	100		87		
	Yellow Perch	PSD				100	19	67		27		4
		PSD-P				40	3	22		0		0
		Wr				92	106	110		108		83
std exp gill net	Black Bullhead	PSD	4	63	74							
		PSD-P	0	0	1							
	Walleye	PSD	0	8	20							
		PSD-P	0	8	0							
		Wr		77	79							
	Yellow Perch	PSD	0	57	70							
		PSD-P	0	0	35							
		Wr	105	100	93							

Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Walleye

				Mean Ler	gth (expa	nded sam	ple numb	er) at captı	ire by age)	
Year	N	1	2	3	4	5	6	7	8	9	10+
2019	79	257 (15)	397 (3)		465 (33)	486 (22)	542 (6)				
2018	29			386 (10)	392 (9)	451 (7)	611 (1)	575 (1)			591 (1)
2017	13		265 (1)	350 (8)	460 (3)	520 (1)					
Species: Y	ellow Pe	erch									
				Mean Ler	gth (expa	nded sam	ple numb	er) at captu	ire by age)	
Year	N	1	2	3	4	5	6	7	8	9	10+
2018	11		188 (11)								
2017	5			226 (3)	266 (1)		301 (1)				
2014	67	147 (67)									

Fish Condition


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).


			Length Groups									
			S-Q		Q-P		P-M		М			
Species	Year	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)			
Walleye Gill Net	2019	9	110 (1.3)	51	98 (0.9)	13	97 (1.9)	0				
	2021	7	87 (2.7)	11	85 (1.2)	4	94 (7.1)	0				
Yellow Perch Gill Net	2019	3	127 (10.5)	4	101 (1.4)	2	101 (0.7)	0				
	2021	55	109 (2.4)	20	106 (2.2)	0		0				
	2023	54	83 (1.2)	2	84 (3.5)	0		0				


Length Frequency Distribution

Length frequency histogram of species sampled by year.


Species: Black Bullhead Gear: AFS std gill net










Species: Walleye Gear: AFS std gill net









Species: Yellow Perch Gear: AFS std gill net









Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.


Species: Black Bullhead Gear: AFS std gill net




Species: Black Bullhead Gear: std exp gill net




Species: Walleye Gear: AFS std gill net




Species: Walleye Gear: std exp gill net



Species: Yellow Perch Gear: AFS std gill net



Species: Yellow Perch Gear: std exp gill net



Fish Stocking

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2018	Walleye	Large Fingerling	2,023
2019	Walleye	Small Fingerling	23,064
2022	Saugeye	Juvenile	33,250
2023	Yellow Perch	Adult	1,800