SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Island South, Minnehaha County LBS-Lake-213-001 2024

Lake Information

Name: Island South
County: Minnehaha

Surface Area: 125 Acres

Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
AFS std gill net	Jul 18, 2024	4 net-nights

Common Fish Species Present

Common Carp

Black Bullhead

Walleye

Smallmouth Bass

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- AFS std gill net 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- std experimental gill net for non-Missouri River waters 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- std experimental gill net for Missouri River reservoirs 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$\mathit{CPUE} = \frac{\mathit{number of fish}}{\mathit{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{number\ of\ fish \ge quality\ length}{number\ of\ fish \ge stock\ length}\right) \times 100$$

$$\textit{PSD} - \textit{P} = \left(\frac{number\ of\ fish\ \geq preferred\ length}{number\ of\ fish\ \geq stock\ length}\right) \ge 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{Ws}\right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (S), quality (Q), preferred (P), memorable (M) and trophy (T). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

	St	ock	Qu	ality	Pref	erred	Mem	orable	Tro	ophy
Species Name	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

* Methods/Species that ignore stock length

			Abun	dance	St	ock Der	sity Indic	es	Condition	
Gear	Species	Sample Size (n)	CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
AFS std gill net	Black Bullhead	71	17.8	6.0	54	9	4			
	Common Carp	94	19.0	3.5	47	8	1			
	Smallmouth Bass	1	0.3	0.4	0		0		115	
	Walleye	29	2.3	1.4	0		0		83	2

10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

^{*} Methods/Species that ignore stock length

							CPUE					
Gear	Species	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	Avg
AFS std frame	Black Bullhead			7.6								7.60
net	Black Crappie			2.4								2.40
	Bluegill			3.0								3.00
	Common Carp			3.0								3.00
	Golden Shiner			0.0								0.00
	Northern Pike			1.0								1.00
	Smallmouth Bass			0.2								0.20
	Walleye			1.6								1.60
	Yellow Perch			1.6								1.60
AFS std gill net	Black Bullhead			5.3	5.0	2.3			15.5		17.8	9.18
	Black Crappie			0.3	1.3	2.0			0.3		0.0	0.78
	Channel Catfish			0.0	0.0	0.0			1.8		0.0	0.36
	Common Carp			22.0	4.0	5.5			5.3		19.0	11.16
	Northern Pike			8.0	0.3	0.0			1.8		0.0	0.58
	Smallmouth Bass			0.0	0.0	0.0			0.3		0.3	0.12
	Walleye			6.3	13.3	15.3			8.8		2.3	9.20
	White Sucker			0.3	0.0	0.0			0.5		0.0	0.16
	Yellow Perch			2.8	2.5	4.0			8.0		0.0	2.02
frame net (std	Black Bullhead				4.3	2.6						3.45
3/4 in)	Black Crappie				7.3	19.0						13.15
	Bluegill				1.5	1.6						1.55
	Common Carp				2.0	0.8						1.40
	Green Sunfish				0.0	0.2						0.10
	Northern Pike				1.3	0.2						0.75
	Sunfish Hybrid				0.0	0.0						0.00
	Walleye				0.5	3.0						1.75
	White Sucker				0.3	0.0						0.15
	Yellow Perch				1.3	1.4						1.35
std exp gill net	Black Bullhead	101.3	58.0									79.65
	Black Crappie	0.0	1.0									0.50
	Common Carp	5.3	2.7									4.00
	Golden Shiner	0.0	0.0									0.00
	Northern Pike	9.0	1.3									5.15
	Walleye	2.3	4.7									3.50

1/20/2025 Page 6

							CPUE					
Gear	Species	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	Avg
std exp gill net	White Sucker	0.0	0.3									0.15
	Yellow Perch	6.0	7.7									6.85

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

							Ye	ear				
Gear	Species	Index	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
AFS std frame	Black Bullhead	PSD			79							
net		PSD-P			16							
	Common Carp	PSD			87							
		PSD-P			87							
	Smallmouth Bass	PSD			100							
		PSD-P			100							
		Wr			114							
	Walleye	PSD			63							
		PSD-P			0							
		Wr			96							
AFS std gill net	Black Bullhead	PSD			95	95	100			5		54
		PSD-P			19	15	0			2		4
	Common Carp	PSD			11	100	68			95		47
		PSD-P			10	6	5			14		1
	Smallmouth Bass	PSD								100		0
		PSD-P								100		0
		Wr								88		115
	Walleye	PSD			60	92	43			100		0
		PSD-P			36	19	10			40		0
		Wr			84	92	92			89		83
frame net (std	Black Bullhead	PSD				82	92					
3/4 in)		PSD-P				24	23					
	Common Carp	PSD				100	75					
		PSD-P				75	75					
	Walleye	PSD				100	93					
		PSD-P				0	87					
		Wr				90	90					
std exp gill net	Black Bullhead	PSD	27	86								
		PSD-P	0	0								
	Common Carp	PSD	13	75								
	·	PSD-P	0	13								
	Walleye	PSD	100	57								
	•											

			Year									
Gear	Species	Index	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
std exp gill net	Walleye	PSD-P	86	36								
		Wr	96	83								

Fish Condition


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).


	S								
		S-Q			Q-P	P-M			M
Species	Year	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Walleye Gill Net	2022	0		21	90 (1.6)	12	88 (1.4)	2	82 (2.8)
	2024	9	83 (1.2)	0		0		0	

Length Frequency Distribution


Length frequency histogram of species sampled by year.


Species: Black Bullhead Gear: AFS std gill net







Species: Common Carp Gear: AFS std gill net









Species: Walleye Gear: AFS std gill net









Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.


Species: Black Bullhead Gear: AFS std gill net




Species: Black Bullhead Gear: std exp gill net




Species: Common Carp Gear: AFS std gill net




Species: Common Carp Gear: std exp gill net



Species: Walleye Gear: AFS std gill net



Species: Walleye Gear: std exp gill net



Fish Stocking

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2013	Yellow Perch	Small Fingerling	138,250
2014	Walleye	Small Fingerling	14,300
2015	Walleye	Juvenile	666
2015	Walleye	Small Fingerling	5,082
2016	Walleye	Juvenile	703
2016	Walleye	Small Fingerling	5,670
2016	Yellow Perch	Adult	10,187
2017	Yellow Perch	Adult	3,362
2018	Yellow Perch	Adult	3,313
2020	Yellow Perch	Adult	1,454
2021	Walleye	Adult	1,074
2023	Walleye	Fry	150,000
2023	Yellow Perch	Adult	900
2024	Walleye	Juvenile	160
2024	Yellow Perch	Juvenile	1,282