# SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Goldsmith, Brookings County

MBS-Lake-236-000

2024

#### Lake Information

Name:	Goldsmith	Maximum Depth:	9 Feet
County:	Brookings	Mean Depth:	6 Feet
Legal Description:	T110N-R51W-Sec 9,16		
Surface Area:	308 Acres		

#### **Surveys and Investigations**

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
AFS std gill net	Jul 24, 2024	6 net-nights

# **Common Fish Species Present**

Yellow Perch Walleye Bigmouth Buffalo Common Carp Saugeye Northern Pike Black Bullhead White Sucker Black Crappie

#### **Terminology**

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- **AFS std gill net** 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- std experimental gill net for non-Missouri River waters 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- std experimental gill net for Missouri River reservoirs 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$CPUE = \frac{number \, off ish}{effort}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{number \ of fish \ge quality \ length}{number \ of \ fish \ge stock \ length}\right) \ge 100$$

$$PSD - P = \left(\frac{number \ offish \ge preferred \ length}{number \ of \ fish \ge stock \ length}\right) \ge 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{Ws}\right) \ge 100$$

Confidence intervals (**CI**) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (**S**), quality (**Q**), preferred (**P**), memorable (**M**) and trophy (**T**). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

	St	ock	Qu	ality	Pref	ferred	Mem	orable	Tro	ophy
Species Name	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

## Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80). \* Methods/Species that ignore stock length

			Abun	dance	St	ock Der	nsity Indic	es	Condition	
Gear	Species	Sample Size (n)	CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
AFS std gill net	Bigmouth Buffalo	203	12.5	3.8	16	6	11	6		
	Black Bullhead	13	2.0	1.0	8		8			
	Black Crappie	2	0.3	0.3	0		0		113	1
	Common Carp	60	8.2	1.7	6		4			
	Northern Pike	26	4.3	2.0	38	15	12		89	3
	Saugeye	28	4.7	1.1	29	13	14		89	1
	White Sucker	3	0.5	0.5	100		100			
	Yellow Perch	7	1.2	0.9	86		43		101	10

## 10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

\* Methods/Species that ignore stock length

							CPUE					
Gear	Species	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	Avg
AFS std gill net	Bigmouth Buffalo			0.0	5.5	1.2		1.8	2.2		12.5	3.87
	Black Bullhead			1.0	2.8	0.5		4.3	0.5		2.0	1.85
	Black Crappie			0.0	0.0	0.0		0.3	0.2		0.3	0.13
	Common Carp			0.2	2.3	1.0		0.5	0.3		8.2	2.08
	Northern Pike			2.0	0.8	1.5		0.8	1.0		4.3	1.73
	Saugeye			0.7	0.5	4.0		5.5	3.5		4.7	3.15
	Walleye			2.0	8.7	3.2		2.0	2.3		0.0	3.03
	White Sucker			1.7	1.5	0.7		1.5	0.5		0.5	1.07
	Yellow Perch			5.0	2.8	0.7		1.0	4.5		1.2	2.53
std exp gill net	Bigmouth Buffalo	0.0	0.0									0.00
	Black Bullhead	9.3	0.0									4.65
	Common Carp	0.3	0.3									0.30
	Northern Pike	1.3	2.3									1.80
	Orangespotted Sunfish	0.0	0.0									0.00
	Saugeye	0.0	0.0									0.00
	Walleye	0.0	6.3									3.15
	White Sucker	1.0	4.0									2.50
	Yellow Perch	7.0	40.3									23.65

## **10-Year Size Structure and Condition Statistics by Gear and Species**

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

							Ye	ar				
Gear	Species	Index	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
AFS std gill net	Bigmouth Buffalo	PSD				12	100		100	69		16
		PSD-P				3	0		36	46		11
	Black Bullhead	PSD			0	18	100		81	0		8
		PSD-P			0	0	33		0	0		8
	Black Crappie	PSD							100	100		0
		PSD-P							50	100		0
		Wr							98	94		113
	Common Carp	PSD			100	79	100		100	0		6
		PSD-P			100	7	50		67	0		4
	Northern Pike	PSD			92	80	56		60	67		38
		PSD-P			8	20	33		0	0		12
		Wr			88	100	88		81	75		89
	Saugeye	PSD			0	100	63		91	76		29
		PSD-P			0	0	25		18	52		14
		Wr			99	93	93		90	81		89
	Walleye	PSD			17	96	100		100	79		
		PSD-P			0	0	32		83	36		
		Wr			89	93	92		84	81		
	White Sucker	PSD			100	100	100		100	67		100
		PSD-P			80	78	75		100	67		100
	Yellow Perch	PSD			67	94	25		100	70		86
		PSD-P			50	12	25		33	26		43
		Wr			111	99	108		103	96		101
std exp gill net	Black Bullhead	PSD	4									
		PSD-P	0									
	Common Carp	PSD	0	100								
		PSD-P	0	100								
	Northern Pike	PSD	50	86								
		PSD-P	0	29								
		Wr	77	84								
	Saugeye	PSD		0								
		PSD-P		0								
	Walleye	PSD	0	0								

							Ye	ar				
Gear	Species	Index	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
std exp gill net	Walleye	PSD-P	0	0								
		Wr		96								
	White Sucker	PSD	100	83								
		PSD-P	100	58								
	Yellow Perch	PSD	43	75								
		PSD-P	33	8								
		Wr	105	107								

## Length at Capture

Mean length at capture by age across years sampled, sample size (N).

## Species: Saugeye

Mean Length (expanded sample number) at capture by age											
Year	Ν	1	2	3	4	5	6	7	8	9	10+
2024	28	310 (20)	386 (3)	447 (2)		556 (2)	565 (1)				
2022	34	226 (18)		468 (13)	533 (1)		613 (2)				
2021	32		369 (27)	493 (1)	544 (3)	488 (1)					
2019	24	273 (9)	401 (5)	464 (10)							
2018	3		423 (3)								

# Species: Walleye

				Mean Ler	ngth (expa	nded sam	ple numb	er) at cap	ture by ag	e	
Year	N	1	2	3	4	5	6	7	8	9	10+
2022	14		385 (7)					594 (2)	514 (4)	610 (1)	
2021	12						548 (3)	547 (9)			
2019	18				455 (2)	487 (15)		628 (1)			
2018	52	298 (1)	411 (12)	409 (5)	452 (34)						
Species: Y	ellow Pe	erch									
				Mean Ler	ngth (expa	nded sam	ple numb	er) at cap	ture by ag	e	
Year	Ν	1	2	3	4	5	6	7	8	9	10+
2019	4	144 (3)		278 (1)							


## Fish Condition


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

					Length	Group	S		
			S-Q	Q-P		P-M			М
Species	Year	N	Wr (SE)	Ν	Wr (SE)	Ν	Wr (SE)	Ν	Wr (SE)
Northern Pike Gill Net	2021	2	92 (3.8)	3	73 (3.9)	0		0	
	2022	2	67 (2.9)	4	79 (2.1)	0		0	
	2024	16	86 (1.7)	7	93 (5.8)	3	97 (0.4)	0	
Saugeye Gill Net	2021	3	89 (4.8)	24	90 (1.0)	4	88 (2.5)	2	90 (7.0)
	2022	5	82 (1.6)	5	76 (2.3)	9	82 (2.6)	2	89 (1.9)
	2024	20	88 (1.3)	4	88 (2.3)	2	93 (3.2)	2	93 (1.6)
Walleye Gill Net	2021	0		2	86 (1.8)	10	84 (1.4)	0	
	2022	3	80 (0.7)	6	80 (2.0)	5	84 (3.6)	0	
Yellow Perch Gill Net	2021	0		4	105 (3.7)	2	100 (0.8)	0	
	2022	8	104 (2.6)	12	94 (2.0)	7	90 (2.3)	0	
	2024	1	93	3	111 (18.4)	3	94 (3.2)	0	


#### **Length Frequency Distribution**

Length frequency histogram of species sampled by year.








Species: Common Carp Gear: AFS std gill net




Species: Northern Pike Gear: AFS std gill net









Species: Walleye Gear: AFS std gill net






Species: Yellow Perch Gear: AFS std gill net




#### **Historic Fish Sizes and Relative Abundance**


Size distribution per net by color for species sampled by year.

#### Species: Black Bullhead Gear: AFS std gill net





Species: Black Bullhead Gear: std exp gill net






Species: Common Carp Gear: std exp gill net







Species: Northern Pike Gear: std exp gill net




Species: Saugeye Gear: AFS std gill net




Species: Saugeye Gear: std exp gill net




Species: Walleye Gear: AFS std gill net




Species: Walleye Gear: std exp gill net




Species: White Sucker Gear: AFS std gill net




Species: White Sucker Gear: std exp gill net



Species: Yellow Perch Gear: AFS std gill net



Species: Yellow Perch Gear: std exp gill net



# Fish Stocking

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2014	Walleye	Fry	300,000
2015	Walleye	Small Fingerling	20,480
2016	Saugeye	Small Fingerling	31,030
2018	Saugeye	Small Fingerling	20,550
2019	Saugeye	Small Fingerling	21,120
2021	Saugeye	Fry	400,000
2021	Saugeye	Juvenile	25,830
2023	Saugeye	Juvenile	20,944