

# SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Minnewasta, Day County

UBS-Lake-411-705

2024

## Lake Information

**Name:** Minnewasta

**County:** Day

**Surface Area:** 606 Acres

## Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
AFS std gill net	Jul 09, 2024	4 net-nights
AFS std gill net	Jul 10, 2024	4 net-nights
AFS std gill net	Jul 11, 2024	4 net-nights

## **Common Fish Species Present**

Yellow Perch

Walleye

Northern Pike

White Sucker

Smallmouth Bass

Rock Bass

---

## Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- **AFS std gill net** – 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- **std experimental gill net for non-Missouri River waters** - 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- **std experimental gill net for Missouri River reservoirs** – 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$CPUE = \frac{\text{number of fish}}{\text{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left( \frac{\text{number of fish} \geq \text{quality length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

$$PSD - P = \left( \frac{\text{number of fish} \geq \text{preferred length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (**W**) compared to a length-specific standard weight (**Ws**) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left( \frac{W}{W_s} \right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (S), quality (Q), preferred (P), memorable (M) and trophy (T). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

## Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

\* **Methods/Species that ignore stock length**

Gear	Species	Sample Size (n)	Abundance		Stock Density Indices			Condition		
			CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
AFS std gill net	Northern Pike	25	2.1	0.5	96		20	13	80	2
	Rock Bass	1	0.1	0.1	100		0		106	
	Smallmouth Bass	3	0.3	0.2	100		67		105	6
	Walleye	37	2.8	0.8	85	10	38	13	86	1
	White Sucker	9	0.8	0.4	100		100		103	3
	Yellow Perch	190	15.8	2.1	69	5	14	4	117	7

## 10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

\* Methods/Species that ignore stock length

Gear	Species	CPUE										Avg	
		2015	2016	2017	2018	2019	2020	2021	2022	2023	2024		
AFS std gill net	Black Bullhead				0.1			0.3				0.0	0.13
	Black Crappie				0.1			0.0				0.0	0.03
	Common Carp				0.4			0.2				0.0	0.20
	Northern Pike				0.8			2.7				2.1	1.87
	Rock Bass				0.2			0.2				0.1	0.17
	Smallmouth Bass				0.0			0.2				0.3	0.17
	Walleye				2.8			4.6				2.8	3.40
	White Bass				0.1			0.3				0.0	0.13
	White Sucker				0.9			0.4				0.8	0.70
	Yellow Perch				3.1			2.8				15.8	7.23
std exp gill net	Black Bullhead	1.7											1.70
	Northern Pike	0.2											0.20
	Walleye	0.8											0.80
	White Sucker	1.2											1.20
	Yellow Perch	1.7											1.70

## 10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

Gear	Species	Index	Year									
			2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
AFS std gill net	Northern Pike	PSD				100			50			96
		PSD-P				30			22			20
		Wr				77			85			80
	Rock Bass	PSD				50			100			100
		PSD-P				0			0			0
		Wr				112			110			106
	Smallmouth Bass	PSD							100			100
		PSD-P							50			67
		Wr							111			105
	Walleye	PSD				38			49			85
		PSD-P				9			5			38
		Wr				90			88			86
	White Sucker	PSD				100			100			100
		PSD-P				100			100			100
		Wr				99			105			103
Yellow Perch	PSD				92			70			69	
	PSD-P				62			6			14	
	Wr				110			114			117	
std exp gill net	Northern Pike	PSD	0									
		PSD-P	0									
		Wr	72									
	Walleye	PSD	40									
		PSD-P	0									
		Wr	82									
	White Sucker	PSD	100									
		PSD-P	100									
		Wr	108									
Yellow Perch	PSD	100										
	PSD-P	80										
	Wr	103										

## **Length at Capture**

Mean length at capture by age across years sampled, sample size (N).

Species: Walleye

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2024	37		241 (4)	351 (5)	438 (5)	454 (4)	508 (2)		535 (7)	562 (2)	626 (8)
2021	55	200 (1)	295 (6)	365 (27)	405 (7)	435 (7)	490 (5)				697 (2)
2018	38	196 (4)	285 (11)	341 (7)	380 (11)		454 (1)		539 (4)		
2015	5				317 (3)	389 (2)					

Species: Yellow Perch

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2024	189	138 (4)	204 (145)	244 (29)	276 (8)	314 (1)	319 (2)				
2021	33		190 (13)	231 (20)							
2018	37	133 (2)	209 (6)	245 (10)		289 (2)	289 (4)	293 (8)	294 (3)	334 (1)	340 (1)
2015	10			215 (1)	283 (7)		317 (2)				



## **Fish Condition**



Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

Species	Year	Length Groups							
		S-Q		Q-P		P-M		M	
		N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Northern Pike Gill Net	2021	16	86 (1.8)	9	83 (1.9)	3	90 (3.6)	4	82 (4.0)
	2024	1	77	19	79 (1.5)	5	81 (1.7)	0	
Walleye Gill Net	2021	28	88 (1.0)	24	88 (0.9)	1	95	2	92 (2.8)
	2024	5	81 (1.5)	16	83 (0.8)	9	91 (1.7)	4	94 (1.1)
White Sucker Gill Net	2021	0		0		0		5	105 (6.0)
	2024	0		0		0		9	103 (2.1)
Yellow Perch Gill Net	2021	10	113 (1.7)	21	114 (1.7)	2	119 (7.0)	0	
	2024	58	113 (1.1)	105	111 (0.8)	23	155 (45.6)	3	103 (2.0)



# Length Frequency Distribution

Length frequency histogram of species sampled by year.



Species: Northern Pike  
Gear: AFS std gill net



Species: Walleye  
Gear: AFS std gill net




Species: Yellow Perch  
Gear: AFS std gill net




## Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.


Species: Northern Pike  
Gear: AFS std gill net




Species: Northern Pike  
Gear: std exp gill net




Species: Walleye  
Gear: AFS std gill net




Species: Walleye  
Gear: std exp gill net




Species: White Sucker  
Gear: AFS std gill net




Species: White Sucker  
Gear: std exp gill net



Species: Yellow Perch  
Gear: AFS std gill net



Species: Yellow Perch  
Gear: std exp gill net



## **Fish Stocking**

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2018	Yellow Perch	Fingerling	26,750
2019	Walleye	Fry	300,000
2021	Walleye	Fry	600,000
2023	Walleye	Fry	600,000