

SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Opitz, Day County

UJA-Lake-866-002

2024

Lake Information

Name: Opitz **Maximum Depth:** 23 Feet
County: Day **Mean Depth:** 14 Feet
Surface Area: 1,452 Acres

Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
fall night EF-WAE	Sep 19, 2024	3600 seconds

Common Fish Species Present

Yellow Perch

Walleye

Northern Pike

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- **AFS std gill net** – 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- **std experimental gill net for non-Missouri River waters** - 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- **std experimental gill net for Missouri River reservoirs** – 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$CPUE = \frac{\text{number of fish}}{\text{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{\text{number of fish} \geq \text{quality length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

$$PSD - P = \left(\frac{\text{number of fish} \geq \text{preferred length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (*W*) compared to a length-specific standard weight (*Ws*) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{W_s} \right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (S), quality (Q), preferred (P), memorable (M) and trophy (T). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

* Methods/Species that ignore stock length

Gear	Species	CPUE										Avg
		2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	
AFS std frame net	Black Bullhead			1.1								1.10
	Common Carp			0.3								0.30
	Northern Pike			0.1								0.10
	Orangespotted Sunfish			0.0								0.00
	Rock Bass			0.9								0.90
	Walleye			1.5								1.50
AFS std gill net	Black Bullhead		0.8	0.4	0.2	0.0		0.3	0.5	0.0		0.31
	Black Crappie		0.0	0.0	0.0	0.0		0.0	0.1	0.0		0.01
	Common Carp		0.1	0.3	0.2	0.3		0.5	0.3	0.1		0.26
	Northern Pike		0.4	0.1	0.1	0.0		0.2	0.0	0.3		0.16
	Rock Bass		1.8	2.8	2.2	2.1		3.4	3.2	0.9		2.34
	Smallmouth Bass		0.2	0.0	0.0	0.0		0.1	0.0	0.0		0.04
	Walleye		5.6	3.3	7.4	5.2		3.5	1.5	0.9		3.91
	White Bass		0.0	0.0	0.0	0.0		0.1	0.0	0.0		0.01
	White Sucker		0.0	0.0	0.2	0.3		0.1	0.5	0.3		0.20
	Yellow Perch		14.3	13.8	11.7	9.4		3.3	3.4	4.2		8.59
boat shocker (night)	Walleye*		360.0									360.00
fall night EF-WAE*	Walleye				327.0	3.0		459.0		105.0	49.0	188.60
std exp gill net	Black Bullhead	0.7										0.70
	Common Carp	0.5										0.50
	Northern Pike	0.5										0.50
	Rock Bass	0.8										0.80
	Walleye	22.5										22.50
	Yellow Perch	33.8										33.80

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

Gear	Species	Index	Year										
			2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	
AFS std frame net	Northern Pike	PSD			100								
		PSD-P			0								
		Wr			81								
	Walleye	PSD			85								
		PSD-P			8								
		Wr			69								
AFS std gill net	Northern Pike	PSD		100	100	100				100		100	
		PSD-P		80	0	0				0		50	
		Wr		93	81	70				90		93	
	Walleye	PSD		27	55	11	6			40	69	55	
		PSD-P		1	0	1	2			0	0	9	
		Wr		82	80	82	83			85	82	86	
	Yellow Perch	PSD		100	91	97	91			97	65	28	
		PSD-P		93	84	59	66			90	59	28	
		Wr		103	108	111	112			106	108	112	
boat shocker (night)	Walleye	PSD		0									
		PSD-P		0									
		Wr		90									
std exp gill net	Northern Pike	PSD	100										
		PSD-P	67										
		Wr	81										
	Walleye	PSD	5										
		PSD-P	0										
		Wr	80										
	Yellow Perch	PSD	96										
		PSD-P	74										
		Wr	108										

Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Walleye

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2023	14	218 (3)	303 (5)			404 (3)		430 (1)			593 (2)
2022	26	194 (10)			384 (8)		422 (6)			336 (1)	447 (1)
2021	42			365 (29)		408 (12)					403 (1)
2019	63	221 (2)		329 (54)		373 (2)	376 (3)	548 (1)	391 (1)		
2018	89		293 (63)		365 (5)	372 (7)	398 (2)	386 (10)			630 (1)
2017	66	205 (26)	317 (1)	355 (2)	385 (19)	384 (3)	386 (15)		405 (1)		
2016	67		320 (6)	349 (21)	374 (10)	380 (30)		520 (1)			
2015	143	201 (7)	276 (24)	324 (24)	356 (88)	395 (2)					

Species: Yellow Perch

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2023	51	144 (35)	186 (2)	254 (1)				318 (8)			333 (5)
2022	38	142 (14)	240 (3)		298 (5)		315 (6)	333 (1)	323 (2)	327 (5)	328 (2)
2021	39	149 (1)		258 (9)	264 (2)	295 (13)			325 (10)	320 (2)	344 (2)
2019	114	138 (11)		244 (44)	273 (9)		300 (21)	299 (9)	307 (22)		
2018	140		220 (56)	267 (12)		294 (34)	297 (20)	303 (18)	342 (1)		
2017	166	141 (15)	231 (10)		272 (38)	289 (11)	298 (90)		325 (2)		
2016	171			254 (37)	277 (23)	300 (90)	309 (10)	305 (11)			
2015	203		197 (14)	229 (28)	274 (149)	278 (11)	325 (1)				

Fish Condition





Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

Species	Year	Length Groups							
		S-Q		Q-P		P-M		M	
		N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Northern Pike Gill Net	2021	0		2	90 (2.5)	0		0	
	2023	0		2	97 (3.1)	2	88 (1.3)	0	
Walleye Gill Net	2021	25	85 (0.8)	17	86 (0.9)	0		0	
	2022	5	83 (1.4)	11	82 (1.5)	0		0	
	2023	5	86 (1.3)	5	89 (2.8)	0		1	72
Yellow Perch Gill Net	2021	1	117	3	119 (4.7)	13	110 (2.9)	22	102 (1.7)
	2022	13	111 (1.9)	2	114 (9.1)	4	113 (1.6)	18	104 (2.2)
	2023	36	115 (1.4)	0		2	113 (3.6)	12	100 (2.8)





Length Frequency Distribution

Length frequency histogram of species sampled by year.

Species: Walleye
Gear: AFS std gill net



Species: Yellow Perch
Gear: AFS std gill net




Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.


Species: Northern Pike

Gear: AFS std gill net




Species: Northern Pike


Gear: std exp gill net




Species: Walleye
Gear: AFS std gill net




Species: Walleye
Gear: std exp gill net



Species: Yellow Perch
Gear: AFS std gill net



Species: Yellow Perch
Gear: std exp gill net



Fish Stocking

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2016	Walleye	Fry	700,000
2018	Walleye	Fry	710,000
2021	Walleye	Fry	800,000
2023	Walleye	Juvenile	106,547