SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Eureka, McPherson County

WMC-Lake-1372-800

2024

Lake Information

Name:	Eureka	Maximum Depth:	15 Feet
County:	McPherson	Mean Depth:	7 Feet
Surface Area:	202 Acres		

Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
AFS std gill net	Jul 09, 2024	4 net-nights
frame net (std 3/4 in)	Jul 09, 2024	5 net-nights
frame net (std 3/4 in)	Jul 10, 2024	5 net-nights

Common Fish Species Present

Yellow Perch

Walleye

Northern Pike

Common Carp

Bluegill

Green Sunfish

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- **AFS std gill net** 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- std experimental gill net for non-Missouri River waters 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- std experimental gill net for Missouri River reservoirs 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$CPUE = \frac{number \, off ish}{effort}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{number \ of fish \ge quality \ length}{number \ of \ fish \ge stock \ length}\right) \ge 100$$

$$PSD - P = \left(\frac{number \ offish \ge preferred \ length}{number \ of \ fish \ge stock \ length}\right) \ge 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{Ws}\right) \ge 100$$

Confidence intervals (**CI**) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (**S**), quality (**Q**), preferred (**P**), memorable (**M**) and trophy (**T**). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

	St	ock	Qu	ality	Preferred		Mem	orable	Tro	ophy
Species Name	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80). * Methods/Species that ignore stock length

			Abuno	dance	St	Stock Density Indices				ndition
Gear	Species	Sample Size (n)	CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
AFS std gill net	Common Carp	12	3.0	0.7	100		75		79	3
	Northern Pike	6	1.5	0.8	83		17		81	2
	Walleye	9	1.3	0.4	60		40		75	2
	Yellow Perch	3	0.8	0.4	67		33		107	5
frame net (std 3/4	Bluegill	4	0.2	0.3	100		50		104	22
in)	Green Sunfish	2	0.2	0.2	0		0		115	19
	Northern Pike	3	0.3	0.3	33		33		82	2
	Walleye	16	1.3	0.6	77		46	23	81	1
	Yellow Perch	5	0.5	0.4	20		0		98	4

10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

* Methods/Species that ignore stock length

	CPUE											
Gear	Species	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	Avg
AFS std gill net	Black Bullhead				0.0			0.3			0.0	0.10
	Channel Catfish				0.0			0.3			0.0	0.10
	Common Carp				2.5			1.0			3.0	2.17
	Northern Pike				1.0			0.8			1.5	1.10
	Walleye				1.5			10.3			1.3	4.37
	Yellow Perch				3.7			0.8			0.8	1.77
frame net (std	Bluegill				6.5			2.0			0.2	2.90
3/4 in)	Channel Catfish				0.0			0.1			0.0	0.03
	Common Carp				0.0			0.1			0.0	0.03
	Green Sunfish				0.0			0.0			0.2	0.07
	Northern Pike				0.3			0.5			0.3	0.37
	Orangespotted Sunfish				0.0			0.0			0.0	0.00
	Walleye				0.6			0.8			1.3	0.90
	Yellow Perch				2.7			0.4			0.5	1.20

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

	Year Species Index 2015 2016 2017 2018 2019 2020 2021 2022 202												
Gear	Species	Index	2015 2	2016	2017	2018	2019	2020	2021	2022	2023	2024	
AFS std gill net	Common Carp	PSD				100			100			100	
		PSD-P				93			25			75	
		Wr				81			85			79	
	Northern Pike	PSD				100			33			83	
		PSD-P				0			0			17	
		Wr				72			78			81	
	Walleye	PSD				56			95			60	
		PSD-P				11			2			40	
		Wr				80			84			75	
	Yellow Perch	PSD				73			33			67	
		PSD-P				0			0			33	
		Wr				95			80			107	
rame net (std /4 in)	Bluegill	PSD				9			40			100	
3/4 in)		PSD-P				2			10			50	
	٧	Wr				132			109			104	
	Common Carp	PSD				0			100				
		PSD-P				0			100				
		Wr							76				
	Green Sunfish	PSD										0	
		PSD-P										0	
		Wr										115	
	Northern Pike	PSD				100			60			33	
		PSD-P				0			0			33	
		Wr				80			87			82	
	Walleye	PSD				67			100			77	
		PSD-P				33			25			46	
		Wr				82			83			81	
	Yellow Perch	PSD				0			75			20	
		PSD-P				0			75			0	
		Wr				104			80			98	

Back-Calculated Lengths

Mean species back-calculated total length (mm) at age, standard error (SE), and sample size (N).

Species: Walleye

			Mean back-calculated length (SE) at age											
Year Class	Age	Ν	1	2	3	4	5	6	7	8	9	10		
2023	1	2	165 (3.2)											
2022	2	2	133 (3)	182 (5)										
2022	2	2	151 (0)	210 (13.5)										
2021	3	1	169	268	310									
2021	3	1	185	247	315									
2020	4	2	155 (16.2)	155 (16.2)	197 (5.4)	197 (5.4)								
2019	5	1	203	231	300	377	435							
2019	5	2	202 (.7)	260 (7.8)	320 (.4)	366 (7.3)	424 (3.5)							
2018	6	2	136 (39.2)	157 (18.1)	196 (23.4)	220 (.5)	312 (30.7)	349 (67.4)						
2018	6	6	151 (22.2)	192 (25.8)	251 (30.4)	300 (29.7)	366 (21.7)	415 (21.2)						
2016	8	1	103	103	144	144	252	252	336	336				
2014	10	2	105 (5.2)	105 (5.2)	153 (5.9)	153 (5.9)	259 (2.4)	259 (2.4)	310 (5.7)	310 (5.7)	390 (2.9)	390 (2.9)		
Weighted Mean		24	153	188	239	262	348	360	319	319	390	390		
Year Class	Age	Ν	11	12	13	14	15	16	17	18	19	20		
2023	1	2												
2022	2	2												
2022	2	2												
2021	3	1												
2021	3	1												
2020	4	2												
2019	5	1												

2019	5	2				
2018	6	2				
2018	6	6				
2016	8	1				
2014	10	2				
Weighted Mean		24				

Species: Yellow Perch

		Mean back-calculated length (SE) at age													
Year Class	Age	Ν	1	2	3	4	5	6	7	8	9	10			
2022	2	1	78	118											
2022	2	2	102 (6.6)	128 (2.1)											
2021	3	2	74 (4.8)	95 (9.5)	121 (2.2)										
2020	4	1	104	140	164	199									
2019	5	1	85	102	129	160	186								
2019	5	1	121	180	197	215	240								
Weighted Mean		8	93	123	146	191	213								
Year Class	Age	Ν	11	12	13	14	15	16	17	18	19	20			
2022	2	1													
2022	2	2													
2021	3	2													
2020	4	1													
2019	5	1													
2019	5	1													
Weighted Mean		8													

Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Bluegill

				Mean Ler	ngth (expa	nded sam	ple numbe	er) at capt	ure by ag	е	
Year	Ν	1	2	3	4	5	6	7	8	9	10+
2021	20	89 (1)	110 (1)	133 (1)	138 (11)	173 (2)	175 (1)	199 (1)	231 (2)		
2018	54	108 (45)	122 (2)	148 (2)	152 (2)			192 (2)		207 (1)	
Species: W	alleye										
				Mean Ler	ngth (expa	nded sam	ple numbe	er) at capt	ure by ag	е	
Year	Ν	1	2	3	4	5	6	7	8	9	10+
2024	8	190 (2)	221 (2)	360 (1)		454 (1)	430 (2)				
2021	41				405 (20)		440 (19)		484 (1)		525 (1)
2018	13	210 (4)		306 (4)		439 (4)		615 (1)			
pecies: Y	ellow Pe	erch									
				Mean Ler	ngth (expa	nded sam	ple numbe	er) at capt	ure by ag	e	
Year	Ν	1	2	3	4	5	6	7	8	9	10+
2024	3		158 (1)		210 (1)	250 (1)					
2021	3				195 (3)						
2018	22	142 (1)	200 (9)	214 (1)	224 (2)	235 (2)	235 (3)	235 (1)	245 (1)	245 (1)	

Fish Condition


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

					Length	Group	S		
			S-Q		Q-P		P-M		М
Species	Year	N	Wr (SE)	Ν	Wr (SE)	Ν	Wr (SE)	Ν	Wr (SE)
Bluegill Frame Net	2021	12	104 (1.6)	6	113 (3.8)	2	124 (2.0)	0	
	2024	0		1	87	1	121	0	
Common Carp Gill Net	2021	0		3	88 (5.0)	1	75	0	
	2024	0		3	87 (2.0)	8	76 (2.9)	1	78
Northern Pike Gill Net	2021	2	83 (1.0)	1	69	0		0	
	2024	1	79	4	80 (1.7)	0		1	87
Walleye Gill Net	2021	2	88 (2.7)	38	83 (0.7)	1	86	0	
	2024	2	80 (1.6)	1	72	2	72 (1.1)	0	
Yellow Perch Gill Net	2021	2	82 (2.9)	1	77	0		0	
	2024	1	112	1	98	1	111	0	


Length Frequency Distribution

Length frequency histogram of species sampled by year.


Species: Bluegill Gear: frame net (std 3/4 in)



Species: Common Carp Gear: AFS std gill net




Species: Walleye Gear: AFS std gill net




Historic Fish Sizes and Relative Abundance


Size distribution per net by color for species sampled by year.


Species: Bluegill Gear: frame net (std 3/4 in)




Species: Common Carp Gear: AFS std gill net








Species: Walleye Gear: AFS std gill net





Fish Stocking

Number of fish stocked by year, species, and size.

2013WalleyeFry2015WalleyeFry2017WalleyeFry2019WalleyeFry2021WalleyeFry2023WalleyeFry	Number
2017WalleyeFry2019WalleyeFry2021WalleyeFry	100,000
2019Walleye2021WalleyeFry	95,000
2021 Walleye Fry	100,000
	100,000
2023 Walleye Fry	100,000
	100,000
2024 Bluegill Adult	350
2024 Largemouth Bass Juvenile	935