Lake Mitchell Survey Summary

Lake Mitchell, located in Mitchell, SD, is managed as a bluegill, largemouth bass, and walleye fishery; other fish species (e.g., black crappie, channel catfish, flathead catfish, freshwater drum, smallmouth bass, and white crappie) provide additional angling opportunities.

- Bluegill. Bluegill abundance increased to 21.4 fish per frame net in 2024 resulting in one of the highest catch rates in the region. Relative abundance was similar to the previous year but well above the long term mean (CPUE = 20.5 and 12.1 fish per net, respectively). Netted fish ranged from 3.1 to 8.5 inches in length with approximately 75% measuring >6 inches. The sample was comprised of six year classes of fish ranging from 1 to 6 years in age. The age 2 cohort accounted for a majority of catches (53%) followed by age 3 fish (32%). Growth was excellent with fish averaging 6.1 inches in length by age 2 and 7.1 inches by age 3. A high mean relative weight score (Wr = 107) indicates the bluegill population is in great condition. Any angler targeting bluegill in the southeast region should be sure to consider trying Lake Mitchell.
- Black Crappie. Black crappie abundance improved considerably from the previous sample year (CPUE = 1.3 and 7.1 fish per frame net in 2023 and 2024, respectively). Netted fish ranged from 3.9 to 10.6 inches in length with approximately 22% measuring in the quality length category (>8 inches). Aging data from previous years indicates black crappie typically grow to a mean length of 6 inches by age 2 and 8 inches by age 3 in Lake Mitchell. A high mean relative weight score (Wr = 96 in 2024) indicates the black crappie population is in good condition.
- **Channel Catfish.** The 2024 net catches of channel catfish (15.4/net) were the highest since 2015 (18.7/net). Size of Lake Mitchell channel catfish is excellent with over 90% of the stock length (11 inch and longer) channel catfish caught in nets at least measuring 16 inches and nearly 20% measuring over 24 inches long. Lake Mitchell provides one of the best channel catfish opportunities in the state.
- Walleye. Sampling efforts produced a catch rate of 0.5 walleye per gill net in 2024. Relative abundance was similar to the previous year and long term mean (CPUE = 0.6 and 0.4 fish per net, respectively). Sampled fish ranged from 15.3 to 20.6 inches in length with 20% measuring in the preferred length category (>20 inches). Fisheries staff began stocking saugeye this year (400,00 fry and 56,000 juveniles) in an attempt to improve catch rates. These walleye/sauger hybrids have been found to be more productive in lakes with marginal water conditions.

For more detailed results see the computer-generated South Dakota Statewide Fisheries Survey For Lake Mitchell (below).

SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Mitchell, Davison County

LJA-Lake-623-000

2024

Lake Information

Name:	Mitchell	Maximum Depth:	29 Feet
County:	Davison	Mean Depth:	12 Feet
Legal Description:	T103W- R60N-Sec 4-6, 9; T104N- R60W-Sec 31-32		
Surface Area:	690 Acres	Watershed Area:	19,821.31 Sq Miles

Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort	
AFS std gill net	Jun 12, 2024	5 net-nights	
AFS std gill net	Jun 13, 2024	5 net-nights	
frame net (std 3/4 in)	Jun 12, 2024	4 net-nights	
frame net (std 3/4 in)	Jun 13, 2024	5 net-nights	

Common Fish Species Present

Largemouth Bass Bluegill Black Crappie Channel Catfish Freshwater Drum White Crappie River Carpsucker Common Carp Smallmouth Bass Walleye

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- **AFS std gill net** 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- std experimental gill net for non-Missouri River waters 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- std experimental gill net for Missouri River reservoirs 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$\textit{CPUE} = \frac{\textit{number of fish}}{\textit{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{number \ of fish \ge quality \ length}{number \ of \ fish \ge stock \ length}\right) \ge 100$$

$$PSD - P = \left(\frac{number \ offish \ge preferred \ length}{number \ of \ fish \ge stock \ length}\right) \ge 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{Ws}\right) \ge 100$$

Confidence intervals (**CI**) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (**S**), quality (**Q**), preferred (**P**), memorable (**M**) and trophy (**T**). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

	St	ock	Qu	ality	Pref	erred	Mem	orable	Tro	ophy
Species Name	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80). * Methods/Species that ignore stock length

			Abun	dance	St	ock Der	es	Cor	ndition	
Gear	Species	Sample Size (n)	CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
AFS std gill net	Bigmouth Buffalo	3	0.1	0.1	0		0			
	Black Crappie	1	0.1	0.1	0		0		116	
	Bluegill	1	0.1	0.1	100		0		104	
	Channel Catfish	18	1.5	1.0	80		7		97	4
	Common Carp	13	0.7	0.5	0		0			
	Freshwater Drum	24	2.0	0.8	85		10			
	River Carpsucker	13	1.3	1.2	100		100			
	Shorthead Redhorse	3	0.3	0.2	100		100			
	Walleye	5	0.5	0.4	100		20		93	2
frame net (std 3/4	Black Crappie	66	7.1	5.1	22	8	5		96	2
in)	Bluegill	193	21.4	12.1	75	4	2		107	2
	Channel Catfish	139	15.4	15.6	94	3	19	5	98	2
	Common Carp	6	0.7	0.5	33		33			
	Freshwater Drum	4	0.0	0.0	0		0			
	Green Sunfish	1	0.1	0.2	0		0			
	Northern Pike	1	0.1	0.2	100		100		102	
	Shorthead Redhorse	1	0.1	0.2	100		100			
	Smallmouth Bass	6	0.7	0.8	17		0		101	3
	White Crappie	18	2.0	1.1	100		89		90	2

10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

* Methods/Species that ignore stock length

							CPUE					
Gear	Species	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	Avg
AFS std frame	Black Crappie			20.3								20.30
net	Bluegill			4.6								4.60
	Channel Catfish			2.3								2.30
	Common Carp			1.3								1.30
	Freshwater Drum			0.1								0.10
	Green Sunfish			0.1								0.10
	Northern Pike			0.1								0.10
	Orangespotted Sunfish			0.0								0.00
	Shorthead Redhorse			0.1								0.10
	Sunfish Hybrid			0.0								0.00
	White Crappie			16.9								16.90
AFS std gill net	Bigmouth Buffalo			0.1	0.0	0.1		0.4	0.1	0.1	0.1	0.13
	Black Bullhead			0.0	0.0	0.4		0.2	0.0	0.0	0.0	0.09
	Black Crappie			7.4	1.5	0.8		0.5	0.4	0.7	0.1	1.63
	Bluegill			0.1	0.2	0.1		0.2	0.1	0.7	0.1	0.21
	Channel Catfish			13.4	8.3	8.9		2.3	4.5	4.5	1.5	6.20
	Common Carp			0.0	0.7	0.7		1.1	0.2	0.5	0.7	0.56
	Flathead Catfish			0.1	0.0	0.1		0.0	0.0	0.0	0.0	0.03
	Freshwater Drum			2.6	8.6	2.8		2.9	1.4	2.6	2.0	3.27
	Northern Pike			0.3	0.0	0.0		1.5	1.7	0.4	0.0	0.56
	Quillback			0.0	0.0	0.0		0.0	0.0	0.0	0.0	0.00
	River Carpsucker			0.0	0.0	0.6		3.4	6.7	2.4	1.3	2.06
	Shorthead Redhorse			0.0	0.0	0.0		0.1	0.6	0.2	0.3	0.17
	Walleye			0.3	0.3	0.4		0.2	0.2	0.6	0.5	0.36
	White Crappie			3.0	0.8	1.5		0.7	0.0	0.3	0.0	0.90
	White Sucker			0.0	0.0	0.0		0.7	0.0	0.3	0.0	0.14
boat shocker	Largemouth Bass		3.5					4.5	3.5			3.83
(night)	Smallmouth Bass		20.0					0.0	0.0			6.67
electrofishing (flathead)	Flathead Catfish							9.3				9.30
fall night EF-	Flathead Catfish	13.6										13.60
WAE*	Walleye	0.2										0.20
frame net (std	Bigmouth Buffalo	0.5	0.0		0.0	0.0		0.0	0.0	0.0	0.0	0.06
3/4 in)	Black Bullhead	0.0	0.1		0.0	0.0		0.0	0.0	0.0	0.0	0.01

							CPUE					
Gear	Species	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	Avg
frame net (std	Black Crappie	7.3	43.9		12.7	2.1		14.3	3.5	1.3	7.1	11.53
3/4 in)	Bluegill	4.0	2.3		16.1	3.3		26.7	2.5	20.5	21.4	12.10
	Channel Catfish	18.7	6.4		9.2	1.8		8.8	1.3	0.6	15.4	7.78
	Common Carp	1.4	1.3		0.8	0.2		0.9	0.2	0.3	0.7	0.73
	Flathead Catfish	0.3	0.0		0.5	0.2		0.2	0.0	0.0	0.0	0.15
	Freshwater Drum	0.0	0.1		0.2	0.1		0.2	0.0	0.2	0.0	0.10
	Green Sunfish	0.0	0.4		0.1	0.0		0.2	0.0	0.0	0.1	0.10
	Largemouth Bass	0.0	0.1		0.0	0.0		0.0	0.0	0.0	0.0	0.01
	Northern Pike	0.1	0.3		0.0	0.0		0.0	0.0	0.2	0.1	0.09
	Orangespotted Sunfish	0.0	0.0		0.0	0.0		0.0	0.0	0.0	0.0	0.00
	Shorthead Redhorse	0.0	0.1		0.0	0.0		0.0	0.0	0.0	0.1	0.03
	Smallmouth Bass	0.0	0.2		0.4	0.1		0.1	0.6	0.0	0.7	0.26
	Sunfish Hybrid	0.0	0.0		0.0	0.1		0.0	0.0	0.1	0.0	0.03
	Walleye	0.0	0.0		0.0	0.0		0.1	0.0	0.1	0.0	0.03
	White Crappie	0.7	15.3		21.5	4.2		13.4	3.1	0.0	2.0	7.53
	White Sucker	0.0	0.0		0.1	0.0		0.0	0.0	0.0	0.0	0.01
hoop net	Black Crappie		7.0	0.0								3.50
	Bluegill		1.3	0.0								0.65
	Channel Catfish		0.7	2.3								1.50
	Smallmouth Bass		0.7	0.0								0.35
	White Crappie		2.3	0.0								1.15
std exp gill net	Bigmouth Buffalo	0.0	1.0									0.50
	Black Crappie	14.0	5.8									9.90
	Bluegill	0.0	0.2									0.10
	Channel Catfish	18.4	22.0									20.20
	Common Carp	0.4	0.8									0.60
	Freshwater Drum	3.8	5.0									4.40
	Largemouth Bass	0.0	0.0									0.00
	Northern Pike	0.0	0.0									0.00
	Shorthead Redhorse	0.0	0.0									0.00
	Smallmouth Bass	0.2	0.0									0.10
	Walleye	0.8	0.8									0.80
	White Crappie	0.0	5.0									2.50
	White Sucker	0.0	0.0									0.00

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

							Ye	ar				
Gear	Species	Index	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
AFS std frame	Black Crappie	PSD			100							
net		PSD-P			0							
		Wr			95							
	Bluegill	PSD			85							
		PSD-P			4							
		Wr			114							
	Channel Catfish	PSD			79							
		PSD-P			4							
		Wr			95							
	Common Carp	PSD			100							
		PSD-P			40							
	White Crappie	PSD			99							
		PSD-P			0							
		Wr			100							
AFS std gill net	Black Crappie	PSD			100	80	25		40	50	29	0
		PSD-P			2	7	13		0	0	29	0
		Wr			102	94	105		111	111	120	116
	Bluegill	PSD			100	100	0		100	100	43	100
		PSD-P			0	0	0		50	0	29	0
		Wr			100	121	132		118	121	113	104
	Channel Catfish	PSD			85	87	82		70	87	80	80
		PSD-P			2	4	3		9	4	4	7
		Wr			86	87	96		93	94	91	97
	Common Carp	PSD				86	100		64	100	100	0
		PSD-P				43	71		18	100	80	0
	River Carpsucker	PSD					83		100	100	100	100
		PSD-P					83		53	97	100	100
	Walleye	PSD			100	100	75		50	50	33	100
		PSD-P			100	33	75		0	0	0	20
		Wr			87	88	95		86	72	92	93
	White Crappie	PSD			96	50	60		43		100	
		PSD-P			0	13	0		0		100	

							Ye	ar				
Gear	Species	Index	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
AFS std gill net	White Crappie	Wr			105	96	105		106		102	
boat shocker	Largemouth Bass	PSD		71					100	86		
(night)		PSD-P		71					100	43		
		Wr		105					104	102		
	Smallmouth Bass	PSD		3								
		PSD-P		3								
		Wr		97								
frame net (std	Black Crappie	PSD	0	49		79	43		43	74	31	22
3/4 in)		PSD-P	0	2		0	5		2	3	23	5
		Wr	108	103		93	94		96	97	109	96
	Bluegill	PSD	29	79		84	91		63	44	40	75
		PSD-P	4	0		6	0		3	0	1	2
		Wr	104	107		105	105		107	113	115	107
	Channel Catfish	PSD	58	62		90	94		78	100	100	94
		PSD-P	13	1		7	6		8	31	33	19
		Wr	89	97		87	93		93	93	87	98
	Common Carp	PSD	100	93		100	50		88	100	100	33
		PSD-P	71	67		50	50		25	100	100	33
	Largemouth Bass	PSD		100								
		PSD-P		0								
		Wr		97								
	Smallmouth Bass	PSD		100		100	100		0	17	0	17
		PSD-P		50		50	100		0	0	0	0
		Wr		91		95	77		106	95		101
	Walleye	PSD							0		100	
		PSD-P							0		0	
		Wr							78		90	
	White Crappie	PSD	13	66		96	90		55	97		100
		PSD-P	0	0		0	2		12	10		89
		Wr	99	99		88	93		100	98		90
hoop net	Black Crappie	PSD		71								
		PSD-P		0								
		Wr		104								
	Bluegill	PSD		50								
		PSD-P		0								
		Wr		103								

							Ye	ar				
Gear	Species	Index	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
hoop net	Channel Catfish	PSD		50	64							
		PSD-P		0	0							
		Wr		87	93							
	Smallmouth Bass	PSD		0								
		PSD-P		0								
		Wr		100								
	White Crappie	PSD		14								
		PSD-P		0								
		Wr		101								
std exp gill net	Black Crappie	PSD	3	62								
		PSD-P	0	0								
		Wr	113	109								
	Bluegill	PSD		100								
		PSD-P		0								
		Wr		102								
	Channel Catfish	PSD	55	44								
		PSD-P	11	0								
		Wr	95	92								
	Common Carp	PSD	100	100								
		PSD-P	100	50								
	Smallmouth Bass	PSD	100									
		PSD-P	100									
		Wr	95									
	Walleye	PSD	75	0								
		PSD-P	0	0								
		Wr	76	93								
	White Crappie	PSD		76								
		PSD-P		0								
		Wr		105								

Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Black Crappie

Mean Length (expanded sample number) at capture by age												
Year	Ν	1	2	3	4	5	6	7	8	9	10+	
2022	35		164 (2)	206 (19)	236 (10)	232 (3)						
2021	117		169 (26)	199 (74)	217 (17)							
2015	87		180 (87)									

Species: Bluegill

				Mean Len	gth (expa	nded sam	ple numbe	er) at capt	ure by age	Э	
Year	N	1	2	3	4	5	6	7	8	9	10+
2024	180	119 (6)	154 (95)	180 (57)	191 (20)	215 (1)	215 (1)				
2023	196	93 (23)	138 (131)	178 (42)							
2022	25		133 (15)	170 (6)	191 (4)						
2021	223		125 (87)	177 (111)	191 (19)	195 (5)	220 (2)				
2015	48		135 (24)	151 (17)	175 (4)	202 (3)					

Species: Walleye

			Mean Ler	ngth (expa	inded sam	ole numbe	er) at cap	ture by age)	
Year	Ν	1 2	3	4	5	6	7	8	9	10+
2022	2		388 (2)							
2018	3				448 (1)			559 (2)		
2017	2									568 (2)
2016	4	281 (2)	290 (2)							
2015	4	262 (1)	387 (1)	438 (2)						
Species: W	hite Crap	pie								

	Mean Length (expanded sample number) at capture by							ure by age	ge		
Year	Ν	1	2	3	4	5	6	7	8	9	10+
2022	31			214 (24)	242 (7)						
2021	104		188 (66)	215 (28)		227 (11)					
2015	8		186 (8)								

Fish Condition


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).


		Length Groups							
			S-Q		Q-P		P-M		М
Species	Year	N	Wr (SE)	Ν	Wr (SE)	Ν	Wr (SE)	Ν	Wr (SE)
Black Crappie Frame Net	2021	73	100 (1.2)	54	91 (1.0)	2		0	
	2022	9	103 (2.3)	25	95 (1.5)	0		1	90
	2023	9	119 (2.8)	1	95	3	84 (5.0)	0	
	2024	50	98 (1.5)	11	89 (1.4)	3	85 (3.0)	0	
Bluegill Frame Net	2021	88	101 (1.8)	145	113 (1.1)	7	107 (5.4)	0	
	2022	14	110 (2.9)	11	116 (2.0)	0		0	
	2023	122	118 (1.7)	81	109 (2.2)	2		0	
	2024	48	106 (3.1)	141	108 (1.3)	4	96 (4.2)	0	
Channel Catfish Gill Net	2021	7	89 (4.7)	14	94 (3.2)	2	102 (4.2)	0	
	2022	6	88 (4.8)	37	95 (3.4)	2	98 (4.2)	0	
	2023	9	94 (2.3)	34	91 (2.3)	2	86 (1.2)	0	
	2024	3	85 (0.8)	11	98 (3.1)	1	116	0	
Largemouth Bass Electro Fishing	2021	0		0		9	104 (1.8)	0	
	2022	1	110	3	104 (4.7)	3	98 (3.4)	0	
Walleye	2021	1	82	1	89	0		0	
Gill Net	2022	1	90	1	54	0		0	
	2023	4	93 (4.7)	2	90 (0.4)	0		0	
	2024	0		4	92 (2.0)	1	97	0	
White Crappie Frame Net	2021	55	104 (0.9)	51	93 (2.2)	8		7	
	2022	1	101	27	98 (1.0)	3	98 (2.9)	0	
	2024	0		2	95 (11.6)	8	88 (1.3)	8	91 (3.0)


Length Frequency Distribution


Length frequency histogram of species sampled by year.

Species: Black Crappie Gear: frame net (std 3/4 in)











Species: White Crappie Gear: frame net (std 3/4 in)







Historic Fish Sizes and Relative Abundance


Size distribution per net by color for species sampled by year.

Species: Black Crappie Gear: AFS std frame net





Species: Black Crappie Gear: frame net (std 3/4 in)







Species: Bluegill Gear: frame net (std 3/4 in)







Species: Channel Catfish Gear: std exp gill net









Species: Common Carp Gear: std exp gill net







Species: Smallmouth Bass Gear: boat shocker (night)




Species: Walleye Gear: AFS std gill net




Species: Walleye Gear: std exp gill net





Species: White Crappie Gear: frame net (std 3/4 in)



Fish Stocking

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2013	Walleye	Small Fingerling	32,080
2015	Walleye	Small Fingerling	52,698
2016	Walleye	Small Fingerling	48,020
2017	Walleye	Fingerling	59,000
2018	Walleye	Small	47,040
2019	Walleye	Small Fingerling	51,000
2021	Walleye	Juvenile	59,570
2022	Largemouth Bass	Juvenile	40,000
2024	Saugeye	Fry	400,000
2024	Saugeye	Juvenile	56,463