Stockade Reservoir

Stockade is a 129-acre reservoir located 4.5 miles east of the town of Custer. It is located within Custer State Park so a South Dakota state parks sticker is needed. The primary fish species are Walleye, Yellow Perch, Smallmouth Bass, Largemouth Bass, Northen Pike, Bluegill and Black Crappie. Stockade is also home to some secondary species including Black Bullhead, White Sucker and Green Sunfish.

Walleye. Stockade's first stocking of Walleye was in 2017. In 2024 surveys had 2 fish per 60 minutes of boat electrofishing. The fish were in the 15–20-inch range.

Yellow Perch. Stockades Yellow Perch population suffers from a parasite commonly called white grub. The fish surveyed in 2024 showed that most of the perch were infected with the parasite. The surveys had 1.9 perch per gillnet and 8.4 fish per trap net. The perch were in the 5.5-8-inch range.

Smallmouth Bass. Stockade is home to some Smallmouth Bass. 4 were sampled via boat electrofishing in 2024, most of the fish were in the 7–11-inch range.

Largemouth Bass. There were 0.4 Largemouth Bass per gillnet and 71 fish were sampled via boat electrofishing in 2024. The oldest age class in stockade will be in the 12–15-inch range.

Northen Pike. The 2024 survey results had 1.8 Northern Pike per gillnet and 1.4 fish per trap net. Most of the pike were in the 21–28-inch range with several fish over 28 inches.

Bluegill. The surveys for 2024 had 11.9 Bluegill per gillnet and 24 fish per trap net. Most of the Bluegill will be in the 6–8-inch range, with several fish over the 8-inch mark.

Black Crappie. Stockade is home to a population of Black Crappie. The 2024 survey results are 3.7 fish per trap net and 1.8 fish per gill net. Most of the fish were over the 10-inch mark.

SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Stockade, Custer County MCS-Lake-12-000 2024

Lake Information

Name: Stockade

County: Custer

Surface Area: 129 Acres

Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
AFS std gill net	Aug 06, 2024	4 net-nights
AFS std gill net	Aug 07, 2024	4 net-nights
boat shocker (day)	Oct 11, 2024	3600 seconds
frame net (std 3/4 in)	Aug 06, 2024	4 net-nights
frame net (std 3/4 in)	Aug 07, 2024	3 net-nights

Common Fish Species Present

Yellow Perch

Smallmouth Bass

Northern Pike

Largemouth Bass

Bluegill

Black Bullhead

Black Crappie

Walleye

Green Sunfish

White Sucker

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

A statewide effort to help make netting efforts comparable to all waters sampled across the state, occurred in 2017, with a switch to American Fisheries Society gill nets. Past gill netting efforts were completed with different style/types of nets and are not comparable side by side.

- AFS std gill net 80 ft experimental gill net containing eight panels (10 ft each) of varying monofilament meshes of 0.75, 1.00, 1.25, 1.50, 1.75, 2.00, 2.25 and 2.50 inches.
- std experimental gill net for non-Missouri River waters 150 ft experimental gill net containing six panels (25 ft each) of varying monofilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.
- std experimental gill net for Missouri River reservoirs 300 ft experimental gill net containing six panels (50 ft each) of varying multifilament meshes of 0.5, 0.75, 1.00, 1.25, 1.50 and 2.00 inches.

$$\mathit{CPUE} = \frac{\mathit{number of fish}}{\mathit{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{number\ of\ fish \ge quality\ length}{number\ of\ fish \ge stock\ length}\right) \times 100$$

$$\textit{PSD} - \textit{P} = \left(\frac{number\ of\ fish\ \geq preferred\ length}{number\ of\ fish\ \geq stock\ length}\right) \ge 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{Ws}\right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (S), quality (Q), preferred (P), memorable (M) and trophy (T). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

	St	ock	Qu	ality	Pref	erred	Mem	orable	Tro	pphy
Species Name	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38
Bluegill	3	8	6	15	8	20	10	25	12	30
Brown Trout	8	20	12	30	16	40	20	50	18	46
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Lake Trout	12	30	20	50	26	65	31	80	39	100
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Rainbow Trout	10	25	16	40	20	50	26	65	31	80
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

* Methods/Species that ignore stock length

			Abun	dance	St	ock Der	nsity Indic	es	Cor	ndition
Gear	Species	Sample Size (n)	CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
AFS std gill net	Black Bullhead	50	6.3	2.7	100		14	8	94	2
	Black Crappie	14	1.8	1.2	71		0		116	17
	Bluegill	95	11.9	6.5	80	6	22	6	102	2
	Golden Shiner	3	0.0	0.0						
	Largemouth Bass	3	0.4	0.4	67		0		130	24
	Northern Pike	14	1.8	0.6	93		29		90	3
	Smallmouth Bass	2	0.3	0.4	100		100		87	8
	White Sucker	4	0.5	0.3	100		100		98	5
	Yellow Perch	15	1.9	1.3	47	21	0		85	2
boat shocker (day)	Largemouth Bass	98	71.0	11.1	56	9	20	7	109	2
	Smallmouth Bass	4	4.0	3.0	0		0		94	2
	Walleye*	2	2.0	3.0	100		100		101	14
frame net (std 3/4	Black Bullhead	36	5.1	5.3	100		8		88	4
in)	Black Crappie	26	3.7	2.9	65	15	12		94	3
	Bluegill	168	24.0	8.4	61	5	11	3	99	4
	Green Sunfish	4	0.6	0.3	100		0		111	5
	Northern Pike	10	1.4	0.9	40		0		87	3
	Walleye	1	0.1	0.2	100		100		84	
	Yellow Perch	62	8.4	3.4	25	8	0		87	5

10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

^{*} Methods/Species that ignore stock length

							CPUE					
Gear	Species	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	Avg
AFS std frame	Black Bullhead			0.0			18.1					9.05
net	Black Crappie			0.0			5.9					2.95
	Bluegill			0.0			8.4					4.20
	Green Sunfish			0.0			0.3					0.15
	Northern Pike			0.0			0.4					0.20
	White Sucker			0.0			0.1					0.05
	Yellow Perch			0.0			0.8					0.40
AFS std gill net	Black Bullhead				38.5		11.3	16.5	15.3	11.0	6.3	16.48
	Black Crappie				0.0		2.3	7.3	5.5	2.5	1.8	3.23
	Bluegill				1.5		5.0	20.5	11.5	6.5	11.9	9.48
	Channel Catfish				0.3		0.0	0.3	0.0	0.0	0.0	0.10
	Golden Shiner				0.0		0.0	0.0	0.0	0.0	0.0	0.00
	Largemouth Bass				0.0		0.0	0.3	0.0	0.3	0.4	0.17
	Northern Pike				1.8		2.3	3.3	2.0	1.8	1.8	2.17
	Smallmouth Bass				0.3		0.3	1.5	1.0	0.5	0.3	0.65
	Walleye				0.0		0.0	0.8	0.3	0.5	0.0	0.27
	White Sucker				0.3		8.0	1.8	0.3	0.5	0.5	0.70
	Yellow Perch				2.5		0.3	7.0	16.5	3.3	1.9	5.25
boat shocker	Largemouth Bass								86.0		71.0	78.50
(day)	Smallmouth Bass								2.0		4.0	3.00
	Walleye*								0.0		2.0	1.00
boat shocker	Largemouth Bass	84.0	68.0	66.9	105.0	134.0	59.0	64.0		47.0		78.49
(night)	Smallmouth Bass	0.0	9.0	13.7	7.0	26.0	1.0	4.0		6.0		8.34
	Walleye*	0.0	0.0	0.0	0.0	4.0	0.0	1.0		1.0		0.75
frame net (std 3/4 in)	Black Bullhead	471.1			91.4					6.9	5.1	143.6 3
	Black Crappie	5.3			2.5					3.1	3.7	3.65
	Bluegill	12.8			9.1					30.1	24.0	19.00
	Bluegill X Gr. Sunfish Hybrid	0.1			0.0					0.0	0.0	0.03
	Golden Shiner	0.0			0.0					0.0	0.0	0.00
	Green Sunfish	0.0			0.0					0.4	0.6	0.25
	Largemouth Bass	0.1			0.0					0.0	0.0	0.03
	Northern Pike	0.6			0.8					0.6	1.4	0.85
	Smallmouth Bass	0.4			0.6					0.0	0.0	0.25

2/26/2025 Page 6

							CPUE					
Gear	Species	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	Avg
frame net (std	Walleye	0.0			0.0					0.0	0.1	0.03
3/4 in)	White Sucker	0.1			0.1					0.0	0.0	0.05
	Yellow Perch	3.0			21.0					13.6	8.4	11.50
std exp gill net	Black Bullhead	41.0	53.5									47.25
	Black Crappie	0.5	0.0									0.25
	Bluegill	1.5	1.0									1.25
	Golden Shiner	0.0	0.0									0.00
	Largemouth Bass	0.5	1.0									0.75
	Northern Pike	5.5	4.5									5.00
	Smallmouth Bass	0.0	0.0									0.00
	White Sucker	0.5	1.0									0.75
	Yellow Perch	61.5	63.5									62.50

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

							Ye	ar				
Gear	Species	Index	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
AFS std frame	Black Bullhead	PSD			100		,	95		,		
net		PSD-P			0			0				
		Wr			97			86				
	Black Crappie	PSD			67			40				
		PSD-P			0			2				
		Wr			93			97				
	Bluegill	PSD						52				
		PSD-P						10				
		Wr						98				
	Green Sunfish	PSD						100				
		PSD-P						50				
		Wr						118				
	Northern Pike	PSD						100				
		PSD-P						0				
		Wr						85				
	White Sucker	PSD			100			100				
		PSD-P			100			100				
		Wr			100			81				
	Yellow Perch	PSD						33				
		PSD-P						0				
		Wr						81				
AFS std gill net	Black Bullhead	PSD				100		98	100	100	100	100
		PSD-P				0		0	2	7	16	14
		Wr				88		95	95		98	94
	Black Crappie	PSD						11	93	77	50	71
		PSD-P						0	0	0	10	0
		Wr						107	110		98	116
	Bluegill	PSD				100		75	94	98	81	80
		PSD-P				33		10	10	33	31	22
		Wr				109		108	106		97	102
	Largemouth Bass	PSD							100		100	67
		PSD-P							100		100	0
		Wr							116		100	130
							0/00	/2025		Pane 8		

2/26/2025 Page 8

							Ye	ar				
Gear	Species	Index	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
AFS std gill net	Northern Pike	PSD				86		100	85	88	100	93
		PSD-P				29		11	38	13	29	29
		Wr				95		91	96		90	90
	Smallmouth Bass	PSD				100		100	100	100	100	100
		PSD-P				100		100	67	100	100	100
		Wr				77		89	102		91	87
	Walleye	PSD							67	100	100	
		PSD-P							0	100	0	
		Wr							88		89	
	White Sucker	PSD				100		100	100	100	100	100
		PSD-P				100		100	100	100	100	100
		Wr				97		97	102		100	98
	Yellow Perch	PSD				60		0	0	5	15	47
		PSD-P				10		0	0	0	0	0
		Wr				84		93	108		81	85
boat shocker	Largemouth Bass	PSD								59		56
(day)		PSD-P								8		20
		Wr								109		109
	Smallmouth Bass	PSD								0		0
		PSD-P								0		0
		Wr								99		94
	Walleye	PSD										100
		PSD-P										100
		Wr										101
boat shocker	Largemouth Bass	PSD	46	72	78	65	66	53	83		55	
(night)	Ü	PSD-P	18	47	19	11	16	14	19		13	
		Wr	125	75	111	103	105	104	101		105	
	Smallmouth Bass	PSD	0	33	56	71	73	100	75		17	
		PSD-P	0	22	19	14	35	100	0		0	
		Wr		66	100	93	90	60	88		86	
	Walleye	PSD					0		0		100	
	•	PSD-P					0		0		100	
		Wr					104		102		100	
frame net (std	Black Bullhead	PSD	0			79					98	100
3/4 in)	DIACK DUIIIICAU	PSD-P	0			0					90	8
		Wr	84			88					91	88
		VVI	64			Øδ					91	ŏŏ

							Υe	ear				
Gear	Species	Index	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
frame net (std	Black Crappie	PSD	69			15					92	65
3/4 in)		PSD-P	2			5					68	12
		Wr	109			102					87	94
	Bluegill	PSD	93			44					80	61
		PSD-P	23			16					16	11
		Wr	111			103					96	99
	Green Sunfish	PSD									100	100
		PSD-P									33	0
		Wr									111	111
	Largemouth Bass	PSD	100									
		PSD-P	0									
		Wr	110									
	Northern Pike	PSD	40			50					60	40
		PSD-P	0			17					0	0
		Wr	81			96					93	87
	Smallmouth Bass	PSD	67			80						
		PSD-P	33			20						
		Wr	95			90						
	Walleye	PSD										100
		PSD-P										100
		Wr										84
	White Sucker	PSD	100			100						
		PSD-P	100			100						
		Wr	102			101						
	Yellow Perch	PSD	92			89					20	25
		PSD-P	42			4					0	0
		Wr	78			76					83	87
std exp gill net	Black Bullhead	PSD	0	12								
		PSD-P	0	0								
		Wr	98	104								
	Black Crappie	PSD	0									
		PSD-P	0									
		Wr	96									
	Bluegill	PSD	100	50								
		PSD-P	100	50								
		Wr	102	123								
	Largemouth Bass	PSD	100	100								

							Ye	ar				
Gear	Species	Index	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
std exp gill net	Largemouth Bass	PSD-P	0	100								
		Wr	131	99								
	Northern Pike	PSD	64	78								
		PSD-P	9	0								
		Wr	90	95								
	White Sucker	PSD	100	100								
		PSD-P	100	100								
		Wr	109	104								
	Yellow Perch	PSD	62	60								
		PSD-P	15	5								
		Wr	89	90								

Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Bluegill

				Mean Len	gth (expa	nded sam	ple numbe	er) at cap	ture by ag	е	
Year	N	1	2	3	4	5	6	7	8	9	10+
2015	100		140 (2)	178 (42)	192 (45)	210 (13)	215 (1)				
Species: L	argemou	th Bass									
				Mean Len	gth (expa	nded sam	ple numbe	er) at cap	ture by ag	е	
Year	N	1	2	3	4	5	6	7	8	9	10+
2016	71	366 (71)									
Species: S	mallmou	th Bass									
				Mean Len	gth (expa	nded sam	ple numbe	er) at cap	ture by ag	е	
Year	N	1	2	3	4	5	6	7	8	9	10+
2016	9	302 (9)									
Species: Y	ellow Pe	rch									
				Mean Len	gth (expa	nded sam	ple numbe	er) at cap	ture by ag	e	
Year	N	1	2	3	4	5	6	7	8	9	10+
2016	127			187 (39)	206 (52)	220 (20)	237		245 (4)	234	250 (5)

Fish Condition

Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).


					Length	Group	s		
			S-Q		Q-P		P-M		М
Species	Year	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Black Bullhead Gill Net	2020	1	89	44	95 (1.4)	0		0	
	2021	0		65	94 (1.2)	1	98	0	
	2023	0		37	100 (1.5)	7	91 (3.1)	0	
	2024	0		43	94 (1.7)	7	93 (4.0)	0	
Black Crappie Frame Net	2020	28	99 (1.2)	18	94 (1.3)	1	77	0	
	2023	2	86 (8.3)	6	98 (5.7)	15	87 (1.4)	2	57 (25.5)
	2024	9	100 (3.6)	14	90 (3.2)	3	89 (7.8)	0	
Bluegill Frame Net	2020	32	99 (7.4)	28	98 (1.9)	7	91 (1.2)	0	
	2023	47	102 (1.8)	155	96 (1.0)	39	86 (1.8)	0	
	2024	66	109 (7.7)	84	95 (1.3)	18	84 (2.1)	0	
Largemouth Bass Electro Fishing	2020	28	108 (2.4)	23	102 (2.0)	8	96 (3.3)	0	
	2021	11	103 (2.7)	41	102 (0.9)	12	96 (2.2)	0	
	2022	35	110 (1.2)	44	109 (0.9)	7	99 (3.6)	0	
	2023	21	108 (2.1)	20	106 (1.5)	5	103 (4.7)	1	2
	2024	31	112 (1.3)	26	108 (1.7)	14	105 (4.5)	0	
Northern Pike Gill Net	2020	0		8	91 (1.3)	1	92	0	
	2021	2	88 (10.3)	6	103 (8.6)	4	90 (4.6)	1	
	2023	0		5	91 (2.8)	1	82	1	
	2024	1	84	9	93 (2.3)	4	85 (5.0)	0	
Smallmouth Bass	2020	0		0		1	60	0	
Electro Fishing	2021	1	80	3	90 (5.1)	0		0	
	2022	2	99 (4.4)	0		0		0	

					Length	Group)S		
			S-Q		Q-P		P-M		М
Species	Year	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Smallmouth Bass Electro Fishing	2023	5	87 (3.9)	1	81	0		0	
	2024	4	94 (1.7)	0		0		0	
Walleye Gill Net	2021	1	75	2	94 (8.3)	0		0	
	2023	0		2	89 (1.2)	0		0	
White Sucker Gill Net	2020	0		0		0		3	97 (6.6)
	2021	0		0		0		7	102 (2.2)
	2023	0		0		0		2	100 (0.6)
	2024	0		0		0		4	98 (3.8)
Yellow Perch	2020	1	93	0		0		0	
Gill Net	2021	28	108 (2.2)	0		0		0	
	2023	11	82 (2.2)	2	73 (2.9)	0		0	
	2024	8	88 (2.7)	7	81 (1.7)	0		0	


Length Frequency Distribution

Length frequency histogram of species sampled by year.


Species: Black Bullhead Gear: AFS std gill net




Species: Black Crappie Gear: AFS std frame net




Species: Black Crappie Gear: frame net (std 3/4 in)







Species: Bluegill Gear: AFS std frame net





Species: Bluegill Gear: frame net (std 3/4 in)






Species: Largemouth Bass Gear: boat shocker (day)







Species: Largemouth Bass Gear: boat shocker (night)




Species: Northern Pike Gear: AFS std gill net








Species: Smallmouth Bass Gear: boat shocker (night)




Species: Yellow Perch Gear: AFS std gill net











Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.


Species: Black Bullhead Gear: AFS std gill net




Species: Black Bullhead Gear: std exp gill net




Species: Black Crappie Gear: AFS std frame net



Species: Black Crappie Gear: frame net (std 3/4 in)




Species: Bluegill Gear: AFS std frame net




Species: Bluegill


Gear: frame net (std 3/4 in)




Species: Largemouth Bass Gear: boat shocker (day)




Species: Largemouth Bass Gear: boat shocker (night)




Species: Northern Pike Gear: AFS std gill net




Species: Northern Pike Gear: std exp gill net




Species: Smallmouth Bass Gear: boat shocker (day)




Species: Smallmouth Bass Gear: boat shocker (night)




Species: Walleye Gear: AFS std gill net




Species: White Sucker Gear: AFS std gill net




Species: White Sucker Gear: std exp gill net



Species: Yellow Perch Gear: AFS std gill net



Species: Yellow Perch Gear: std exp gill net



Fish Stocking

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2013	Smallmouth Bass	Fingerling	4,770
2016	Smallmouth Bass	Juvenile	300
2017	Walleye	Large Fingerling	9,500
2019	Walleye	Small Fingerling	12,240
2023	Walleye	Juvenile	10,000
2024	Walleye	Juvenile	30,253