SOUTH DAKOTA STATEWIDE FISHERIES SURVEY Glad Valley Dam, Ziebach County GRA-Lake-271-000 2017

Lake Information

County: Ziebach

Surface Area: 25 Acres

Surveys and Investigations

Survey methods used by gear type, date, and effort.

| Gear | Date | Effort | |
|-------------------|---------------|--------------|--|
| AFS std frame net | July 18, 2017 | 5 net-nights | |
| AFS std frame net | July 19, 2017 | 5 net-nights | |
| AFS std gill net | July 18, 2017 | 2 net-nights | |
| AFS std gill net | July 19, 2017 | 2 net-nights | |

Common Fish Species Present

Yellow Perch

Black Bullhead

Green Sunfish

Largemouth Bass

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

$$CPUE = \frac{number \ off ish}{effort}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{number \, offish \ge quality \, length}{number \, of \, fish \ge stock \, length}\right) \ge 100$$

$$PSD - P = \left(\frac{number \ offish \ge preferred \ length}{number \ of \ fish \ge stock \ length}\right) \ge 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{Ws}\right) \ge 100$$

Confidence intervals (**CI**) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (**S**), quality (**Q**), preferred (**P**), memorable (**M**) and trophy (**T**). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

| | St | Stock | | Quality | | Preferred | | Memorable | | ophy |
|------------------|------|-------|------|---------|------|-----------|------|-----------|------|------|
| Species Name | (in) | (cm) | (in) | (cm) | (in) | (cm) | (in) | (cm) | (in) | (cm) |
| Bigmouth Buffalo | 11 | 28 | 18 | 46 | 24 | 61 | 30 | 76 | 37 | 94 |
| Black Bullhead | 6 | 15 | 9 | 23 | 12 | 30 | 15 | 38 | 18 | 46 |
| Black Crappie | 5 | 13 | 8 | 20 | 10 | 25 | 12 | 30 | 15 | 38 |

7/16/2018 Page 3

| | Stock | | Qu | ality | Pref | erred | Mem | orable | Trophy | |
|----------------------------------|-------|------|------|-------|------|-------|------|--------|--------|------|
| Species Name | (in) | (cm) | (in) | (cm) | (in) | (cm) | (in) | (cm) | (in) | (cm) |
| Blue Catfish | 12 | 30 | 20 | 51 | 30 | 76 | 35 | 89 | 45 | 114 |
| Bluegill | 3 | 8 | 6 | 15 | 8 | 20 | 10 | 25 | 12 | 30 |
| Bluegill X Gr. Sunfish Hybrid | 3 | 8 | 6 | 15 | 8 | 20 | 10 | 25 | 12 | 30 |
| Brown Bullhead | 5 | 13 | 8 | 20 | 11 | 28 | 14 | 36 | 17 | 43 |
| Burbot | 8 | 20 | 15 | 38 | 21 | 53 | 26 | 67 | 32 | 82 |
| Channel Catfish | 11 | 28 | 16 | 41 | 24 | 61 | 28 | 71 | 36 | 91 |
| Common Carp | 11 | 28 | 16 | 41 | 21 | 53 | 26 | 66 | 33 | 84 |
| Flathead Catfish | 14 | 35 | 20 | 51 | 28 | 71 | 34 | 86 | 40 | 102 |
| Freshwater Drum | 8 | 20 | 12 | 30 | 15 | 38 | 20 | 51 | 25 | 63 |
| Gizzard Shad | 7 | 18 | 11 | 28 | | | | | | |
| Green Sunfish | 3 | 8 | 6 | 15 | 8 | 20 | 10 | 25 | 12 | 30 |
| Lake Herring | 5 | 13 | 8 | 20 | 11 | 28 | 14 | 35 | 17 | 43 |
| Largemouth Bass | 8 | 20 | 12 | 30 | 15 | 38 | 20 | 51 | 25 | 63 |
| Longnose Gar | 16 | 41 | 27 | 69 | 36 | 91 | 45 | 114 | 55 | 140 |
| Muskellunge | 20 | 51 | 30 | 76 | 38 | 97 | 42 | 107 | 50 | 127 |
| Northern Pike | 14 | 35 | 21 | 53 | 28 | 71 | 34 | 86 | 44 | 112 |
| Paddlefish | 16 | 41 | 26 | 66 | 33 | 84 | 41 | 104 | 51 | 130 |
| Pumpkinseed | 3 | 8 | 6 | 15 | 8 | 20 | 10 | 25 | 12 | 30 |
| Redear Sunfish | 4 | 10 | 7 | 18 | 9 | 23 | 11 | 28 | 13 | 33 |
| River Carpsucker | 7 | 18 | 11 | 28 | 14 | 36 | 18 | 46 | 22 | 56 |
| Rock Bass | 4 | 10 | 7 | 18 | 9 | 23 | 11 | 28 | 13 | 33 |
| Rudd | 6 | 15 | 10 | 25 | 12 | 30 | 15 | 38 | 19 | 48 |
| Sauger | 8 | 20 | 12 | 30 | 15 | 38 | 20 | 51 | 25 | 63 |
| Saugeye | 9 | 23 | 14 | 35 | 18 | 46 | 22 | 56 | 27 | 69 |
| Shorthead Redhorse | 6 | 15 | 10 | 25 | 13 | 33 | 16 | 41 | 20 | 51 |
| Smallmouth Bass | 7 | 18 | 11 | 28 | 14 | 35 | 17 | 43 | 20 | 51 |
| Smallmouth Buffalo | 11 | 28 | 18 | 46 | 24 | 61 | 30 | 76 | 37 | 94 |
| Spotted Bass | 7 | 18 | 11 | 28 | 14 | 35 | 17 | 43 | 20 | 51 |
| Striped Bass | 12 | 30 | 20 | 51 | 30 | 76 | 35 | 89 | 45 | 114 |
| Striped Bass Hybrid (wiper) | 8 | 20 | 12 | 30 | 15 | 38 | 20 | 51 | 25 | 63 |
| Walleye | 10 | 25 | 15 | 38 | 20 | 51 | 25 | 63 | 30 | 76 |
| White Bass | 6 | 15 | 9 | 23 | 12 | 30 | 15 | 38 | 18 | 46 |
| White Crappie | 5 | 13 | 8 | 20 | 10 | 25 | 12 | 30 | 15 | 38 |
| White Perch | 5 | 13 | 8 | 20 | 10 | 25 | 12 | 30 | 15 | 38 |
| White Sucker | 6 | 15 | 10 | 25 | 13 | 33 | 16 | 41 | 20 | 51 |
| Yellow Bass | 4 | 10 | 7 | 18 | 9 | 23 | 11 | 28 | 13 | 33 |
| Yellow Bullhead | 4 | 10 | 7 | 18 | 9 | 23 | 11 | 28 | 14 | 36 |
| Yellow Perch | 5 | 13 | 8 | 20 | 10 | 25 | 12 | 30 | 15 | 38 |

Catch Summary of Stock Length Fish

| | | Abur | idance | Stock Density Indices | | | | | ndition |
|-------------------|-----------------|------|--------|-----------------------|-------|-------|-------|-----|---------|
| Gear | Species | CPUE | CI-80 | PSD | CI-80 | PSD-P | CI-80 | Wr | CI-80 |
| AFS std frame net | Black Bullhead | 5.5 | 3.2 | 98 | | 62 | 10 | 99 | 2 |
| | Green Sunfish | 1.2 | 1.0 | 100 |) | 58 | 24 | 129 | 3 |
| | Largemouth Bass | 0.4 | 0.2 | 50 |) | 0 | | 98 | 2 |
| | Yellow Perch | 1.2 | 0.8 | 92 | | 42 | 24 | 105 | 3 |
| AFS std gill net | Black Bullhead | 4.5 | 4.7 | 100 |) | 83 | | 108 | 4 |
| | Largemouth Bass | 0.5 | 0.8 | 0 | | 0 | | 111 | 1 |
| | Yellow Perch | 13.0 | 9.7 | 98 | ł | 83 | 8 | 101 | 3 |

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

10-Year Catch Per Unit Effort by Gear and Species

| | | | | | | | CPUE | | | | | |
|------------------|-----------------|------|------|------|------|------|------|------|------|------|------|------|
| Gear | Species | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | Avg |
| AFS std frame | Black Bullhead | | | | | | | | | | 5.5 | 5.5 |
| net | Green Sunfish | | | | | | | | | | 1.2 | 1.2 |
| | Largemouth Bass | | | | | | | | | | 0.4 | 0.4 |
| | Yellow Perch | | | | | | | | | | 1.2 | 1.2 |
| AFS std gill net | Black Bullhead | | | | | | | | | | 4.5 | 4.5 |
| | Largemouth Bass | | | | | | | | | | 0.5 | 0.5 |
| | Yellow Perch | | | | | | | | | | 13.0 | 13.0 |
| rod and reel | Largemouth Bass | | | | | | | | 1.8 | | | 1.8 |

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

| | | Year |
|------------------|--------------|---|
| Gear | Species | Index 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017 |
| AFS std frame | Yellow Perch | PSD 92 |
| net | | PSD-P 42 |
| | | Wr 105 |
| AFS std gill net | Yellow Perch | PSD 98 |
| | | PSD-P 83 |
| | | Wr 101 |

Back-Calculated Lengths

Mean species back-calculated total length (mm) at age, standard error (SE), and sample size (N).

| | - | | | | | | | d length (S | | | | |
|------------------|-----|----|--------------|---------------|---------------|---------------|---------------|---------------|-----|----|----|----|
| Year Class | Age | Ν | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| 2016 | 1 | 1 | 76 | | | | | | | | | |
| 2015 | 2 | 5 | 89 (1.9) | 177 (2) | | | | | | | | |
| 2015 | 2 | 5 | 102 (5.6) | 192 (9.2) | | | | | | | | |
| 2014 | 3 | 3 | 108 (.9) | 185 (10.8) | 242 (10.8) | | | | | | | |
| 2014 | 3 | 6 | 107 (3.4) | 184 (9.9) | 230 (13.8) | | | | | | | |
| 2012 | 5 | 15 | 86 (3.4) | 155 (3.4) | 209 (2.7) | 251 (3) | 283 (4) | | | | | |
| 2011 | 6 | 3 | 77 (2.5) | 132 (5.6) | 191 (7) | 229 (10.5) | 263 (13.9) | 278 (12.7) | | | | |
| 2011 | 6 | 9 | 79 (3.6) | 143 (6.1) | 203 (7.7) | 242 (8.5) | 272 (9.2) | 292 (10.1) | | | | |
| 2010 | 7 | 1 | 80 | 146 | 197 | 224 | 239 | 255 | 269 | | | |
| Weighted Mean | | 48 | 90 | 163 | 212 | 245 | 276 | 286 | 269 | | | |
| Year Class | Age | Ν | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 |
| 2016 | 1 | 1 | | | | | | | | | | |
| 2015 | 2 | 5 | | | | | | | | | | |
| 2015 | 2 | 5 | | | | | | | | | | |
| 2014 | 3 | 3 | | | | | | | | | | |
| 2014 | 3 | 6 | | | | | | | | | | |
| 2012 | 5 | 15 | | | | | | | | | | |
| 2011 | 6 | 3 | | | | | | | | | | |
| 2011 | 6 | 9 | | | | | | | | | | |
| 2010 | 7 | 1 | | | | | | | | | | |
| Weighted Mean | | 48 | | | | | | | | | | |

Species: Yellow Perch

Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Yellow Perch

| | Mean Length (expanded sample number) at capture by age | | | | | | | | | | | | |
|------|--|---|------------|------------|---|-------------|-------------|------------|---|---|-----|--|--|
| Year | N | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10+ | | |
| 2017 | 52 | | 216 (6) | 249 (7) | | 297 (22) | 298 (14) | 273 (2) | | | | | |

Fish Condition


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

| | | | Length Groups | | | | | | | | | | |
|--------------------------|------|---|---------------|---|--------------|----|--------------|----|-------------|--|--|--|--|
| | | | S-Q | | Q-P | | P-M | М | | | | | |
| Species | Year | Ν | Wr (SE) | Ν | Wr (SE) | Ν | Wr (SE) | Ν | Wr (SE) | | | | |
| Yellow Perch Gill Net | 2017 | 1 | 93 | 8 | 113 (9.7) | 26 | 103 (2.0) | 17 | 92 (1.4) | | | | |

Length Frequency Distribution

Length frequency histogram of species sampled by year.

Species: Yellow Perch Gear: AFS std gill net



Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.

Species: Yellow Perch Gear: Gill Net

