SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Coal Springs, Perkins County LMO-Lake-1689-000 2017

Lake Information

Name: Coal Springs

County: Perkins

Surface Area: 91 Acres

Surveys and Investigations

Survey methods used by gear type, date, and effort.

| Gear | Date | Effort |
|-------------------|---------------|--------------|
| AFS std frame net | July 18, 2017 | 5 net-nights |
| AFS std frame net | July 19, 2017 | 5 net-nights |
| AFS std gill net | July 18, 2017 | 2 net-nights |
| AFS std gill net | July 19, 2017 | 2 net-nights |

Common Fish Species Present

Bluegill

Black Bullhead

Northern Pike

Walleye

Yellow Perch

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

$$\mathit{CPUE} = \frac{\mathit{number of fish}}{\mathit{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$\textit{PSD} = \left(\frac{number\ of\ fish \geq quality\ length}{number\ of\ fish \geq stock\ length}\right) \ge 100$$

$$\textit{PSD} - \textit{P} = \left(\frac{number\ of\ fish\ \geq preferred\ length}{number\ of\ fish\ \geq stock\ length}\right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{Ws}\right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (**S**), quality (**Q**), preferred (**P**), memorable (**M**) and trophy (**T**). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

| | St | ock | Quality Preferred | | erred | Memorable | | Tro | ophy | |
|------------------|------|------|-------------------|------|-------|-----------|------|------|------|------|
| Species Name | (in) | (cm) | (in) | (cm) | (in) | (cm) | (in) | (cm) | (in) | (cm) |
| Bigmouth Buffalo | 11 | 28 | 18 | 46 | 24 | 61 | 30 | 76 | 37 | 94 |
| Black Bullhead | 6 | 15 | 9 | 23 | 12 | 30 | 15 | 38 | 18 | 46 |
| Black Crappie | 5 | 13 | 8 | 20 | 10 | 25 | 12 | 30 | 15 | 38 |

7/16/2018 Page 3

| | St | ock | Qu | ality | Preferred | | Mem | orable | Trophy | |
|----------------------------------|------|------|------|-------|-----------|------|------|--------|--------|------|
| Species Name | (in) | (cm) | (in) | (cm) | (in) | (cm) | (in) | (cm) | (in) | (cm) |
| Blue Catfish | 12 | 30 | 20 | 51 | 30 | 76 | 35 | 89 | 45 | 114 |
| Bluegill | 3 | 8 | 6 | 15 | 8 | 20 | 10 | 25 | 12 | 30 |
| Bluegill X Gr. Sunfish Hybrid | 3 | 8 | 6 | 15 | 8 | 20 | 10 | 25 | 12 | 30 |
| Brown Bullhead | 5 | 13 | 8 | 20 | 11 | 28 | 14 | 36 | 17 | 43 |
| Burbot | 8 | 20 | 15 | 38 | 21 | 53 | 26 | 67 | 32 | 82 |
| Channel Catfish | 11 | 28 | 16 | 41 | 24 | 61 | 28 | 71 | 36 | 91 |
| Common Carp | 11 | 28 | 16 | 41 | 21 | 53 | 26 | 66 | 33 | 84 |
| Flathead Catfish | 14 | 35 | 20 | 51 | 28 | 71 | 34 | 86 | 40 | 102 |
| Freshwater Drum | 8 | 20 | 12 | 30 | 15 | 38 | 20 | 51 | 25 | 63 |
| Gizzard Shad | 7 | 18 | 11 | 28 | | | | | | |
| Green Sunfish | 3 | 8 | 6 | 15 | 8 | 20 | 10 | 25 | 12 | 30 |
| Lake Herring | 5 | 13 | 8 | 20 | 11 | 28 | 14 | 35 | 17 | 43 |
| Largemouth Bass | 8 | 20 | 12 | 30 | 15 | 38 | 20 | 51 | 25 | 63 |
| Longnose Gar | 16 | 41 | 27 | 69 | 36 | 91 | 45 | 114 | 55 | 140 |
| Muskellunge | 20 | 51 | 30 | 76 | 38 | 97 | 42 | 107 | 50 | 127 |
| Northern Pike | 14 | 35 | 21 | 53 | 28 | 71 | 34 | 86 | 44 | 112 |
| Paddlefish | 16 | 41 | 26 | 66 | 33 | 84 | 41 | 104 | 51 | 130 |
| Pumpkinseed | 3 | 8 | 6 | 15 | 8 | 20 | 10 | 25 | 12 | 30 |
| Redear Sunfish | 4 | 10 | 7 | 18 | 9 | 23 | 11 | 28 | 13 | 33 |
| River Carpsucker | 7 | 18 | 11 | 28 | 14 | 36 | 18 | 46 | 22 | 56 |
| Rock Bass | 4 | 10 | 7 | 18 | 9 | 23 | 11 | 28 | 13 | 33 |
| Rudd | 6 | 15 | 10 | 25 | 12 | 30 | 15 | 38 | 19 | 48 |
| Sauger | 8 | 20 | 12 | 30 | 15 | 38 | 20 | 51 | 25 | 63 |
| Saugeye | 9 | 23 | 14 | 35 | 18 | 46 | 22 | 56 | 27 | 69 |
| Shorthead Redhorse | 6 | 15 | 10 | 25 | 13 | 33 | 16 | 41 | 20 | 51 |
| Smallmouth Bass | 7 | 18 | 11 | 28 | 14 | 35 | 17 | 43 | 20 | 51 |
| Smallmouth Buffalo | 11 | 28 | 18 | 46 | 24 | 61 | 30 | 76 | 37 | 94 |
| Spotted Bass | 7 | 18 | 11 | 28 | 14 | 35 | 17 | 43 | 20 | 51 |
| Striped Bass | 12 | 30 | 20 | 51 | 30 | 76 | 35 | 89 | 45 | 114 |
| Striped Bass Hybrid (wiper) | 8 | 20 | 12 | 30 | 15 | 38 | 20 | 51 | 25 | 63 |
| Walleye | 10 | 25 | 15 | 38 | 20 | 51 | 25 | 63 | 30 | 76 |
| White Bass | 6 | 15 | 9 | 23 | 12 | 30 | 15 | 38 | 18 | 46 |
| White Crappie | 5 | 13 | 8 | 20 | 10 | 25 | 12 | 30 | 15 | 38 |
| White Perch | 5 | 13 | 8 | 20 | 10 | 25 | 12 | 30 | 15 | 38 |
| White Sucker | 6 | 15 | 10 | 25 | 13 | 33 | 16 | 41 | 20 | 51 |
| Yellow Bass | 4 | 10 | 7 | 18 | 9 | 23 | 11 | 28 | 13 | 33 |
| Yellow Bullhead | 4 | 10 | 7 | 18 | 9 | 23 | 11 | 28 | 14 | 36 |
| Yellow Perch | 5 | 13 | 8 | 20 | 10 | 25 | 12 | 30 | 15 | 38 |

Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

| | | Abundance | | St | ock De | es | Coi | ndition | |
|-------------------|----------------|-----------|-------|-----|--------|-------|-------|---------|-------|
| Gear | Species | CPUE | CI-80 | PSD | CI-80 | PSD-P | CI-80 | Wr | CI-80 |
| AFS std frame net | Black Bullhead | 32.4 | 17.9 | 43 | 4 | 0 | | 75 | 2 |
| | Bluegill | 4.0 | 2.5 | 38 | 12 | 10 | | 126 | 3 |
| | Northern Pike | 2.4 | 0.7 | 92 | | 4 | | 61 | 3 |
| AFS std gill net | Black Bullhead | 3.5 | 1.4 | 64 | 22 | 0 | | 86 | 5 |
| | Northern Pike | 5.8 | 3 2.0 | 100 | | 0 | | 70 | 1 |
| | Walleye | 0.5 | 0.5 | 100 | | 100 | | 106 | 2 |
| | Yellow Perch | 0.3 | 0.4 | 0 | | 0 | | 93 | |

10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

| | | | | | | | CPUE | | | | | |
|------------------|----------------|------|------|------|------|------|------|------|-------|------|------|------|
| Gear | Species | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | Avg |
| AFS std frame | Black Bullhead | | | | | | | | ' | | 32.4 | 32.4 |
| net | Bluegill | | | | | | | | | | 4.0 | 4.0 |
| | Northern Pike | | | | | | | | | | 2.4 | 2.4 |
| AFS std gill net | Black Bullhead | | | | | | | | | | 3.5 | 3.5 |
| | Northern Pike | | | | | | | | | | 5.8 | 5.8 |
| | Walleye | | | | | | | | | | 0.5 | 0.5 |
| | Yellow Perch | | | | | | | | | | 0.3 | 0.3 |
| frame net (1/2 | Black Bullhead | | | 0.3 | | | | | | | | 0.3 |
| inch) | Northern Pike | | | 7.8 | | | | | | | | 7.8 |
| | Yellow Perch | | | 0.0 | | | | | | | | 0.0 |
| frame net (std | Black Bullhead | | | | 0.4 | | 40.3 | 37.6 | 142.3 | | | 55.2 |
| 3/4 in) | Bluegill | | | | | | 8.0 | 2.0 | 22.8 | | | 8.5 |
| | Northern Pike | | | | 3.8 | | 1.8 | 8.0 | 4.5 | | | 4.5 |
| | Yellow Perch | | | | 0.9 | | 0.3 | 0.6 | 0.4 | | | 0.6 |
| std exp gill net | Northern Pike | | | 4.0 | | | | | | | | 4.0 |
| std exp gill net | Black Bullhead | | | | | | 26.0 | 46.0 | 31.0 | | | 34.3 |
| (150 ft) | Northern Pike | | | | 15.0 | | 27.0 | 3.5 | 6.5 | | | 13.0 |
| | Yellow Perch | | | | 3.5 | | 2.0 | 4.5 | 1.0 | | | 2.8 |

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

| | | | | Year | | | | | | | | |
|------------------|---------------|-------|------|------|------|------|------|------|------|------|------|------|
| Gear | Species | Index | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 |
| AFS std frame | Northern Pike | PSD | | | | | | | | | | 92 |
| net | | PSD-P | | | | | | | | | | 4 |
| | | Wr | | | | | | | | | | 61 |
| AFS std gill net | Northern Pike | PSD | | | | | | | | | | 100 |
| | | PSD-P | | | | | | | | | | 0 |
| | | Wr | | | | | | | | | | 70 |
| | Walleye | PSD | | | | | | | | | | 100 |
| | | PSD-P | | | | | | | | | | 100 |
| | | Wr | | | | | | | | | | 106 |
| | Yellow Perch | PSD | | | | | | | | | | 0 |
| | | PSD-P | | | | | | | | | | 0 |
| | | Wr | | | | | | | | | | 93 |
| frame net (1/2 | Northern Pike | PSD | | | 3 | | | | | | | |
| inch) | | PSD-P | | | 0 | | | | | | | |
| | | Wr | | | 80 | | | | | | | |
| | Yellow Perch | PSD | | | 0 | | | | | | | |
| | | PSD-P | | | 0 | | | | | | | |
| frame net (std | Northern Pike | PSD | | | | 7 | | 43 | 23 | 69 | | |
| 3/4 in) | | PSD-P | | | | 3 | | 14 | 0 | 0 | | |
| | | Wr | | | | 82 | | 77 | 66 | 75 | | |
| | Yellow Perch | PSD | | | | 86 | | 100 | 100 | 100 | | |
| | | PSD-P | | | | 29 | | 100 | 100 | 67 | | |
| | | Wr | | | | 100 | | 102 | 90 | 87 | | |
| std exp gill net | Northern Pike | PSD | | | 0 | | | | | | | |
| | | PSD-P | | | 0 | | | | | | | |
| | | Wr | | | 81 | | | | | | | |
| std exp gill net | Northern Pike | PSD | | | | 10 | | 20 | 43 | 92 | | |
| (150 ft) | | PSD-P | | | | 3 | | 4 | 0 | 8 | | |
| | | Wr | | | | 90 | | 81 | 70 | 77 | | |
| | Yellow Perch | PSD | | | | 86 | | 100 | 78 | 0 | | |

7/16/2018 Page 7

| | Year | | | | | | | | | | | |
|------------------------------|--------------|-------|------|------|------|------|------|------|------|------|------|------|
| Gear | Species | Index | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 |
| std exp gill net (150 ft) | Yellow Perch | PSD-P | | | | 57 | | 100 | 33 | 0 | | |
| | | Wr | | | | 110 | | 113 | 87 | 90 | | |

Back-Calculated Lengths

Mean species back-calculated total length (mm) at age, standard error (SE), and sample size (N).

Species: Walleye

| | | | | | Me | an back-d | calculated | d length (| SE) at ag | е | | |
|------------------|-----|---|-----|-----|-----|-----------|------------|------------|-----------|-----|----|----|
| Year Class | Age | N | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| 2010 | 7 | 1 | 161 | 246 | 377 | 424 | 450 | 484 | 500 | | | |
| 2009 | 8 | 1 | 154 | 230 | 300 | 333 | 426 | 497 | 526 | 540 | | |
| Weighted Mean | | 2 | 158 | 238 | 339 | 379 | 438 | 491 | 513 | 540 | | |
| Year Class | Age | N | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 |
| 2010 | 7 | 1 | | | | | | | | | | |
| 2009 | 8 | 1 | | | | | | | | | | |
| Weighted Mean | | 2 | | | | | | | | | | |

Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Walleye

| 1 | • | | | | | | | | | | |
|------------|-----------|-----|-------------|----------|------------|------------|----------|------------|-------------|-------------|-----|
| | | | N | Mean Ler | ngth (expa | nded sam | ole numb | er) at cap | ture by age |) | |
| Year | N | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10+ |
| 2017 | 2 | | | | | | | 510 (1) | 549 (1) | | |
| Species: Y | ellow Per | rch | <u> </u> | Mean Ler | ooth (expa | nded sami | ole numb | er) at can | ture by age | <u> </u> | |
| Year | N | | 2 | 3 | | 5 | 6 | 7 | 8 | 9 | 10+ |
| ı eai | | | | <u></u> | 4 | | | | | | |
| 2014 | 18 | | 200 (10) | | 261 (6) | 289 (2) | | | | | |

Fish Condition


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

| | | Length Groups | | | | | | | | |
|---------------------------|------|---------------|-------------|----|-------------|---|--------------|---|--------------|--|
| | | - | S-Q | | Q-P | | P-M | | М | |
| Species | Year | N | Wr (SE) | N | Wr (SE) | N | Wr (SE) | N | Wr (SE) | |
| Northern Pike Gill Net | 2013 | 86 | 81 (0.6) | 18 | 76 (2.8) | 4 | 94 (3.1) | 0 | | |
| | 2014 | 8 | 70 (1.5) | 6 | 71 (0.6) | 0 | | 0 | | |
| | 2015 | 2 | 87 (0.0) | 22 | 75 (0.8) | 0 | | 2 | 100 (0.0) | |
| | 2017 | 0 | | 23 | 70 (1.1) | 0 | | 0 | | |
| Walleye Gill Net | 2017 | 0 | | 0 | | 2 | 106 (1.8) | 0 | | |
| Yellow Perch Gill Net | 2013 | 0 | | 0 | | 8 | 113 (2.7) | 0 | | |
| | 2014 | 4 | 86 (0.0) | 8 | 85 (2.4) | 6 | 90 (4.5) | 0 | | |
| | 2015 | 4 | 90 (0.4) | 0 | | 0 | | 0 | | |
| | 2017 | 1 | 93 | 0 | | 0 | | 0 | | |


Length Frequency Distribution


Length frequency histogram of species sampled by year.


Species: Northern Pike Gear: AFS std gill net




Species: Northern Pike Gear: std exp gill net (150 ft)









Species: Walleye Gear: AFS std gill net





Species: Yellow Perch Gear: AFS std gill net



Species: Yellow Perch Gear: std exp gill net (150 ft)









Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.


Species: Northern Pike Gear: Gill Net



Species: Walleye Gear: Gill Net



Species: Yellow Perch Gear: Gill Net



Fish Stocking

Number of fish stocked by year, species, and size.

| Year | Species | Size | Number |
|------|-----------------|------------------|--------|
| 2009 | Walleye | Fingerling | 40,100 |
| 2009 | Yellow Perch | Adult | 1,000 |
| 2010 | Largemouth Bass | Fingerling | 5,000 |
| 2010 | Walleye | Small Fingerling | 8,200 |
| 2011 | Bluegill | Adult | 125 |
| 2011 | Golden Shiner | Adult | 60 |
| 2011 | Yellow Perch | Adult | 585 |
| 2012 | Bluegill | Adult | 470 |
| 2012 | Largemouth Bass | Adult | 150 |
| 2012 | Yellow Perch | Adult | 572 |
| 2014 | Walleye | Fingerling | 20,000 |
| 2014 | Walleye | Large Fingerling | 1,592 |
| | | | |