

SOUTH DAKOTA STATEWIDE FISHERIES SURVEY
Campbell, Brookings County
MBS-Lake-234-000
2017

Lake Information

Name:	Campbell	Maximum Depth:	8 Feet
County:	Brookings	Mean Depth:	3 Feet
Legal Description:	T109n-R50W-Sec.28, 29, 32, 33; T108N-R50W-Sec. 5	OHWM Elevation:	1,576
Surface Area:	798 Acres	Outlet Elevation:	1,575

Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
AFS std gill net	June 29, 2017	6 net-nights

Common Fish Species Present

Yellow Perch

Walleye

Black Bullhead

White Sucker

Common Carp

Channel Catfish

White Bass

Bigmouth Buffalo

Saugeye

Northern Pike

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

$$CPUE = \frac{\text{number of fish}}{\text{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{\text{number of fish} \geq \text{quality length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

$$PSD - P = \left(\frac{\text{number of fish} \geq \text{preferred length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{W_s} \right) \times 100$$

Confidence intervals (**CI**) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (**S**), quality (**Q**), preferred (**P**), memorable (**M**) and trophy (**T**). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Bigmouth Buffalo	11	28	18	46	24	61	30	76	37	94
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Blue Catfish	12	30	20	51	30	76	35	89	45	114
Bluegill	3	8	6	15	8	20	10	25	12	30
Bluegill X Gr. Sunfish Hybrid	3	8	6	15	8	20	10	25	12	30
Brown Bullhead	5	13	8	20	11	28	14	36	17	43
Burbot	8	20	15	38	21	53	26	67	32	82
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Common Carp	11	28	16	41	21	53	26	66	33	84
Flathead Catfish	14	35	20	51	28	71	34	86	40	102
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Gizzard Shad	7	18	11	28						
Green Sunfish	3	8	6	15	8	20	10	25	12	30
Lake Herring	5	13	8	20	11	28	14	35	17	43
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Longnose Gar	16	41	27	69	36	91	45	114	55	140
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Paddlefish	16	41	26	66	33	84	41	104	51	130
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Redear Sunfish	4	10	7	18	9	23	11	28	13	33
River Carpsucker	7	18	11	28	14	36	18	46	22	56
Rock Bass	4	10	7	18	9	23	11	28	13	33
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Saugeye	9	23	14	35	18	46	22	56	27	69
Shorthead Redhorse	6	15	10	25	13	33	16	41	20	51
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Smallmouth Buffalo	11	28	18	46	24	61	30	76	37	94
Spotted Bass	7	18	11	28	14	35	17	43	20	51
Striped Bass	12	30	20	51	30	76	35	89	45	114
Striped Bass Hybrid (wiper)	8	20	12	30	15	38	20	51	25	63
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
White Perch	5	13	8	20	10	25	12	30	15	38
White Sucker	6	15	10	25	13	33	16	41	20	51
Yellow Bass	4	10	7	18	9	23	11	28	13	33
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

Gear	Species	Abundance		Stock Density Indices			Condition		
		CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
AFS std gill net	Bigmouth Buffalo	1.5	1.7	0		0			
	Black Bullhead	10.8	2.3	29	8	3			
	Channel Catfish	2.3	1.2	50	22	14		98	3
	Common Carp	4.3	1.5	85		58	15		
	Northern Pike	0.8	1.0	80		40		85	6
	Saugeye	1.3	0.7	0		0		88	3
	Shorthead Redhorse	0.2	0.2	100		100			
	Walleye	10.8	2.7	20	7	0		82	1
	White Bass	1.5	0.9	100		100		92	3
	White Sucker	9.5	2.2	100		98			
Yellow Perch	3.3	1.0	95		80		91	3	

10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

Gear	Species	CPUE										Avg
		2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	
AFS std gill net	Bigmouth Buffalo										1.5	1.5
	Black Bullhead										10.8	10.8
	Channel Catfish										2.3	2.3
	Common Carp										4.3	4.3
	Northern Pike										0.8	0.8
	Saugeye										1.3	1.3
	Shorthead Redhorse										0.2	0.2
	Walleye										10.8	10.8
	White Bass										1.5	1.5
	White Sucker										9.5	9.5
	Yellow Perch									3.3	3.3	
large frame net	Bigmouth Buffalo	3.0		34.9	11.4		1.2					12.6
	Black Bullhead	139.5		158.3	322.9		32.3					163.3
	Black Crappie			0.4	0.1							0.3
	Channel Catfish	0.1		2.5	8.6		6.1					4.3
	Common Carp	17.0		6.1	1.7		8.2					8.3
	Green Sunfish	0.4										0.4
	Northern Pike	2.6		1.5	2.9		0.2					1.8
	Orangespotted Sunfish	0.0										0.0
	Shorthead Redhorse				0.1		0.1					0.1
	Walleye	0.9		0.3	0.3							0.5
	White Bass				0.5							0.5
	White Sucker	4.1		6.9	3.0		0.8					3.7
	Yellow Bullhead	1.3		0.3	2.3							1.3
	Yellow Perch	1.8		4.0	0.6							2.1
std exp gill net	Bigmouth Buffalo			0.7				0.7	2.0			1.1
	Black Bullhead	36.0		41.0	26.0		21.3	27.7	39.7	61.0		36.1
	Channel Catfish			3.3	4.3		7.3	3.7	3.0	5.3		4.5
	Common Carp	16.3			0.3		1.3	0.0	4.0	1.3		3.9
	Common Shiner	0.0							0.0			0.0
	Emerald Shiner	0.0										0.0
	Northern Pike	2.0		1.7	8.0		1.0	8.7	2.7	1.3		3.6
	Orangespotted Sunfish	0.0		0.0					0.0	0.0		0.0

Gear	Species	CPUE										
		2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	Avg
std exp gill net	Shorthead Redhorse						1.0			0.3		0.7
	Walleye	15.0		1.3	0.7		0.3	3.0	0.3	22.7		6.2
	White Bass				1.0				2.7	2.0		1.9
	White Sucker	16.3		41.3	5.7		2.3	3.3	7.7	9.3		12.3
	Yellow Bullhead	0.3			0.3							0.3
	Yellow Perch	5.3		146.7	5.3			2.3	26.3	6.0		32.0

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

Gear	Species	Index	Year											
			2008	2009	2010	2011	2012	2013	2014	2015	2016	2017		
AFS std gill net	Northern Pike	PSD											80	
		PSD-P											40	
		Wr											85	
	Walleye	PSD												20
		PSD-P												0
		Wr												82
	Yellow Perch	PSD												95
		PSD-P												80
		Wr												91
large frame net	Black Crappie	PSD			0	100								
		PSD-P			0	0								
		Wr			105	115								
	Northern Pike	PSD	88		100	55			100					
		PSD-P	27		53	7			0					
		Wr	95		95	89			72					
	Walleye	PSD	56		100	100								
		PSD-P	33		33	0								
		Wr	93		106	97								
	Yellow Perch	PSD	22		48	50								
		PSD-P	22		38	0								
		Wr	102		95	85								
	std exp gill net	Northern Pike	PSD	83		40	42		100	35	75	100		
			PSD-P	17		0	4		0	8	0	25		
			Wr	89		98	91		79	98	97	88		
Walleye		PSD	47		100	50		0	100	0	0			
		PSD-P	7		0	0		0	11	0	0			
		Wr	99		105	97		100	105	113	90			
Yellow Perch		PSD	56		25	31			0	100	100			
		PSD-P	44		17	6			0	43	83			
		Wr	106		98	97			112	103	91			

Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Walleye

		Mean Length (expanded sample number) at capture by age									
Year	N	1	2	3	4	5	6	7	8	9	10+
2017	66	270 (28)	376 (37)					505 (1)			

Fish Condition

Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).


Species	Year	Length Groups							
		S-Q		Q-P		P-M		M	
		N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Northern Pike Gill Net	2013	0		3	79 (10.4)	0		0	
	2014	17	99 (2.7)	7	95 (2.3)	2	93 (4.1)	0	
	2015	2	95 (4.2)	6	98 (3.6)	0		0	
	2016	0		3	91 (0.1)	1	79	0	
	2017	1	96	2	74 (4.2)	2	90 (0.8)	0	
Walleye Gill Net	2013	1	100	0		0		0	
	2014	0		8	104 (2.4)	1	108	0	
	2015	1	113	0		0		0	
	2016	68	90 (0.7)	0		0		0	
	2017	52	82 (0.8)	13	84 (1.6)	0		0	
Yellow Perch Gill Net	2014	7	112 (3.5)	0		0		0	
	2015	0		45	103 (1.1)	34	103 (1.0)	0	
	2016	0		3	87 (3.7)	12	91 (2.5)	3	92 (3.9)
	2017	1	81	3	104 (2.5)	9	90 (2.2)	7	88 (4.8)

Length Frequency Distribution

Length frequency histogram of species sampled by year.




Species: Northern Pike


Gear: AFS std gill net



Species: Northern Pike


Gear: std exp gill net






2016

Species: Walleye
Gear: AFS std gill net




2017


Species: Walleye
Gear: std exp gill net




2013



2014




2015




2016

Species: Yellow Perch
Gear: AFS std gill net





2017

Species: Yellow Perch
Gear: std exp gill net




2014




Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.


Species: Black Crappie
Gear: Frame Net




Species: Northern Pike
Gear: Gill Net



Species: Walleye
Gear: Gill Net



Species: Yellow Perch
Gear: Gill Net



Fish Stocking

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2006	Walleye	Fry	926,316
2009	Yellow Perch	Fry	4,584,000
2010	Walleye	Small Fingerling	91,320
2014	Walleye	Fry	553,320
2015	Walleye	Fry	450,000
2016	Saugeye	Small Fingerling	46,310
2017	Saugeye	Small Fingerling	62,500
2017	Yellow Perch	Small Fingerling	510,590