

SOUTH DAKOTA STATEWIDE FISHERIES SURVEY
Goldsmith, Brookings County
MBS-Lake-236-000
2017

Lake Information

Name:	Goldsmith	Maximum Depth:	9 Feet
County:	Brookings	Mean Depth:	6 Feet
Legal Description:	T110N-R51W-Sec 9,16		
Surface Area:	308 Acres		

Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
AFS std gill net	June 28, 2017	6 net-nights

Common Fish Species Present

Yellow Perch

Walleye

Northern Pike

White Sucker

Black Bullhead

Saugeye

Common Carp

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

$$CPUE = \frac{\text{number of fish}}{\text{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{\text{number of fish} \geq \text{quality length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

$$PSD - P = \left(\frac{\text{number of fish} \geq \text{preferred length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{W_s} \right) \times 100$$

Confidence intervals (**CI**) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (**S**), quality (**Q**), preferred (**P**), memorable (**M**) and trophy (**T**). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Bigmouth Buffalo	11	28	18	46	24	61	30	76	37	94
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Blue Catfish	12	30	20	51	30	76	35	89	45	114
Bluegill	3	8	6	15	8	20	10	25	12	30
Bluegill X Gr. Sunfish Hybrid	3	8	6	15	8	20	10	25	12	30
Brown Bullhead	5	13	8	20	11	28	14	36	17	43
Burbot	8	20	15	38	21	53	26	67	32	82
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Common Carp	11	28	16	41	21	53	26	66	33	84
Flathead Catfish	14	35	20	51	28	71	34	86	40	102
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Gizzard Shad	7	18	11	28						
Green Sunfish	3	8	6	15	8	20	10	25	12	30
Lake Herring	5	13	8	20	11	28	14	35	17	43
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Longnose Gar	16	41	27	69	36	91	45	114	55	140
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Paddlefish	16	41	26	66	33	84	41	104	51	130
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Redear Sunfish	4	10	7	18	9	23	11	28	13	33
River Carpsucker	7	18	11	28	14	36	18	46	22	56
Rock Bass	4	10	7	18	9	23	11	28	13	33
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Saugeye	9	23	14	35	18	46	22	56	27	69
Shorthead Redhorse	6	15	10	25	13	33	16	41	20	51
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Smallmouth Buffalo	11	28	18	46	24	61	30	76	37	94
Spotted Bass	7	18	11	28	14	35	17	43	20	51
Striped Bass	12	30	20	51	30	76	35	89	45	114
Striped Bass Hybrid (wiper)	8	20	12	30	15	38	20	51	25	63
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
White Perch	5	13	8	20	10	25	12	30	15	38
White Sucker	6	15	10	25	13	33	16	41	20	51
Yellow Bass	4	10	7	18	9	23	11	28	13	33
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

Gear	Species	Abundance		Stock Density Indices			Condition		
		CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
AFS std gill net	Black Bullhead	1.0	0.5	0		0			
	Common Carp	0.2	0.2	100		100			
	Northern Pike	2.0	0.9	92		8		88	3
	Saugeye	0.7	0.5	0		0		99	5
	Walleye	2.0	0.7	17		0		89	2
	White Sucker	1.7	1.1	100		80			
	Yellow Perch	5.0	2.5	67	13	50	14	111	2

10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

Gear	Species	CPUE										Avg
		2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	
AFS std gill net	Black Bullhead										1.0	1.0
	Common Carp										0.2	0.2
	Northern Pike										2.0	2.0
	Saugeye										0.7	0.7
	Walleye										2.0	2.0
	White Sucker										1.7	1.7
	Yellow Perch										5.0	5.0
large frame net	Bigmouth Buffalo		3.6		0.2		6.6					3.5
	Black Bullhead		3.6		7.2		10.8					7.2
	Bluegill						0.4					0.4
	Common Carp		2.0		0.8		1.2					1.3
	Northern Pike		0.4		2.2		1.4					1.3
	Walleye		2.0		3.8		3.4					3.1
	White Bass		3.8		2.4		1.2					2.5
	White Crappie		0.6				0.6					0.6
	White Sucker		1.2		0.4		0.2					0.6
	Yellow Bullhead		0.6									0.6
	Yellow Perch		0.2									0.2
std exp gill net	Bigmouth Buffalo						0.0	0.0				0.0
	Black Bullhead				0.7		1.0	7.7	9.3			4.7
	Common Carp		0.3		0.3		0.7	0.3	0.3	0.3		0.4
	Northern Pike		1.0		3.7		3.0	1.0	1.3	2.3		2.1
	Orangespotted Sunfish							0.0				0.0
	Saugeye									0.0		0.0
	Walleye		2.7		3.7		5.3		0.0	6.3		3.6
	White Bass		2.0									2.0
	White Sucker		2.3		1.3		2.7	4.0	1.0	4.0		2.6
Yellow Perch		4.7		4.3		1.3	5.0	7.0	40.3		10.4	

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

Gear	Species	Index	Year											
			2008	2009	2010	2011	2012	2013	2014	2015	2016	2017		
AFS std gill net	Northern Pike	PSD											92	
		PSD-P											8	
		Wr											88	
	Walleye	PSD												17
		PSD-P												0
		Wr												89
	Yellow Perch	PSD												67
		PSD-P												50
		Wr												111
large frame net	Northern Pike	PSD		50		73		86						
		PSD-P		0		18		0						
		Wr		89		85		88						
	Walleye	PSD		90		37		41						
		PSD-P		30		11		12						
		Wr		84		85		100						
	Yellow Perch	PSD		100										
		PSD-P		0										
		Wr		105										
std exp gill net	Northern Pike	PSD		33		36		89	0	50	86			
		PSD-P		0		9		33	0	0	29			
		Wr		67		84		97	94	77	84			
	Walleye	PSD		88		64		31		0	0			
		PSD-P		0		0		6		0	0			
		Wr		92		93		94			96			
	Yellow Perch	PSD		57		23		50	60	43	75			
		PSD-P		0		0		0	0	33	8			
		Wr		104		106		111	106	105	107			

Fish Condition


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

Species	Year	Length Groups							
		S-Q		Q-P		P-M		M	
		N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Northern Pike Gill Net	2013	1	76	5	89 (4.0)	3	117 (4.8)	0	
	2014	3	94 (6.3)	0		0		0	
	2015	2	74 (2.1)	2	79 (1.4)	0		0	
	2016	1	83	4	85 (2.9)	2	82 (0.5)	0	
	2017	1	107	10	86 (2.3)	1	93	0	
Walleye Gill Net	2013	11	95 (1.1)	4	89 (2.0)	0		1	100
	2015	0		0		0		0	
	2016	19	96 (1.6)	0		0		0	
	2017	10	89 (1.8)	2	90 (2.3)	0		0	
Yellow Perch Gill Net	2013	2	123 (1.1)	2	99 (4.9)	0		0	
	2014	6	107 (2.8)	9	106 (2.6)	0		0	
	2015	12	105 (2.5)	2	100 (3.2)	7	106 (3.4)	0	
	2016	30	106 (1.8)	81	109 (0.7)	9	102 (1.8)	1	92
	2017	10	106 (3.8)	5	117 (4.6)	14	112 (1.9)	1	100




Length Frequency Distribution


Length frequency histogram of species sampled by year.

Species: Northern Pike
Gear: AFS std gill net




Species: Northern Pike
Gear: std exp gill net








Species: Walleye
Gear: AFS std gill net




Species: Walleye
Gear: std exp gill net









Species: Yellow Perch
Gear: AFS std gill net



Species: Yellow Perch
Gear: std exp gill net







Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.


Species: Northern Pike
Gear: Gill Net



Species: Walleye
Gear: Gill Net



Species: Yellow Perch
Gear: Gill Net



Fish Stocking

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2006	Walleye	Small Fingerling	31,200
2008	Walleye	Small Fingerling	30,000
2009	Walleye	Fingerling	2,991
2009	Walleye	Large Fingerling	846
2010	Walleye	Small Fingerling	32,640
2011	Walleye	Large Fingerling	172
2011	Yellow Perch	Adult	2,280
2011	Yellow Perch	Small Fingerling	145,920
2012	Walleye	Juvenile	1,350
2014	Walleye	Fry	300,000
2015	Walleye	Small Fingerling	20,480
2016	Saugeye	Small Fingerling	31,030