#### SOUTH DAKOTA STATEWIDE FISHERIES SURVEY

Nine Mile, Marshall County UJA-Lake-882-001 2017

#### **Lake Information**

Name: Nine Mile Maximum Depth: 10 Feet

County: Mean Depth: 7 Feet

**OHWM Elevation:** 1,826

Surface Area: 248 Acres Outlet Elevation: 1,825

#### **Surveys and Investigations**

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
AFS std gill net	May 31, 2017	3 net-nights
AFS std gill net	June 01, 2017	3 net-nights

# **Common Fish Species Present**

Yellow Perch Walleye

Northern Pike

Black Bullhead

Bluegill

Black Crappie

#### **Terminology**

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

$$\mathit{CPUE} = \frac{\mathit{number of fish}}{\mathit{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$\textit{PSD} = \left(\frac{number\ of\ fish \geq quality\ length}{number\ of\ fish \geq stock\ length}\right) \ge 100$$

$$\textit{PSD} - \textit{P} = \left(\frac{number\ of\ fish\ \geq preferred\ length}{number\ of\ fish\ \geq stock\ length}\right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{Ws}\right) \times 100$$

Confidence intervals (CI) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (**S**), quality (**Q**), preferred (**P**), memorable (**M**) and trophy (**T**). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

	St	ock	Qu	Quality Preferred		Memorable		Trophy		
Species Name	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Bigmouth Buffalo	11	28	18	46	24	61	30	76	37	94
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38

7/16/2018 Page 3

	St	ock	Qu	ality	Pref	erred	Mem	orable	Tro	pphy
Species Name	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Blue Catfish	12	30	20	51	30	76	35	89	45	114
Bluegill	3	8	6	15	8	20	10	25	12	30
Bluegill X Gr. Sunfish Hybrid	3	8	6	15	8	20	10	25	12	30
Brown Bullhead	5	13	8	20	11	28	14	36	17	43
Burbot	8	20	15	38	21	53	26	67	32	82
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Common Carp	11	28	16	41	21	53	26	66	33	84
Flathead Catfish	14	35	20	51	28	71	34	86	40	102
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Gizzard Shad	7	18	11	28						
Green Sunfish	3	8	6	15	8	20	10	25	12	30
Lake Herring	5	13	8	20	11	28	14	35	17	43
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Longnose Gar	16	41	27	69	36	91	45	114	55	140
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Paddlefish	16	41	26	66	33	84	41	104	51	130
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Redear Sunfish	4	10	7	18	9	23	11	28	13	33
River Carpsucker	7	18	11	28	14	36	18	46	22	56
Rock Bass	4	10	7	18	9	23	11	28	13	33
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Saugeye	9	23	14	35	18	46	22	56	27	69
Shorthead Redhorse	6	15	10	25	13	33	16	41	20	51
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Smallmouth Buffalo	11	28	18	46	24	61	30	76	37	94
Spotted Bass	7	18	11	28	14	35	17	43	20	51
Striped Bass	12	30	20	51	30	76	35	89	45	114
Striped Bass Hybrid (wiper)	8	20	12	30	15	38	20	51	25	63
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
White Perch	5	13	8	20	10	25	12	30	15	38
White Sucker	6	15	10	25	13	33	16	41	20	51
Yellow Bass	4	10	7	18	9	23	11	28	13	33
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

### **Catch Summary of Stock Length Fish**

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

			dance	Sto	ock De	Co	ndition		
Gear	Species	CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
AFS std gill net	Black Bullhead	2.0	0.5	0		0		92	4
	Black Crappie	0.3	0.3	0		0		108	7
	Bluegill	0.5	0.3	67		0		111	2
	Northern Pike	2.5	0.7	80		0		84	3
	Walleye	0.2	0.2	0		0		95	
	Yellow Perch	6.0	3.3	0		0		92	2

# 10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

							CPUE					
Gear	Species	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	Avg
AFS std gill net	Black Bullhead										2.0	2.0
	Black Crappie										0.3	0.3
	Bluegill										0.5	0.5
	Northern Pike										2.5	2.5
	Walleye										0.2	0.2
	Yellow Perch										6.0	6.0
frame net (std	Black Bullhead					91.6						91.6
3/4 in)	Northern Pike					0.7						0.7
	Walleye					0.3						0.3
	White Sucker					0.1						0.1
	Yellow Perch					44.9						44.9
std exp gill net	Black Bullhead					29.7						29.7
	Northern Pike					7.3						7.3
	White Sucker					1.7						1.7
	Yellow Perch					32.3						32.3

### 10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

							Υe	ar				
Gear	Species	Index	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017
AFS std gill net	Black Crappie	PSD										0
		PSD-P										0
		Wr										108
	Northern Pike	PSD										80
		PSD-P										0
		Wr										84
	Walleye	PSD										0
		PSD-P										0
		Wr										95
	Yellow Perch	PSD										0
		PSD-P										0
		Wr										92
frame net (std	Northern Pike	PSD					75					
3/4 in)		PSD-P					0					
		Wr					95					
	Walleye	PSD					100					
		PSD-P					75					
		Wr					95					
	Yellow Perch	PSD					1					
		PSD-P					0					
		Wr					100					
std exp gill net	Northern Pike	PSD					41					
		PSD-P					9					
		Wr					96					
	Yellow Perch	PSD					0					
		PSD-P					0					
		Wr					99					

# **Length at Capture**

Mean length at capture by age across years sampled, sample size (N).

Species: Walleye

Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+
2017	1				335 (1)						
Species: Y	ellow Pe	rch									
Magazil angth (avanaded comple number) at conture by one											

	Mean Length (expanded sample number) at capture by age											
Year	N	1	2	3	4	5	6	7	8	9	10+	
2017	36		150 (14)	178 (23)								
2012	111	91 (14)		148 (96)		185 (1)						

### **Fish Condition**


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

			Length Groups										
			S-Q	Q-P		P-M			M				
Species	Year	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)				
Northern Pike Gill Net	2017	3	84 (5.7)	12	85 (2.8)	0		0					
Walleye Gill Net	2017	1	95	0		0		0					
Yellow Perch Gill Net	2017	36	92 (1.1)	0		0		0					


### **Length Frequency Distribution**

Length frequency histogram of species sampled by year.


Species: Northern Pike Gear: AFS std gill net



Species: Walleye Gear: AFS std gill net




Species: Yellow Perch Gear: AFS std gill net




### **Historic Fish Sizes and Relative Abundance**

Size distribution per net by color for species sampled by year.


Species: Northern Pike Gear: Gill Net



Species: Walleye Gear: Gill Net



Species: Yellow Perch Gear: Gill Net



# Fish Stocking

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2010	Walleye	Fry	260,000
2012	Walleye	Fry	260,000
2013	Walleye	Fry	1,300,000
2014	Yellow Perch	Small	5,000