

SOUTH DAKOTA STATEWIDE FISHERIES SURVEY
Marindahl, Yankton County
VER-Lake-276-000
2017

Lake Information

Name:	Marindahl	Maximum Depth:	30 Feet
County:	Yankton	Mean Depth:	13 Feet
Legal Description:	T95N-R54W-Sec. 7, 17, 18, 20		
Surface Area:	147 Acres		

Surveys and Investigations

Survey methods used by gear type, date, and effort.

Gear	Date	Effort
AFS std frame net	August 07, 2017	5 net-nights
AFS std gill net	August 07, 2017	6 net-nights

Common Fish Species Present

Largemouth Bass

Bluegill

Black Crappie

Black Bullhead

White Sucker

Yellow Perch

Channel Catfish

Gizzard Shad

Common Carp

Smallmouth Bass

Terminology

Catch per unit effort (**CPUE**) refers to the relative abundance of a species. It is defined as the number of fish captured per unit of effort (i.e., number of fish captured per net-night or number of fish captured per hour electrofishing). In this report CPUE is typically given for only stock-length fish (see length categories table for stock lengths).

$$CPUE = \frac{\text{number of fish}}{\text{effort}}$$

Population size structure is quantified using the indices proportional size distribution of quality-length fish (**PSD**) and proportional size distribution of preferred-length fish (**PSD-P**). These indices indicate the proportion of stock-length fish that are equal to or greater than a given length. Minimum lengths for stock, quality and preferred length fish are given in the length categories table.

$$PSD = \left(\frac{\text{number of fish} \geq \text{quality length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

$$PSD - P = \left(\frac{\text{number of fish} \geq \text{preferred length}}{\text{number of fish} \geq \text{stock length}} \right) \times 100$$

Relative weight (**Wr**) is used to quantify fish plumpness. Relative weight is the ratio of what a fish weighs (W) compared to a length-specific standard weight (Ws) multiplied by 100. Relative weight values of 95-105 are commonly cited as optimum values, but values in the 80s are common during summer sampling in South Dakota.

$$Wr = \left(\frac{W}{W_s} \right) \times 100$$

Confidence intervals (**CI**) are provided for many of the estimates calculated in this report. The confidence interval provides a range in which the true mean is expected to fall. For example, with an 80% CI we are 80% confident that the interval contains the true value.

Length categories include stock (**S**), quality (**Q**), preferred (**P**), memorable (**M**) and trophy (**T**). Length categories for most species have been defined based on a percentage of the world record length for that species. Some species mentioned in this report do not have defined length categories. Length categories for species used in this report are provided in the following table. Measurements are the minimum total length for each category and are reported in inches (in) and centimeters (cm).

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Bigmouth Buffalo	11	28	18	46	24	61	30	76	37	94
Black Bullhead	6	15	9	23	12	30	15	38	18	46
Black Crappie	5	13	8	20	10	25	12	30	15	38

Species Name	Stock		Quality		Preferred		Memorable		Trophy	
	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)	(in)	(cm)
Blue Catfish	12	30	20	51	30	76	35	89	45	114
Bluegill	3	8	6	15	8	20	10	25	12	30
Bluegill X Gr. Sunfish Hybrid	3	8	6	15	8	20	10	25	12	30
Brown Bullhead	5	13	8	20	11	28	14	36	17	43
Burbot	8	20	15	38	21	53	26	67	32	82
Channel Catfish	11	28	16	41	24	61	28	71	36	91
Common Carp	11	28	16	41	21	53	26	66	33	84
Flathead Catfish	14	35	20	51	28	71	34	86	40	102
Freshwater Drum	8	20	12	30	15	38	20	51	25	63
Gizzard Shad	7	18	11	28						
Green Sunfish	3	8	6	15	8	20	10	25	12	30
Lake Herring	5	13	8	20	11	28	14	35	17	43
Largemouth Bass	8	20	12	30	15	38	20	51	25	63
Longnose Gar	16	41	27	69	36	91	45	114	55	140
Muskellunge	20	51	30	76	38	97	42	107	50	127
Northern Pike	14	35	21	53	28	71	34	86	44	112
Paddlefish	16	41	26	66	33	84	41	104	51	130
Pumpkinseed	3	8	6	15	8	20	10	25	12	30
Redear Sunfish	4	10	7	18	9	23	11	28	13	33
River Carpsucker	7	18	11	28	14	36	18	46	22	56
Rock Bass	4	10	7	18	9	23	11	28	13	33
Rudd	6	15	10	25	12	30	15	38	19	48
Sauger	8	20	12	30	15	38	20	51	25	63
Saugeye	9	23	14	35	18	46	22	56	27	69
Shorthead Redhorse	6	15	10	25	13	33	16	41	20	51
Smallmouth Bass	7	18	11	28	14	35	17	43	20	51
Smallmouth Buffalo	11	28	18	46	24	61	30	76	37	94
Spotted Bass	7	18	11	28	14	35	17	43	20	51
Striped Bass	12	30	20	51	30	76	35	89	45	114
Striped Bass Hybrid (wiper)	8	20	12	30	15	38	20	51	25	63
Walleye	10	25	15	38	20	51	25	63	30	76
White Bass	6	15	9	23	12	30	15	38	18	46
White Crappie	5	13	8	20	10	25	12	30	15	38
White Perch	5	13	8	20	10	25	12	30	15	38
White Sucker	6	15	10	25	13	33	16	41	20	51
Yellow Bass	4	10	7	18	9	23	11	28	13	33
Yellow Bullhead	4	10	7	18	9	23	11	28	14	36
Yellow Perch	5	13	8	20	10	25	12	30	15	38

Catch Summary of Stock Length Fish

Catch per unit effort (CPUE), proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) for species sampled in survey with 80% confidence interval (CI-80).

Gear	Species	Abundance		Stock Density Indices			Condition		
		CPUE	CI-80	PSD	CI-80	PSD-P	CI-80	Wr	CI-80
AFS std frame net	Black Bullhead	41.8	19.5	13	3	1			
	Black Crappie	68.0	40.8	94	2	4	2	94	1
	Common Carp	0.0	0.0	0		0			
	White Sucker	5.6	3.2	96		64	14		
	Yellow Perch	0.2	0.3	100		100		75	
AFS std gill net	Black Bullhead	12.0	3.6	18	7	1			
	Black Crappie	2.8	1.0	94		0		96	1
	Bluegill	0.2	0.2	0		0		85	
	Channel Catfish	3.8	1.5	91		9		103	3
	Common Carp	1.2	1.2	0		0			
	Gizzard Shad	3.3	2.3	90					
	Smallmouth Bass	0.3	0.5	100		0		83	2
	White Sucker	5.0	2.1	90		63	14		
	Yellow Perch	4.2	2.6	96		80	13	82	2

10-Year Catch Per Unit Effort by Gear and Species

Catch per unit effort (CPUE) and average (Avg) of species across 10 years using different gear types.

Gear	Species	CPUE										Avg
		2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	
AFS std frame net	Black Bullhead										41.8	41.8
	Black Crappie										68.0	68.0
	Common Carp										0.0	0.0
	White Sucker										5.6	5.6
	Yellow Perch										0.2	0.2
AFS std gill net	Black Bullhead										12.0	12.0
	Black Crappie										2.8	2.8
	Bluegill										0.2	0.2
	Channel Catfish										3.8	3.8
	Common Carp										1.2	1.2
	Gizzard Shad										3.3	3.3
	Smallmouth Bass										0.3	0.3
	White Sucker										5.0	5.0
	Yellow Perch										4.2	4.2
boat shocker (night)	Largemouth Bass		5.5				6.0	8.1	7.5	6.0		6.6
	Smallmouth Bass								0.5			0.5
large frame net	Black Bullhead		0.3		3.9		11.0					5.1
	Black Crappie		70.0		46.1		3.6					39.9
	Bluegill		28.3		81.9		6.3					38.8
	Channel Catfish		0.5		1.8		1.4					1.2
	Common Carp		0.1		0.9		0.0					0.3
	Green Sunfish		0.3		3.5		0.5					1.4
	Largemouth Bass				0.0							0.0
	Sunfish Hybrid						0.0					0.0
	White Sucker		25.4		43.7		4.8					24.6
	Yellow Perch		0.1		0.3							0.2
std frame net (3/8 inch)	Black Bullhead							52.1	185.2	51.3		96.2
	Black Crappie							5.6	19.0	49.4		24.7
	Bluegill							4.7	21.9	17.9		14.8
	Channel Catfish							0.5	1.3	0.6		0.8
	Common Carp							1.4	0.1	0.6		0.7
	Green Sunfish							0.3	1.2	1.2		0.9
	Largemouth Bass								0.0			0.0

		CPUE										
Gear	Species	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	Avg
std frame net	White Sucker							5.5	19.8	12.6		12.6

10-Year Size Structure and Condition Statistics by Gear and Species

Species proportional size distribution (PSD), proportional size distribution of preferred length fish (PSD-P), and relative weight (Wr) collected by different gear types across 10 years.

Gear	Species	Index	Year											
			2008	2009	2010	2011	2012	2013	2014	2015	2016	2017		
AFS std frame net	Black Crappie	PSD											94	
		PSD-P											4	
		Wr											94	
	Yellow Perch	PSD												100
		PSD-P												100
		Wr												75
AFS std gill net	Black Crappie	PSD											94	
		PSD-P											0	
		Wr											96	
	Yellow Perch	PSD												96
		PSD-P												80
		Wr												82
large frame net	Black Crappie	PSD		33		93		47						
		PSD-P		0		0		3						
		Wr		95		98		105						
	Yellow Perch	PSD		100		33								
		PSD-P		0		0								
		Wr		89		67								
std frame net (3/8 inch)	Black Crappie	PSD								98	2	9		
		PSD-P								0	1	7		
		Wr								104	115	99		

Back-Calculated Lengths

Mean species back-calculated total length (mm) at age, standard error (SE), and sample size (N).

Species: Black Crappie

Year Class	Age	N	Mean back-calculated length (SE) at age											
			1	2	3	4	5	6	7	8	9	10		
2015	2	4	81 (.6)	160 (3.1)										
2014	3	9	76 (2.7)	155 (2.9)	202 (4.3)									
2013	4	17	82 (1.8)	164 (1.7)	202 (2.6)	226 (2.6)								
2012	5	1	63	98	172	205	251							
2010	7	2	90 (5.9)	181 (2.4)	224 (10.5)	245 (12.4)	265 (12.7)	279 (7.9)	297 (4.6)					
2009	8	1	74	124	167	221	250	266	278	286				
Weighted Mean		34	80	159	201	227	258	275	291	286				
Year Class	Age	N	11	12	13	14	15	16	17	18	19	20		
2015	2	4												
2014	3	9												
2013	4	17												
2012	5	1												
2010	7	2												
2009	8	1												
Weighted Mean		34												

Length at Capture

Mean length at capture by age across years sampled, sample size (N).

Species: Black Crappie

Year	N	Mean Length (expanded sample number) at capture by age									
		1	2	3	4	5	6	7	8	9	10+
2017	340		174 (16)	213 (60)	226 (249)	256 (4)		302 (8)	294 (4)		
2015	191	147 (188)	216 (1)	241 (1)	250 (1)						
2014	56		176 (1)	227 (54)	216 (1)						
2013	36		198 (30)	204 (5)	261 (1)						
2011	461	167 (23)		215 (438)							
2009	700	156 (349)	194 (76)	204 (166)	212 (108)						

Fish Condition


Mean relative weight (Wr) by sample size (N), length category stock to quality (S-Q), quality to preferred (Q-P), preferred to memorable (P-M), and memorable (M) for species collected across survey years with standard error (SE).

Species	Year	Length Groups							
		S-Q		Q-P		P-M		M	
		N	Wr (SE)	N	Wr (SE)	N	Wr (SE)	N	Wr (SE)
Black Crappie Frame Net	2013	19	107 (0.8)	16	104 (3.6)	1	103	0	
	2014	1	58	55	105 (0.9)	0		0	
	2015	187	117 (2.3)	2	89 (1.7)	1	84	0	
	2016	449	101 (0.8)	10	83	35	81 (1.0)	0	
	2017	19	98 (1.3)	306	94 (0.5)	11	93 (1.5)	4	93
Yellow Perch Gill Net	2017	1	89	4	89 (4.0)	16	83 (1.5)	4	74 (2.6)


Length Frequency Distribution

Length frequency histogram of species sampled by year.



Species: Black Crappie
Gear: AFS std frame net




Species: Black Crappie
Gear: large frame net




Species: Black Crappie
Gear: std frame net (3/8 inch)






Species: Yellow Perch
 Gear: AFS std gill net




Historic Fish Sizes and Relative Abundance

Size distribution per net by color for species sampled by year.

Species: Black Crappie
Gear: Frame Net



Species: Yellow Perch
Gear: Gill Net



Fish Stocking

Number of fish stocked by year, species, and size.

Year	Species	Size	Number
2006	Largemouth Bass	Adult	320
2009	Largemouth Bass	Juvenile	2,025
2011	Largemouth Bass	Fingerling	2,880
2013	Largemouth Bass	Large Fingerling	3,104
2013	Rainbow Trout	Fingerling	3,424
2014	Bluegill	Adult	144
2014	Channel Catfish	Adult	3
2015	Gizzard Shad	Adult	74
2015	Largemouth Bass	Juvenile	1,590
2016	Gizzard Shad	Adult	360
2017	Walleye	Fingerling	1,200
2017	Walleye	Juvenile	225
2017	Yellow Perch	Adult	7,437